

5% Dextrose Injection, USP

2.5 grams per 50 mL
(50 mg per mL)

For intravenous use.

Rx only

Each 50 mL contains: Dextrose monohydrate, 2.5 g; water for injection, 50 mL.

252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5).

Single Dose Only. Discard Unused Portion.

Dextrose solutions without salts should not be used in blood transfusions because of possible Rouleau formation. Additives may be incompatible. Consult with pharmacist, if available. When introducing additives, use aseptic technique, mix thoroughly and do not store.

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Protect from freezing.

The container closure is not made with natural rubber latex.

Non-PVC,
Non-DEHP,
Sterile.



(01)00365219456056



EXP: MMYYYY

LOT 14XXZZZZ

Sample Label. Please see package insert for complete prescribing information.



**FRESENIUS
KABI**

Lake Zurich, IL 60047

Made in USA

www.fresenius-kabi.com/us

403954

**Sample label. Please see package insert
for complete prescribing information.**

5% Dextrose Injection, USP NDC 63323-624-05

**2.5 grams per 50 mL
(50 mg per mL)**

For intravenous use.

Rx only

Each 50 mL contains: Dextrose monohydrate, 2.5 g;
water for injection, 50 mL.

252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5).

Single Dose Only. Discard Unused Portion.

Dextrose solutions without salts should not be used in
blood transfusions because of possible Rouleau formation.

Additive compatibility, consult pharmacist. When
introducing additives, use aseptic technique, mix
thoroughly and do not store.

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

**STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled
Room Temperature]. Protect from freezing.**

The container closure is not made with natural rubber
latex. Non-PVC, Non-DEHP, Sterile.



Mfd. for:



**FRESENIUS
KABI**

Lake Zurich, IL 60047

Made in Norway

www.fresenius-kabi.com/us

LOT

EXP

403407

FUE 0076 01-69-04-005

1 2 3 4 5 6 7 8 9 0

freeflex®

NDC 65219-464-05

100 mL

5% Dextrose Injection, USP

5 grams per 100 mL
(50 mg per mL)

For intravenous use.

Rx only

Each 100 mL contains: Dextrose monohydrate, 5 g; water for injection, 100 mL.

252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5).

Single Dose Only. Discard Unused Portion.

Dextrose solutions without salts should not be used in blood transfusions because of possible Rouleau formation. Additives may be incompatible. Consult with pharmacist, if available. When introducing additives, use aseptic technique, mix thoroughly and do not store.

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Protect from freezing.

The container closure is not made with natural rubber latex.

Non-PVC,
Non-DEHP,
Sterile.



(01)00365219464051



EXP: MMYYYY

LOT 14XXZZZZ

Sample Label. Please see package insert for complete prescribing information.



FRESENIUS
KABI

Lake Zurich, IL 60047

Made in USA

www.fresenius-kabi.com/us

403953

5% Dextrose Injection, USP

NDC 63323-624-06

5 grams per 100 mL

(50 mg per mL)

For intravenous use.

Rx only

Each 100 mL contains: Dextrose monohydrate, 5 g;
water for injection, 100 mL.

252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5).

Single Dose Only. Discard Unused Portion.

Dextrose solutions without salts should not be used in blood transfusions because of possible Rouleau formation. Additive compatibility, consult pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store.

Usual dosage: See package insert. The overwrap is a moisture barrier.
Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex.

Non-PVC, Non-DEHP, Sterile.

(01)00363323624064



XXXX XX



EXP: MM - YYYY

LOT: 0000000

Mfd. for:

 FRESENIUS
KABI

Lake Zurich, IL 60047

Made in Norway

www.fresenius-kabi.com/us

Sample Label. Please see
package insert for complete
prescribing information.

1 2 3 4 5 6 7 8 9 0

403428B

FUH 0075 01-62-04-001A

freeflex[®] NDC 65219-458-05 **250 mL**

5% Dextrose Injection, USP

12.5 grams per 250 mL
(50 mg per mL)

For intravenous use.

Rx only

50

Each 100 mL contains: Dextrose monohydrate, 5 g; water for injection, 100 mL.
252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5).
Single Dose Only. Discard Unused Portion.

Dextrose solutions without salts should not be used in blood transfusions because of possible Rouleau formation. Additives may be incompatible. Consult with pharmacist, if available. When introducing additives, use aseptic technique, mix thoroughly and do not store.

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

200

R2 →



EXP: MMYYYY

LOT: 14XXZZZZ



**FRESENIUS
KABI**

Lake Zurich, IL 60047
Made in USA

www.fresenius-kabi.com/us

Sample Label.
Please see package
insert for complete
prescribing information.

mL

403952

5% Dextrose Injection, USP

**12.5 grams per 250 mL
(50 mg per mL)**

50

For intravenous use. Rx only

Each 100 mL contains: Dextrose monohydrate, 5 g; water for injection, 100 mL. 252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5). Single Dose Only. Discard Unused Portion.

100

Dextrose solutions without salts should not be used in blood transfusions because of possible Rouleau formation. Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store.

150

Usual dosage: See package insert.

The overwrap is a moisture barrier.
Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

200

(01)00363323624040



XXXXXX



EXP: MM - YYYY

LOT: 0000000

Mfd. for:

FRESENIUS
KABI

Lake Zurich, IL 60047

Made in Norway

www.fresenius-kabi.com/us

403427B

Sample Label. Please see
package insert for complete
prescribing information.

1 2 3 4 5 6 7 8 9 0

FUH 0072 01-62-04-002A

freeflex[®]

NDC 65219-460-05

500 mL

5% Dextrose Injection, USP

25 grams per 500 mL

(50 mg per mL)

100

For intravenous use.

Rx only

Each 100 mL contains: Dextrose monohydrate, 5 g; water for injection, 100 mL.

252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5).

200

Single Dose Only. Discard Unused Portion.

Dextrose solutions without salts should not be used in blood transfusions because of possible Rouleau formation. Additives may be incompatible. Consult with pharmacist, if available. When introducing additives, use aseptic technique, mix thoroughly and do not store.

300

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

400



(01)00365219460053

mL



EXP: MMYYYY

LOT: 14XXZZZZ



**FRESENIUS
KABI**

Lake Zurich, IL 60047
Made in USA

www.fresenius-kabi.com/us

**Sample Label.
Please see package
insert for complete
prescribing information.**

403950

5% Dextrose Injection, USP

25 grams per 500 mL

(50 mg per mL)

100 For intravenous use.

Rx only

Each 100 mL contains: Dextrose monohydrate, 5 g; water for injection, 100 mL.

252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5).

Single Dose Only. Discard Unused Portion.

200 Dextrose solutions without salts should not be used in blood transfusions because of possible Rouleau formation.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store.

Usual dosage: See package insert.


The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

300 **STORE AT:** 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex.

Non-PVC, Non-DEHP, Sterile.



(01)00363323624071

400 Mfd. for:



**FRESENIUS
KABI**

Lake Zurich, IL 60047

Made in Germany

www.fresenius-kabi.com/us

403213B

LOT

EXP

**Sample label.
Please see
package insert
for complete
prescribing
information.**

100

eeflex®

NDC 65219-462-05

1,000 mL

5% Dextrose Injection, USP

200

50 grams per 1,000 mL

(50 mg per mL)

For intravenous use.

Rx only

300

Each 100 mL contains: Dextrose monohydrate, 5 g; water for injection, 100 mL.

252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5).

Single Dose Only. Discard Unused Portion.

400

Dextrose solutions without salts should not be used in blood transfusions because of possible Rouleau formation. Additives may be incompatible. Consult with pharmacist, if available. When introducing additives, use aseptic technique, mix thoroughly and do not store.

500

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

600



(01)00365219462057

700



EXP: MMYYYY

LOT: YYMULLXXX

800



Lake Zurich, IL 60047
Made in USA

www.fresenius-kabi.com/us

Sample Label. Please see package insert for complete prescribing information.

900

403949A

100

5% Dextrose Injection, USP

200**50 grams per 1,000 mL**

(50 mg per mL)

For intravenous use.

Rx only

300**Each 100 mL contains:** Dextrose monohydrate, 5 g; water for injection, 100 mL.

252 mOsmol/LITER (CALC.) pH 4.3 (3.2 to 6.5).

Single Dose Only. Discard Unused Portion.

Dextrose solutions without salts should not be used in blood transfusions because of possible Rouleau formation.

400

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store.


Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

500**STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from freezing.**

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

600

(01)00363323624033

700

Mfd. for:

**FRESENIUS
KABI**

Lake Zurich, IL 60047

Made in Germany

www.fresenius-kabi.com/us

403209A

LOT

EXP

**Sample label.
Please see
package insert
for complete
prescribing
information.**

900

0717151/01 US