

100

freeflex[®]

NDC 65219-479-05

1,000 mL

Lactated Ringer's Injection, USP

For intravenous use.

Rx only

200

Each 100 mL contains:

		mEq/L:	
Sodium Chloride, USP	600 mg	Na ⁺	130
Sodium Lactate, USP	310 mg	K ⁺	4
Potassium Chloride, USP	30 mg	Ca ²⁺	2.7
Calcium Chloride, USP	20 mg	Cl ⁻	109
		Lactate	28

300

273 mOsmol/L (calc.) pH 6.5 (6.0 to 7.5)

Single Dose Only. Discard Unused Portion.

Not for use in the treatment of lactic acidosis.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

400

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

500

STORE AT: 20° to 25° C (68° to 77°F) [see USP Controlled Room Temperature]. Avoid excessive heat.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

600



(01)00365219479055



EXP: MMYYYY

LOT: YYMULLXXX

700

**FRESENIUS
KABI**
Lake Zurich, IL 60047
Made in USA

www.fresenius-kabi.com/us

Sample Label.
Please see package insert
for complete prescribing information.

mL

403955

freeflex®

NDC 65219-477-05

500 mL

Lactated Ringer's Injection, USP

For intravenous use.

Rx only

		mEq/L:	
Each 100 mL contains:		Na ⁺	130
Sodium Chloride, USP	600 mg	K ⁺	4
Sodium Lactate, USP	310 mg	Ca ²⁺	2.7
Potassium Chloride, USP	30 mg	Cl ⁻	109
Calcium Chloride, USP	20 mg	Lactate	28
273 mOsmol/L (calc.)		pH 6.5 (6.0 to 7.5)	

Single Dose Only. Discard Unused Portion.

Not for use in the treatment of lactic acidosis.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

STORE AT: 20° to 25° C (68° to 77°F) [see USP Controlled Room Temperature]. Avoid excessive heat.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.



(01)00365219477051



EXP: MMYYYY

LOT: 14XXZZZZ

**FRESENIUS
KABI**Lake Zurich, IL 60047
Made in USA

www.fresenius-kabi.com/us

Sample Label.
Please see package
insert for complete
prescribing information.

403956

free flex[®] NDC 65219-475-05 **250 mL**

Lactated Ringer's Injection, USP

For intravenous use.

Rx only

Each 100 mL contains:		mEq/L:	
Sodium Chloride, USP	600 mg	Na ⁺	130
Sodium Lactate, USP	310 mg	K ⁺	4
Potassium Chloride, USP	30 mg	Ca ²⁺	2.7
Calcium Chloride, USP	20 mg	Cl ⁻	109
		Lactate	28

273 mOsmol/L (calc.) pH 6.5 (6.0 to 7.5)

Single Dose Only. Discard Unused Portion.

Not for use in the treatment of lactic acidosis.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

STORE AT: 20° to 25° C (68° to 77°F) [see USP Controlled Room Temperature]. Avoid excessive heat.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.



(01)00365219475057



EXP: MMYYYY

LOT: 14XXZZZZ



**FRESENIUS
KABI**

Lake Zurich, IL 60047
Made in USA

www.fresenius-kabi.com/us

Sample Label.
Please see package
insert for complete
prescribing information.

403957

50

100

150

200

mL



Sample label. Please see package insert
for complete prescribing information.



free *flex*[®] NDC 63323-164-24 **250 mL**

Lactated Ringer's Injection, USP

For intravenous use.

Rx only

50
100
150
200
mL

Each 100 mL contains:

			mEq/L:	
Sodium Chloride, USP	600 mg	Na ⁺	130	
Sodium Lactate, USP	310 mg	K ⁺	4	
Potassium Chloride, USP	30 mg	Ca ²⁺	2.7	
Calcium Chloride, USP	20 mg	Cl ⁻	109	
		Lactate	28	
273 mOsmol/L (calc.)		pH 6.5 (6.0 to 7.5)		

Single Dose Only. Discard Unused Portion.

Not for use in the treatment of lactic acidosis.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

Usual dosage: See package insert. The overwrap is a moisture barrier. Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Avoid excessive heat.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.



(01)00363323164249

Mfd. for: **LOT**



Lake Zurich, IL 60047 |

Made in Norway

www.fresenius-kabi.com/us

403292

1 2 3 4 5 6 7 8 9 0

FUH 2342 01-62-12-024

