

100

freeflex®

NDC 65219-479-05

1,000 mL

Lactated Ringer's Injection, USP

For intravenous use.

Rx only

200

Each 100 mL contains:

Sodium Chloride, USP	600 mg
Sodium Lactate, USP	310 mg
Potassium Chloride, USP	30 mg
Calcium Chloride, USP	20 mg

mEq/L:

Na ⁺	130
K ⁺	4
Ca ²⁺	2.7
Cl ⁻	109
Lactate	28

300

273 mOsmol/L (calc.) pH 6.5 (6.0 to 7.5)

Single Dose Only. Discard Unused Portion.

Not for use in the treatment of lactic acidosis.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

400

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

500

STORE AT: 20° to 25° C (68° to 77°F) [see USP Controlled Room Temperature]. Avoid excessive heat.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

600



(01)00365219479055



EXP: MMYYYY

LOT: YYMULLXXX

700



**FRESENIUS
KABI**

Lake Zurich, IL 60047
Made in USA

www.fresenius-kabi.com/us

Sample Label.

**Please see package insert
for complete prescribing information.**

mL

403955



Sample label. Please see package insert
for complete prescribing information.



free *flex*[®]

NDC 63323-164-25

500 mL

Lactated Ringer's Injection, USP

100

For intravenous use.

Rx only

200

Each 100 mL contains:		mEq/L:	
Sodium Chloride, USP	600 mg	Na ⁺	130
Sodium Lactate, USP	310 mg	K ⁺	4
Potassium Chloride, USP	30 mg	Ca ²⁺	2.7
Calcium Chloride, USP	20 mg	Cl ⁻	109
		Lactate	28
273 mOsmol/L (calc.)		pH 6.5 (6.0 to 7.5)	

Single Dose Only. Discard Unused Portion.

Not for use in the treatment of lactic acidosis.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

300

Usual dosage: See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Avoid excessive heat.

400

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.



(01)00363323164256

Mfd. for:

LOT



**FRESENIUS
KABI**

EXP

Lake Zurich, IL 60047

Made in Norway

www.fresenius-kabi.com/us

1 2 3 4 5 6 7 8 9 0

mL

403293

FUH 2343 01-62-12-025





Sample label. Please see package insert
for complete prescribing information.



free *flex*[®]

NDC 63323-164-24

250 mL

Lactated Ringer's Injection, USP

For intravenous use.

Rx only

50

Each 100 mL contains:

Sodium Chloride, USP	600 mg	Na ⁺	130
Sodium Lactate, USP	310 mg	K ⁺	4
Potassium Chloride, USP	30 mg	Ca ²⁺	2.7
Calcium Chloride, USP	20 mg	Cl ⁻	109
		Lactate	28
273 mOsmol/L (calc.)		pH 6.5 (6.0 to 7.5)	

Single Dose Only. Discard Unused Portion.

Not for use in the treatment of lactic acidosis.

100

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

150

Usual dosage: See package insert.
The overwrap is a moisture barrier.
Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Avoid excessive heat.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

200



(01)00363323164249

Mfd. for:

LOT



**FRESENIUS
KABI EXP**

Lake Zurich, IL 60047

Made in Norway

www.fresenius-kabi.com/us

1 2 3 4 5 6 7 8 9 0

mL

403292

FUH 2342 01-62-12-024

