Sample label. Please see package insert for complete prescribing information.

 $\bigcirc$ 

free flex®

For intravenous use.

NDC 63323-164-24

250 mL

Rx only

## Lactated Ringer's Injection, USP

			mEq/L:	
Each 100 mL contains:			Na+	130
Sodium Chloride, USP	600	mg	K+	4
Sodium Lactate, USP	310	mg	Ca <sup>2+</sup>	2.7
Potassium Chloride, USP	30	mg	CI-	109
Calcium Chloride, USP	20	mg	Lactate	28
273 mOsmol/L (calc.)	nН	65	(6 0 to 7 5	5)

Single Dose Only. Discard Unused Portion.

## Not for use in the treatment of lactic acidosis.

100

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

150

**Usual dosage:** See package insert. The overwrap is a moisture barrier. Use immediately once removed from overwrap.

**STORE AT: 20°** to **25°**C (**68°** to **77°**F) [see USP Controlled Room Temperature]. Avoid excessive heat.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

200



(01)00363323164249

Mfd. for:

403292

. . . |

FRESENIUS
KABI EXP
Lake Zurich, IL 60047

Made in Norway

1234567890

www.fresenius-kabi.com/us

FUH 2342 01-62-12-024

mL



free flex®

NDC 63323-164-25

500 mL

mFa/I ·

## Lactated Ringer's Injection, USP

100 For intravenous use. Rx only

		IIIE4/ E.	
Each 100 mL contains:		Na <sup>+</sup>	130
Sodium Chloride, USP	600 mg	K <sup>+</sup>	4
Sodium Lactate, USP	310 mg	Ca <sup>2+</sup>	2.7
Potassium Chloride, USP	30 mg	CI-	109
Calcium Chloride, USP	20 mg	Lactate	28
273 mOsmol/L (calc.)	pH 6.5 (6	.0 to 7.5)	

Single Dose Only. Discard Unused Portion.

Not for use in the treatment of lactic acidosis.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

Usual dosage: See package insert. The overwrap is a moisture barrier. Use immediately once removed from overwrap.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperaturel. Avoid excessive heat.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

Mfd. for:

LOT

**FRESENIUS** KABI

**EXP** 

Lake Zurich, IL 60047 Made in Norway www.fresenius-kabi.com/us

1234567890

403293

FUH 2343 01-62-12-025

400

200

300



free flex®

100

200

300

NDC 63323-164-26

1,000 mL

Rx only

## Lactated Ringer's Injection, USP

For intravenous use.

		mEq/L:	
Each 100 mL contains:		Na <sup>+</sup>	130
Sodium Chloride, USP	600 mg	K <sup>+</sup>	4
Sodium Lactate, USP	310 mg	Ca <sup>2+</sup>	2.7
Potassium Chloride, USP	30 mg	CI-	109
Calcium Chloride, USP	20 mg	Lactate	28
273 mOsmol/L (calc.)	pH 6.5 (6	.0 to 7.5)	

Single Dose Only. Discard Unused Portion.

Not for use in the treatment of lactic acidosis.

Additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic technique, mix thoroughly and do not store. Use only if solution is clear and container is undamaged. Must not be used in series connections. Do not administer simultaneously with blood.

**Usual dosage:** See package insert.

The overwrap is a moisture barrier.

Use immediately once removed from overwrap.

**STORE AT: 20°** to **25°**C (**68°** to **77°**F) [see USP Controlled Room Temperature]. Avoid excessive heat.

The container closure is not made with natural rubber latex. Non-PVC, Non-DEHP, Sterile.

600

500



(01)00363323164263

Mfd. for:

FRESENIUS KABI LOT

Lake Zurich, IL 60047 EXP Made in Norway www.fresenius-kabi.com/us

1234567890

mL

403294 FUH 2344 01-62-12-026