HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Imipenem and Cilastatin for Injection, USP (I.V.) safely and effectively. See full prescribing information for Imipenem and Cilastatin for Injection, USP (I.V.).

IMIPENEM AND CILASTATIN FOR INJECTION, USP, for intravenous use Initial U.S. Approval: 1985

RECENT MAJOR CHANGES	
Indications and Usage (1.9)	12/2016
Dosage and Administration (2)	12/2016
INDICATIONS AND USAGE	
Imipenem and Cilastatin for Injection, USP for intravenous use is a combination	on of imipenem.

a penem antibacterial, and cilastatin, a renal dehydropeptidase inhibitor, indicated for the treatment of the following serious infections caused by designated susceptible bacteria: • Lower respiratory tract infections. (1.1)

- Urinary tract infections. (1.2)
- Intra-abdominal infections. (1.3)
- Gynecologic infections. (1.4) • Bacterial septicemia. (1.5)
- Bone and joint infections (1.6)
- Skin and skin structure infections. (1.7) Endocarditis. (1.8)

Limitations of Use:

- Imipenem and Cilastatin for Injection, USP (I.V.) is not indicated in patients with meningitis because safety and efficacy have not been established (1.9).
- Imipenem and Cilastatin for Injection, USP (I.V.) is not recommended in pediatric patients with CNS infections because of the risk of seizures (1.9).
- Impenem and Cilastatin for Injection, USP (I.V.) is not recommended in pediatric patients weighing less than 30 kg with impaired renal function (1.9).

Usage: To reduce the development of drug resistant bacteria and maintain the effectiveness of Imipenem and Cilastatin for Injection, USP (I.V.) and other antibacterial drugs, Imipenem and Cilastatin for Injection, USP (I.V.) should be used only to treat infections that are proven or strongly suspected to be caused by bacteria (1.10).

- The dosage of Imipenem and Cilastatin for Injection (I.V.) in adult patients should be based
- on suspected or confirmed pathogen susceptibility (2.1). For adult patients with normal renal function (creatinine clearance of greater than or equal to 90 mL/min), the recommended dosage regimens are: 500 mg every 6 hours OR 1,000 mg
- every 8 hours OR 1,000 mg every 6 hours (2.1).
- See full prescribing information for dosage recommendations in pediatric patients (2.2).
 A reduction in dose must be made for a patient with a creatinine clearance of less than
- 90 mL/min (2.3). Patients with creatinine clearances of less than 15 mL/min should not receive Imipenem and
- Cilastatin for Injection (I.V.) unless hemodialysis is instituted within 48 hours (2.4). Reconstitute Imipenem and Cilastatin for Injection, USP (I.V.) vial with appropriate diluent
- and dilute the reconstituted suspension with an appropriate infusion solution before administering by intravenous infusion (2.5). ---DOSAGE FORMS AND STRENGTHS----

For Injection: Imipenem and Cilastatin for Injection (I.V.) is a sterile powder mixture for reconstitution in single-dose vials containing: • 250 mg imipenem (anhydrous equivalent) and 250 mg cilastatin sodium (3)

• 500 mg imipenem (anhydrous equivalent) and 500 mg cilastatin sodium (3)

FULL PRESCRIBING INFORMATION: CONTENTS*

- INDICATIONS AND USAGE
- Lower Respiratory Tract Infections
- Urinary Tract Infections (complicated and uncomplicated) 1.2 Intra-Ábdominal Infections
- 1.4 Gynecologic Infections
- 1.5 Bacterial Septicemia
- 1.6 Bone and Joint Infections
- 17 Skin and Skin Structure Infections 1.8 Endocarditis
- 1.9 Limitations of Use
- 1 10 Usade
- 2 DOSAGE AND ADMINISTRATION
- Dosage in Adults 2.2 Dosage in Pediatric Patients
- Dosage in Adult Patients with Renal Impairment 2.3
- 2.4 Dosage in Hemodialysis Patients Reconstitution and Preparation of Imipenem and Cilastatin for Injection (I.V.) 2.5
- Solution for Intravenous Administration Storage of Reconstituted Solutions 2.6
- Incompatibility and Compatibility of Imipenem and Cilastatin for Injection (I.V.) 2.7 with other Antibacterial Drugs
- DOSAGE FORMS AND STRENGTHS CONTRAINDICATIONS
- WARNINGS AND PRECAUTIONS
- Hypersensitivity Reactions
- 5.2 5.3 Seizure Potential
- Increased Seizure Potential Due to Interaction with Valproic Acid 5.4 Clostridium difficile-Associated Diarrhea (CDAD)
- 5.5 Development of Drug-Resistant Bacteria

FULL PRESCRIBING INFORMATION

INDICATIONS AND USAGE

1.1 Lower Respiratory Tract Infections Imipenem and Cilastatin for Injection, USP (I.V.) for intravenous use is indicated for the treatment of lower respiratory tract infections caused by susceptible strains of Staphylococcus aureus (penicillinase-producing isolates), Acinetobacter species, Enterobacter species, Escherichia coli, Haemophilus influenzae, Haemophilus parainfluenzae, Klebsiella species, Serratia

1.2 Urinary Tract Infections (complicated and uncomplicated)

Imipenem and Cilastatin for Injection, USP (I.V.) is indicated for the treatment of urinary tract infections (complicated and uncomplicated) caused by susceptible strains of Enterococcus faecalis, Staphylococcus aureus (penicillinase-producing isolates), Enterobacter species, Escherichia coli, Klebsiella species, Morganella morganii, Proteus vulgaris, Providencia rettgeri Pseudomonas aeruginosa

1.3 Intra-Abdominal Infections

marcescens

Imipenem and Cilastatin for Injection, USP (I.V.) is indicated for the treatment of intraabdominal infections caused by susceptible strains of Enterococcus faecalis, Staphylococcus aureus (penicillinase-producing isolates), Staphylococcus epidermidis, Citrobacter species Enterobacter species, Escherichia coli, Klebsiella species, Morganella morganii, Proteus species, Pseudomonas aeruginosa, Bifidobacterium species, Clostridium species, Eubacterium species, *Peptococus* species, *Peptostreptococus* species, *Propionibacterium* species, *Bacteroides* species including *B. fragilis, Fusobacterium* species.

1.4 Gynecologic Infections

Imipenem and Cilastatin for Injection, USP (I.V.) is indicated for the treatment of gynecologic infections caused by susceptible strains of Enterococcus faecalis, Staphylococcus aureus (penicillinase-producing isolates), Staphylococcus epidermidis, Streptococcus agalactiae (Group B streptococci), Enterobacter species, Escherichia coli, Gardnerella vaginalis, Klebsiella species, Proteus species, Bifidobacterium species, Peptococcus species, Peptostreptococcus species, Propionibacterium species, Bacteroides species including B. fragilis

1.5 Bacterial Septicemia

Imipenem and Cilastatin for Injection, USP (I.V.) is indicated for the treatment of bacterial septicemia caused by susceptible strains of Enterococcus faecalis. Staphylococcus aureus (penicillinase-producing isolates), Enterobacter species, Escherichia coli, Klebsiella species, Pseudomonas aeruginosa, Serratia species, Bacteroides species including B. fragilis.

1.6 Bone and Joint Infections

Imperent and Cilastatin for Injection, USP (I.V.) is indicated for the treatment of bone and joint infections caused by susceptible strains of *Enterococcus faecalis*, *Staphylococcus aureus* (penicillinase-producing isolates), Staphylococcus epidermidis, Enterobacter species, Pseudomonas aeruginosa.

- ---CONTRAINDICATIONS--• Known hypersensitivity to any component of Imipenem and Cilastatin for Injection (I.V.) (4)
- --WARNINGS AND PRECAUTIONS -• Hypersensitivity Reactions: Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients receiving therapy with beta-lactams If an allergic reaction to Imipenem and Cilastatin for Injection (I.V.) occurs, discontinue the drug immediately (5.1).
- Seizure Potential: Seizures and other CNS adverse reactions, such as confusional states and myoclonic activity, have been reported during treatment with Imipenem and Cilastatin for Injection (I.V.). If focal tremors, myoclonus, or seizures occur, patients should be evaluated neurologically, placed on anticonvulsant therapy if not already instituted, and the dosage of Imipenem and Cilastatin for Injection (I.V.) re-examined to determine whether it should be decreased or the antibacterial drug discontinued (5.2).
- Increased Seizure Potential Due to Interaction with Valproic Acid: Co-administration of Imipenem and Cilastatin for Injection (I.V.), to patients receiving valproic acid or divalproex sodium results in a reduction in valproic acid concentrations. The valproic acid concentrations may drop below the therapeutic range as a result of this interaction, therefore increasing the risk of breakthrough seizures. The concomitant use of Imipenem and Cilastatin for Injection
- (I.V.) and valproic acid/divalproex sodium is generally not recommended (5.3, 7.3).
 Clostridium difficile-Associated Diarrhea (CDAD): has been reported with use of Imipenem and Cilastatin for Injection (I.V.) and may range in severity from mild diarrhea to fatal colitis.

Evaluate if diarrhea occurs (5.4). ---ADVERSE REACTIONS---

- The most frequently occurring adverse reactions (≥ 0.2%) in adults were phlebitis, nausea, diarrhea, vomiting, rash, pain injection site, fever, hypotension, seizures, erythema at
- injection site dizziness pruritus vein induration urticaria somnolence (6.1) The most frequently occurring adverse reactions (> 1%) in pediatric patients greater than or equal to 3 months of age were diarrhea, rash, phlebitis, gastroenteritis, vomiting, IV site
- irritation, urine discoloration (6.1). The most frequently occurring adverse reactions (> 1%) in neonates to 3 months of age
- were convulsions, diarrhea, oliguria/anuria, oral candidiasis, rash, tachycardia (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

- -DRUG INTERACTIONS-Ganciclovir: Generalized seizures have been reported in patients who received ganciclovir. Do not co-administer unless benefit outweighs risk (7.1).
- Probenecid: Concomitant administration of Imipenem and Cilastatin for Injection (I.V.) and probenecid results in increases in the plasma level and half-life of imipenem. Concomitant administration is not recommended (7.2).
- Valproic acid/divalproex sodium: Concomitant use with Imipenem and Cilastatin for Injection (I.V.) is generally not recommended. Consider other antibacterial drugs to treat infections in patients whose seizures are well-controlled on valproic acid or divalproex sodium (5.3, 7.3).
- --- USE IN SPECIFIC POPULATIONS--Renal Impairment: Dosage adjustment is necessary in patients with renal impairment (2.3). Adult patients with creatinine clearances of less than or equal to 30 mL/min, whether or not undergoing hemodialysis, had a higher risk of seizure activity than those without
- impairment of renal function (5.2). Therefore, close adherence to the dosing guidelines and regular monitoring of creatinine clearance for these patients is recommended (8.6).

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Females: (0.85) x (value calculated for males) Revised: 10/2017 Table 3: Dosage of Imipenem and Cilastatin for Injection (I.V.) for Adult Patients in Various Renal Function Groups Based on Estimated Creatinine Clearance (CLcr)

Suspected or Proven Pathogen Susceptibility

susceptible bacterial species

Microbiology (12.4)]

below

Aae

If the infection is suspected or proven to be due to a

If the infection is suspected or proven to be due to

exceed 4 g/day [see Dosage and Administration (2.1)].

mipenem and Cilastatin for Injection (I.V.) is not recommended in pediatric patients with CNS

Imipenem and Cilastatin for Injection (I.V.) is not recommended in pediatric patients < 30 kg with

Based on studies in adults, the maximum total daily dose in pediatric patients should not

The recommended dosage for pediatric patients with non-CNS infections is shown in Table 2

Table 2: Recommended Imipenem and Cilastatin for Injection (I.V.) Dosage in Pediatric

Less than or equal to 3 months of age (Greater than or equal to 1,500 g body weight)

Doses less than or equal to 500 mg should be given by intravenous infusion over 20 to 30 minutes

Doses greater than 500 mg should be given by intravenous infusion over 40 to 60 minutes

Patients with creatinine clearance less than 90 mL/min require dosage reduction of Imipenem

and Cilastatin for Injection (I.V.) as indicated in Table 3. The serum creatinine should represent a

steady state of renal function. Use the Cockroft-Gault method described below to calculate the

6 hours

end that the maximum total daily dosage not exceed 4 g/day

2.3 Dosage in Adult Patients with Renal Impairment

(weight in kg) x (140-age in years)

(72) x serum creatinine (mg/100 mL)

equal to 90

6 hours

1.000 ma

1.000 ma

every 8 hours 6 hours

every 6 hours 8 hours

* Administer doses less than or equal to 500 mg by intravenous infusion over 20 to 30 minutes.

patients who develop nausea during the infusion, the rate of infusion may be slowed

Administer doses greater than 500 mg by intravenous infusion over 40 to 60 minutes. In

In patients with creatinine clearances of less than 30 to greater than or equal to 15 mL/min, there may

be an increased risk of seizures *[see Warnings and Precautions (5.2) and Use in Specific Populations* (8.6)]. Patients with creatinine clearance less than 15 mL/min should not receive Imipenem and Cilastatin

for Injection (I.V.) unless hemodialysis is instituted within 48 hours. There is inadequate information to

recommend usage of Imipenem and Cilastatin for Injection (I.V.) for patients undergoing peritoneal dialysis.

When treating patients with creatinine clearances of less than 15 mL/min who are undergoing

30 to greater than or equal to 15 mL/min in Table 3 above *[see Dosage and Administration (2.3)*]

Both imipenem and cilastatin are cleared from the circulation during hemodialysis. The patient

should receive Imipenem and Cilastatin for Injection (I.V.) after hemodialysis and at intervals timed

from the end of that hemodialysis session. Dialysis patients, especially those with background CNS

disease, should be carefully monitored; for patients on hemodialysis, Imipenem and Cilastatin for

njection (I.V.) is recommended only when the benefit outweighs the potential risk of seizures. [See

Reconstitution and Preparation of Imipenem and Cilastatin for Injection (I.V.)

Do not use diluents containing benzyl alcohol to reconstitute Imipenem and Cilastatin for Injection

(I.V.) for administration to neonates because it has been associated with toxicity in neonates.

While toxicity has not been demonstrated in pediatric patients greater than three months of age.

Contents of the vials must be reconstituted by adding approximately 10 mL of the appropriate

Reconstituted Solutions of Imigenem and Cilastatin for Injection (I.V.) range from colorless to

After reconstitution, shake vial well and transfer the resulting suspension to 100 mL of an

Repeat transfer of the resulting suspension with an additional 10 mL of infusion solution

to ensure complete transfer of vial contents to the infusion solution. Agitate the resulting

Parenteral drug products should be inspected visually for particulate matter and discoloration

Imipenem and Cilastatin for Injection (I.V.), as supplied in single dose vials and reconstituted with

the appropriate diluents *[see Dosage and Administration (2,5)]*, maintains satisfactory potency

for 4 hours at room temperature or for 24 hours under refrigeration (5°C). Do not freeze solutions

yellow. Variations of color within this range do not affect the potency of the product.

The reconstituted suspension must not be administered by direct Intravenous Infusion

appropriate infusion solution before administering by intravenous infusion.

small pediatric patients in this age range may also be at risk for benzyl alcohol toxicity.

hemodialysis, use the dosage recommendations for patients with creatinine clearances of less that

500 mg every

infections because of the risk of seizures [see Use in Specific Populations (8.4)].

renal impairment, as no data are available [see Use in Specific Populations (8.4)].

bacterial species with intermediate susceptibility

(identified under column "I" in Table 10) /See

2.2 Dosage in Pediatric Patients

Patients for Non-CNS Infections

4 weeks to 3 months of age

I to 4 weeks of age

creatinine clearance

Dosage of Imipenem

Injection (I.V.)*,† If the

infection is suspected

or proven to be due to

a susceptible bacterial

Dosage of Imipenem

Injection (I.V.)*,† If the

infection is suspected

or proven to be due

to bacterial species

susceptibility (identified

with intermediate

under column "I"

in Table 10) [See

Microbiology (12.4)]:

2.4 Dosage in Hemodialysis Patients

Solution for Intravenous Administration

diluent to the vial. List of appropriate diluents are as follows:

• 5% Dextrose Injection with 0.225% or 0.45% saline solution

prior to administration, whenever solution and container permit.

5% Dextrose Injection with 0.15% potassium chloride solution

5% Dextrose and 0.9% Sodium Chloride Injection

Impenem and Cilastatin for Injection (I.V.) Vials

0.9% Sodium Chloride Injection

2.6 Storage of Reconstituted Solutions

of Imipenem and Cilastatin for Injection (I.V.).

5% or 10% Dextrose Injection

Mannitol 5% and 10%

mixture until clear.

Vials (After Reconstitution)

Warnings and Precautions (5.2)1.

2.5

and Cilastatin for

and Cilastatin for

species:

Males:

Less than 1 week of age

Greater than or equal to 3 Months of Age

- ADVERSE REACTIONS Clinical Trials Experience
- 6.2 Postmarketing Experience
- DRUG INTERACTIONS Ganciclovir
- Probenecid
- Valproic Acid 8 USE IN SPECIFIC POPULATIONS
 - Pregnancy 8.1 8.3 Nursing Mothers
 - 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment 10 OVERDOSAGE
- 11 DESCRIPTION

1.8 Endocarditis

1.9

1.10 Usage

empiric selection of therapy.

Dosage in Adults

For Intravenous Injection Only

Limitations of Use

- 12 CLINICAL PHARMACOLOGY
- Mechanism of Action 12.3 Pharmacokinetics
- Microbiology 12.4
- 13 NONCLINICAL TOXICOLOGY
- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility 15 REFERENCES

because safety and efficacy have not been established.

Populations (8.4), and Dosage and Administration (2.2)].

is advisable during prolonged therapy.

DOSAGE AND ADMINISTRATION

Warnings and Precautions (5.2), and Use in Specific Populations (8.4)].

- 16 HOW SUPPLIED/STORAGE AND HANDLING
- How Supplied 16.1
- Storage and Handling
- 17 PATIENT COUNSELING INFORMATION

1.7 Skin and Skin Structure Infections

*Sections or subsections omitted from the full prescribing information are not listed.

Imipenem and Cilastatin for Injection, USP (I.V.) is indicated for the treatment of skin and skin

structure infections caused by susceptible strains of *Enterococcus faecalis, Staphylococcus aureus*

(penicillinase-producing isolates), Staphylococcus epidermidis, Acinetobacter species, Citrobacter

species, Enterobacter species, Escherichia coli, Klebsiella species, Morganella morganii, Proteus

vulgaris, Providencia rettgeri, Pseudomonas aeruginosa, Serratia species, Peptococcus species,

Impenem and Cilastatin for Injection, USP (I.V.) is indicated for the treatment of endocarditis

Imipenem and Cilastatin for Injection, USP (I.V.) is not indicated in patients with meningitis

CNS infections because of the risk of seizures [see Dosage and Administration (2.2),

than 30 kg with impaired renal function, as no data are available [see Use in Specific

Impenent and Cilastatin for Injection LISP (LV) is not recommended in pediatric patients with

Impenent and Cilastatin for Injection, USP (I.V.) is not recommended in pediatric patients less

• Periodic assessment of organ system functions, including renal, hepatic and hematopoietic,

To reduce the development of drug-resistant bacteria and maintain the effectiveness of

Imipenem and Cilastatin for Injection, USP (I.V.) and other antibacterial drugs, Imipenem and

Cilastatin for Injection, USP (I.V.) should be used only to treat infections that are proven or strongly

suspected to be caused by susceptible bacteria. When culture and susceptibility information

are available, they should be considered in selecting or modifying antibacterial therapy. In the

absence of such data, local epidemiology and susceptibility patterns may contribute to the

• The dosage of Imipenem and Cilastatin for Injection (I.V.) in adult patients should be based on

suspected or confirmed pathogen susceptibility as shown in Table 1 below. The dosage

recommendations for Imipenem and Cilastatin for Injection (I.V.) represent the quantity of imipenem to be administered. An equivalent amount of cilastatin is also present in the solution

These doses should be used for patients with creatinine clearance of greater than or equal

to 90 mL/min. A reduction in dose must be made for patients with creatinine clearance less

In patients who develop nausea during the infusion, the rate of infusion may be slowed.

than 90 mL/min as shown in Table 3 Isee Dosage and Administration (2.3)].

Recommend that the maximum total daily dosage not exceed 4 g/day.

Administer 500 mg by intravenous infusion over 20 to 30 minutes

Administer 1,000 mg by intravenous infusion over 40 to 60 minutes

Peptostreptococcus species, Bacteroides species including B. fragilis, Fusobacterium species.

caused by susceptible strains of Staphylococcus aureus (penicillinase-producing isolates).

 Table 1: Dosage of Imipenem and Cilastatin for Injection (I.V.) in Adult Patients with Creatinine Clearance Greater than or Equal to 90 mL/min
 2.7
 Incompatibility and Compatibility of Imipenem and Cilastatin for Injection (I.V.) with other Antibacterial Drugs

 Suspected or Proven Pathogen Suspentibility
 Desage of Imipenem and Cilastatin
 0 not mix Imipenem and Cilastatin for Injection (I.V.)

antibacterial drugs Imipenem and Cilastatin for Injection (I.V.) may be administered concomitantly with other antibacterial drugs, such as aminoglycosides.

DOSAGE FORMS AND STRENGTHS

For Injection Imipenem and Cilastatin for Injection (I.V.) is a sterile powder mixture for reconstitution in single-dose vials containing: 250 mg imipenem (anhydrous equivalent) and 250 mg cilastatin sodium

• 500 mg imipenem (anhydrous equivalent) and 500 mg cilastatin sodium

CONTRAINDICATIONS

Dosage of Imipenem and Cilastatin

500 mg every 6 hours

1,000 mg every 8 hours

1,000 mg every 6 hours

for Injection (I.V.)

Dose (mg/kg) *,[†] Frequency (hours)

Every 6 hours

Every 6 hours

Every 8 hours

Every 12 hours

15-25 mg/kg

25 mg/kg

25 mg/kg

25 mg/kg

Creatinine clearance (mL/min)

Greater than or Less than 90 to Less than 60 to Less than 30 to

OR

750 mg every 500 mg every

greater than greater than greater than or

or equal to 60 or equal to 30 equal to 15

400 mg every 300 mg every 200 mg every

500 mg every 500 mg every 500 mg every

6 hours

12 hours

500 mg everv

12 hours

6 hours

8 hours

6 hours

Imipenem and Cilastatin for Injection (I.V.) is contraindicated in patients who have shown hypersensitivity to any component of this product.

WARNINGS AND PRECAUTIONS 5.1

Hypersensitivity Reactions Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients receiving therapy with beta-lactams. These reactions are more likely to occur in

individuals with a history of sensitivity to multiple allergens. There have been reports of individuals with a history of penicillin hypersensitivity who have experienced severe hypersensitivity reactions when treated with another beta-lactam. Before initiating therapy with Imipenem and Cilastatin for Injection (I.V.), careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, other betalactams and other allergens. If an allergic reaction to Imipenem and Cilastatin for Injection (I.V.) occurs, discontinue the drug immediately. Serious anaphylactic reactions require immediate emergency treatment as clinically indicated.

5.2 Seizure Potential

Seizures and other CNS adverse experiences, such as confusional states and myoclonic activity. have been reported during treatment with Imipenem and Cilastatin for Injection (I.V.), especially when recommended dosages were exceeded [see Adverse Reactions (6.1, 6.2)]. These experiences have occurred most commonly in patients with CNS disorders (e.g., brain lesions or history of seizures) and/or compromised renal function [see Use in Specific Populations (8.6)]. However, there have been reports of CNS adverse experiences in patients who had no recognized or documented underlying CNS disorder or compromised renal function.

Anticonvulsant therapy should be continued in patients with known seizure disorders. If focal tremors, myoclonus, or seizures occur, patients should be evaluated neurologically, placed on anticonvulsant therapy if not already instituted, and the dosage of Imipenem and Cilastatin for Injection (I.V.) re-examined to determine whether it should be decreased or the antibacterial drug discontinued

5.3 Increased Seizure Potential Due to Interaction with Valproic Acid

Case reports in the literature have shown that co-administration of carbapenems, including Imipenem and Cilastatin for Injection (I.V.), to patients receiving valproic acid or divalproex sodium results in a reduction in valproic acid concentrations. The valproic acid concentrations may drop below the therapeutic range as a result of this interaction, therefore increasing the risk of breakthrough seizures. Increasing the dose of valproic acid or divalproex sodium may not be sufficient to overcome this interaction. The concomitant use of Imipenem and Cilastatin for Injection (I.V.) and valproic acid/divalproex sodium is generally not recommended

Antibacterials other than carbapenems should be considered to treat infections in patients whose seizures are well controlled on valproic acid or divalproex sodium. If administration of Imipenem and Cilastatin for Injection (I.V.) is necessary, supplemental anti-convulsant therapy should be considered [see Drug Interactions (7.3)]. Close adherence to the recommended dosage and dosage schedules is urged, especially in patients with known factors that predispose to convulsive activity.

5.4 Clostridium difficile-Associated Diarrhea (CDAD)

Clostridium difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including Imipenem and Cilastatin for Injection (I.V.), and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of C. difficile.

C. difficile produces toxins A and B which contribute to the development of CDAD.

Hypertoxin producing strains of C. difficile cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial drug use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibacterial drug use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial drug treatment of C. difficile, and surgical evaluation should be instituted as clinically indicated

5.5 Development of Drug-Resistant Bacteria

As with other antibacterial drugs, prolonged use of Imipenem and Cilastatin for Injection (I.V.) may result in overgrowth of nonsusceptible organisms. Repeated evaluation of the patient's condition is essential. If superinfection occurs during therapy, appropriate measures should be taken.

Prescribing Impenem and Cilastatin for Injection (I.V.) in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

ADVERSE REACTIONS

The following serious adverse reactions are described in greater detail in the Warnings and Precautions section.

- Hypersensitivity Reactions [see Warnings and Precautions (5.1)]
- Seizure Potential [see Warnings and Precautions (5.2)]
 Increased Seizure Potential Due to Interaction with Valproic Acid [see Warnings and
- Precautions (5.3) Clostridium difficile-Associated Diarrhea (CDAD) [see Warnings and Precautions (5.4)]
- Development of Drug-Resistant Bacteria [see Warnings and Precautions (5.5)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adult Patients

During clinical investigations 1,723 patients were treated with Imipenem and Cilastatin for Injection (LV). Table 4 shows the incidence of adverse reactions reported during the clinical nvestigations of adult patients treated with Imipenem and Cilastatin for Injection (I.V.).

Table 4: Incidence (%)* of Adverse Reactions Reported During Clinical Investigations of Adult Patients Treated with Imipenem and Cilastatin for Injection (I.V.)

Body System	Adverse Reactions	Frequency (%)
Local Administration site	Phlebitis/ thrombophlebitis	3.1%
	Pain at the injection site	0.7%
	Erythema at the injection site	0.4%
	Vein induration	0.2%
Gastrointestinal	Nausea	2%
	Diarrhea	1.8%
	Vomiting	1.5%
Skin	Rash	0.9%
	Pruritus	0.3%
	Urticaria	0.2%
Vascular	Hypotension	0.4%
Body as a Whole	Fever	0.5%
Nervous system	Seizures	0.4%
	Dizziness	0.3%
	Somnolence	0.2%

* Adverse reactions with an incidence $\geq 0.2\%$ of Imipenem and Cilastatin for Injection (I.V.) -treated adult patients.

Additional adverse reactions reported in less than 0.2% of the patients or reported since the drug was marketed are listed within each body system in order of decreasing severity (see Table 5).

Table 5: Additional Adverse Reactions Occurring in Less than 0.2% of Adult Patients Listed within Each Body System in Order of Decreasing Severity

Body System	Adverse Reactions
Gastrointestinal	Pseudomembranous Colitis (the onset of Pseudomembranous colitis symptoms), Hemorrhagic Colitis
	Gastroenteritis
	Abdominal Pain
	Glossitis
	Tongue Papillar
	Hypertrophy
	Heartburn
	Pharyngeal Pain
	Increased Salivation
CNS	Encephalopathy
	Confusion
	Myoclonus
	Paresthesia
	Vertigo
	Headache
Special Senses	Hearing Loss
	Tinnitus
Respiratory	Chest Discomfort
	Dyspnea
	Hyperventilation
	Thoracic Spine Pain
Cardiovascular	Palpitations
	Tachycardia
Skin	Erythema Multiforme
	Angioneurotic Edema
	Flushing
	Cyanosis
	Hyperhidrosis
	Skin Texture Changes
	Candidiasis
	Pruritus Vulvae
Local Administration site	Infused vein infection
Body as a Whole	Polyarthralgia
	Asthenia/Weakness
Renal	Oliguria/Anuria
	Polyuria

FOR INJECTION, USP (I.V.)



451205D/Revised: October 2017

451205D/Revised: October 2017

IMIPENEM AND CILASTATIN

FOR INJECTION. USP (I.V.)



Adverse Laboratory Changes

The following adverse laboratory changes were reported during clinical trials: *Hepatic:* Increased alanine aminotransferase (ALT or SGPT), aspartate aminotransferase (AST or

SGOT), alkaline phosphatase, bilirubin, and lactate dehydrogenase (LDH Hemic: Increased eosinophils, positive Coombs test, increased WBC, increased platelets, decreased hemoglobin and hematocrit, increased monocytes, abnormal prothrombin time, increased lymphocytes, increased basophils

Electrolytes: Decreased serum sodium, increased potassium, increased chloride *Renal:* Increased BUN, creatinine

Urinalysis: Presence of urine protein, urine red blood cells, urine white blood cells, urine casts, urine bilirubin, and urine urobilinoger

<u>Pediatric Patients</u> Table 6: Incidence (%)* of Adverse Reactions Reported During Clinical Investigations of Pediatric Patients Greater Than or Equal to 3 Months of Age Treated with Imipenem and Cilastatin for Injection (I,V)

Body System	Adverse Reactions	Frequency (%)	
Local Administration Site	Phlebitis	2.2%	
	Intravenous Site Irritation	1.1%	
Gastrointestinal	Diarrhea	3.9%	
	Gastroenteritis	1.1%	
	Vomiting	1.1%	
Skin	Rash	2.2%	
Renal	Urine Discoloration	1.1%	

^t Adverse reactions that occurred in > 1 % of Imipenem and Cilastatin for Injection (I.V.)-treated pediatric patients (greater than or equal to 3 months of age)

Table 7: Incidence (%)* of Adverse Reactions Reported During Clinical Investigations of Pediatric Patients Neonates to 3 Months of Age Treated with Imipenem and Cilastatin for Injection (I.V.)

Body System	Adverse Reactions	Frequency (%)	
Gastrointestinal	Diarrhea	3%	
CNS	Convulsions	5.9%	
Cardiovascular	Tachycardia	1.5%	
Skin	Rash	1.5%	
Body as a Whole	Oral Candidiasis	1.5%	
Renal	Oliguria/Anuria	2.2%	

 * Adverse reactions that occurred in > 1 % of Imipenem and Cilastatin for Injection (I.V.)-treated pediatric patients (neonates to 3 months of age)

Adverse Laboratory Changes

The following adverse laboratory changes were reported in studies of 178 pediatric patients 3 months of age: increased AST (SGOT), decreased hemoglobin/hematocrit, increased platelets, increased eosinophils, increased ALT (SGPT), increased urine protein, decreased neutrophils.

The following adverse laboratory changes were reported in studies of 135 patients (neonates to 3 months of age): increased eosinophils, increased AST (SGPT), increased serum creatinine, increased/decreased platelet count, increased/decreased bilirubin, increased ALT (SGPT), increased alkaline phosphatase, increased/decreased hematocrit.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of Imipenem and Cilastatin for Injection (I.V). Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Table 8: Adverse Reactions Identified During Post Approval Use of Imipenem and Cilastatin for Injection (I.V.)

Body System	Adverse Reactions
Gastrointestinal	Hepatitis (including fulminant hepatitis)
	Hepatic failure
	Jaundice
	Staining of the teeth and/or tongue
Hematologic	Pancytopenia
	Bone marrow depression
	Thrombocytopenia
	Neutropenia
	Leukopenia
	Hemolytic anemia
CNS	Tremor
	Psychic disturbances including hallucinations
	Dyskinesia
	Agitation
Special Senses	Taste perversion
Skin	Stevens-Johnson syndrome
	Toxic epidermal necrolysis
Body as a whole	Drug fever
Renal	Acute renal failure
	Urine discoloration

Adverse Laboratory Changes

Adverse laboratory changes reported since the drug was marketed were: *Hematologic:* agranulocytosis.

Examination of published literature and spontaneous adverse reactions reports suggested a similar spectrum of adverse reactions in adult and pediatric patients.

7 DRUG INTERACTIONS 7.1 Ganciclovir

Generalized seizures have been reported in patients who received ganciclovir and Imipenem and Cilastatin for Injection (I.V.). These drugs should not be used concomitantly with Imipenem and Cilastatin for Injection (I.V.) unless the potential benefits outweigh the risks.

7.2 Probenecid

Concomitant administration of Imipenem and Cilastatin for Injection (I.V.) and probenecid results in increases in the plasma level and half-life of imipenem. Therefore, it is not recommended that probenecid be given concomitantly with Imipenem and Cilastatin for Injection (I.V.).

7.3 Valproic Acid

Case reports in the literature have shown that co-administration of carbapenems, including Imipenem and Cilastatin for Injection (I.V.), to patients receiving valproic acid or divalproex sodium results in a reduction in valproic acid concentrations. The valproic acid concentrations may drop below the therapeutic range as a result of this interaction, therefore increasing the risk of breakthrough seizures. Although the mechanism of this interaction is unknown, data from *in vitro* and animal studies suggest that carbapenems may inhibit the hydrolysis of valproic acid's glucuronide metabolite (VPA-g) back to valproic acid, thus decreasing the serum concentrations of valproic acid *[see Warnings and Precautions (5.3)]*. The concomitant use of Imipenem and Cilastatin for Injection (I.V.) and valproic acid/divalproex sodium is generally not recommended. Antibacterials other than carbapenems should be considered to treat infections in patients whose seizures are well-controlled on valproic acid or divalproex sodium.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy Pregnancy Category C.

There are no adequate and well-controlled studies of Imipenem and Cilastatin for Injection (I.V.) in pregnant women. Imipenem and Cilastatin for Injection (I.V.) should be used during pregnancy only if the potential benefit justifies the potential risk to the mother and fetus.

Developmental toxicity studies with imipenem and cilastatin sodium (alone or in combination) administered to monkeys, rabbits, rats, and mice revealed no evidence of teratogenicity. Imipenem was administered intravenously to rabbits and rats at doses up to 60 and 900 mg/kg/day, respectively, up to approximately 0.4 and 2.9 time the maximum recommended human daily dose as a component of Imipenem and Cilastatin for Injection (I.V), based on body surface area. Cilastatin sodium was given intravenously to rabbits at doses up to 300 mg/kg/day and to rats subcutaneously at doses up to 1,000 mg/kg/day, up to approximately 1.9 and 3.2 times the maximum recommended human daily dose as a component of Imipenem and Cilastatin for Injection (I.V), based on body surface area. Imipenem-cilastatin sodium was given intravenously at doses up to 80 mg/kg/day and subcutaneously at doses up to 320 mg/kg/day to mice and rats (the higher dose is approximately equal to the highest recommended human daily dose based on body surface area). Intravenous doses of imipenem-cilastatin sodium at approximately 100 mg/kg/day (0.6 times the maximum recommended human daily dose, based on body surface area) administered to pregnant cynomolgus monkeys at an infusion rate mimicking human clinical use were not associated with teratogenicity, but there was an increase in embryonic loss relative to controls. However, an imipenem-cilastatin dose of 40 mg/kg given to pregnant cynomolgus monkeys by bolus intravenous injection caused significant maternal toxicity including death and embryofetal loss.

No adverse effects on the fetus or on lactation were observed when imipenem-cilastatin sodium was administered subcutaneously to rats late in gestation at dosages up to 320 mg/kg/day, approximately equal to the highest recommended human dose (based on body surface area). Although a slight decrease in live fetal body weight was observed at the high dose, there were no adverse effects on fetal viability, growth or postnatal development of pups.

8.3 Nursing Mothers

It is not known whether imipenem-cilastatin sodium is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Imipenem and Cilastatin for Injection (I.V.) is administered to a nursing woman.

Pediatric Use

Use of Imipenem and Cilastatin for Injection (I.V.) in pediatric patients is supported by evidence from adequate and well-controlled trials of Imipenem and Cilastatin for Injection (I.V.) in adults and clinical studies in pediatric patients [see Dosage and Administration (2.2) and Clinical Pharmacology (12.3)].

Imipenem and Cilastatin for Injection (I.V.) is not recommended in pediatric patients with CNS

infections because of the risk of seizures.

Imipenem and Cilastatin for Injection (I.V.) is not recommended in pediatric patients less than 30 kg with renal impairment, as no data are available.

8.5 Geriatric Use

Of the approximately 3,600 subjects \geq 18 years of age in clinical studies of Imipenem and Cilastatin for Injection (I.V.), including postmarketing studies, approximately 2,800 received Imipenem and Cilastatin for Injection (I.V.). Of the subjects who received Imipenem and Cilastatin for Injection (I.V.), data are available on approximately 800 subjects who were 65 and over, including approximately 300 subjects who were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

No dosage adjustment is required based on age [see Clinical Pharmacology (12.3)]. Dosage adjustment in the case of renal impairment is necessary [see Dosage and Administration (2.3)].

8.6 Renal Impairment

Dosage adjustment is necessary in patients with renal impairment [see Dosage and Administration (2.3)]. Adult patients with creatinine clearances of less than or equal to 30 mL/min, whether or not undergoing hemodialysis, had a higher risk of seizure activity than those without impairment of renal function [see Warnings and Precautions (5.2)]. Therefore, close adherence to the dosing guidelines and regular monitoring of creatinine clearance for these patients is recommended.


10 OVERDOSAGE

In the case of overdosage, discontinue Imipenem and Cilastatin for Injection (I.V.), treat symptomatically, and institute supportive measures as required. Imipenem and Cilastatin for Injection (I.V.) is hemodialyzable.


11 DESCRIPTION

Imipenem and Cilastatin for Injection, USP (I.V.) (imipenem and cilastatin) for Injection is a sterile formulation of imipenem, a penem antibacterial, and cilastatin, a renal dehydropeptidase inhibitor with sodium bicarbonate added as a buffer. Imipenem and Cilastatin for Injection, USP (I.V.) is an antibacterial drug for intravenous administration.

Imipenem (N-formimidoylthienamycin monohydrate) is a crystalline derivative of thienamycin, which is produced by *Streptomyces cattleya*. Its chemical name is (5R,6S)-3-[[2-(formimidoylamino)ethyl]thio]-6-[(R)-1-hydroxyethyl]-7-oxo-1-azabicyclo[3.2.0]hept-2-ene-2-carboxylic acid monohydrate. It is an off-white, nonhygroscopic crystalline compound with a molecular weight of 317.37. It is sparingly soluble in water and slightly soluble in methanol. Its empirical formula is $C_{12}H_1$, N_3O_4 S- H_2O , and its structural formula is:



Cilastatin sodium is the sodium salt of a derivatized heptenoic acid. Its chemical name is sodium (*Z*)-7[[(*R*)-2-amino-2-carboxyethyl]thio]-2-[(*S*)-2,2-dimethylcyclopropanecarboxamido]-2-heptenoate. It is an off-white to yellowish-white, hygroscopic, amorphous compound with a molecular weight of 380.43. It is very soluble in water and in methanol. Its empirical formula is $C_{1,e}H_{2,e}N_2O_{e}SNa$, and its structural formula is:



Imipenem and Cilastatin for Injection, USP (I.V.) is buffered to provide solutions in the pH range of 6.5 to 8.5. There is no significant change in pH when solutions are prepared and used as directed. *[See How Supplied/ Storage and Handling (16.1),]* Each Imipenem and Cilastatin for Injection, USP (I.V.) 250 mg/250 mg vial contains imipenem USP 250 mg (anhydrous equivalent) and cilastatin sodium USP equivalent to 250 mg cilastatin and each 500 mg/500 mg vial contains imipenem USP 500 mg (anhydrous equivalent) and cilastatin sodium USP equivalent to 500 mg/250 mg vial contains 10 mg of sodium bicarbonate and the 500 mg/500 mg vial contains 20 mg of sodium bicarbonate. The sodium content of the 250 mg/250 mg vial is 18.8 mg (0.8 mEq) and the sodium content for the 500 mg/500 mg vial is 37.5 mg (1.6 mEq). Solutions of Imipenem and Cilastatin for Injection, USP (I.V.) range from colorless to yellow. Variations of color within this range do not affect the potency of the product.

2 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Imipenem and Cilastatin for Injection (I.V.) is a combination of imipenem and cilastatin. Imipenem is a penem antibacterial drug *[see Microbiology (12.4)]*. Cilastatin sodium is a renal dehydropeptidase inhibitor that limits the renal metabolism of imipenem.

12.3 Pharmacokinetics

Intravenous infusion of Imipenem and Cilastatin for Injection (I.V.) over 20 minutes results in peak plasma levels of imipenem antimicrobial activity that range from 21 to 58 mcg/mL for the 500 mg dose, and from 41 to 83 mcg/mL for the 1,000 mg dose. At these doses, plasma levels of imipenem antimicrobial activity decline to below 1 mcg/mL or less in 4 to 6 hours. Peak plasma levels of cilastatin following a 20-minute intravenous infusion of Imipenem and Cilastatin for Injection (I.V.) range from 31 to 49 mcg/mL for the 500 mg dose, and from 56 to 88 mcg/mL for the 1,000 mg dose.

The binding of imipenem to human serum proteins is approximately 20% and that of cilastatin is approximately 40%.

Imipenem has been shown to penetrate into human tissues, including vitreous humor, aqueous humor, lung, peritoneal fluid, CSF, bone, interstitial fluid, skin, and fascia. As there are no adequate and well-controlled studies of imipenem treatment in these additional body sites, the clinical significance of these tissue concentration data is unknown.

After a 1 gram dose of Imipenem and Cilastatin for Injection (I.V.), the following average levels of imipenem were measured (usually at 1 hour post dose except where indicated) in the tissues and fluids listed in Table 9:

Table 9: Average Levels of Imipenen

Tuble 5. Arenuge Leve	io or minp	onom	
Tissue or Fluid	Ν	Imipenem Level mcg/mL or mcg/g	Range
Vitreous Humor	3	3.4 (3.5 hours post dose)	2.88 to 3.6
Aqueous Humor	5	2.99 (2 hours post dose)	2.4 to 3.9
Lung Tissue	8	5.6 (median)	3.5 to 15.5
Sputum	1	2.1	—
Pleural	1	22	—
Peritoneal	12	23.9 S.D.±5.3 (2 hours post dose)	_
Bile	2	5.3 (2.25 hours post dose)	4.6 to 6
CSF (uninflamed)	5	1 (4 hours post dose)	0.26 to 2
CSF (inflamed)	7	2.6 (2 hours post dose)	0.5 to 5.5
Fallopian Tubes	1	13.6	—
Endometrium	1	11.1	—
Myometrium	1	5	—
Bone	10	2.6	0.4 to 5.4
Interstitial Fluid	12	16.4	10 to 22.6
Skin	12	4.4	NA
Fascia	12	4.4	NA

Metabolism

Imipenem, when administered alone, is metabolized in the kidneys by dehydropeptidase I, resulting in relatively low levels in urine. Cilastatin sodium, an inhibitor of this enzyme, effectively prevents renal metabolism of imipenem so that when imipenem and cilastatin sodium are given concomitantly, adequate antibacterial levels of imipenem are achieved in the urine.

<u>Elimination</u>

The plasma half-life of each component is approximately 1 hour. Approximately 70% of the administered imipenem is recovered in the urine within 10 hours after which no further urinary excretion is detectable. Urine concentrations of imipenem in excess of 10 mcg/mL can be maintained for up to 8 hours with Imipenem and Cilastatin for Injection (I.V.) at the 500-mg dose. Approximately 70% of the cilastatin sodium dose is recovered in the urine within 10 hours of administration of Imipenem and Cilastatin for Injection (I.V.). Imipenem-cilastatin sodium is hemodialyzable *[see Overdosage (10)]*.

No accumulation of imipenem/cilastatin in plasma or urine is observed with regimens administered as frequently as every 6 hours in patients with normal renal function.

Specific Populations Geriatric Patients

In healthy elderly volunteers (65 to 75 years of age with normal renal function for their age), the pharmacokinetics of a single dose of imipenem 500 mg and cilastatin 500 mg administered intravenously over 20 minutes are consistent with those expected in subjects with slight renal impairment for which no dosage alteration is considered necessary. The mean plasma half-lives of imipenem and cilastatin are 91 ± 7 minutes and 69 ± 15 minutes, respectively. Multiple dosing has no effect on the pharmacokinetics of either imipenem or cilastatin, and no accumulation of imipenem/cilastatin is observed.

Pediatric Patients

Doses of 25 mg/kg/dose in patients 3 months to < 3 years of age, and 15 mg/kg/dose in patients 3 to 12 years of age were associated with mean trough plasma concentrations of imipenem of 1.1 ± 0.4 mcg/mL and 0.6 ± 0.2 mcg/mL following multiple 60-minute infusions, respectively; trough urinary concentrations of imipenem were in excess of 10 mcg/mL for both doses. These doses have provided adequate plasma and urine concentrations for the treatment of non-CNS infections.

In a dose-ranging study of smaller premature infants (670 to 1,890 g) in the first week of life, a dose of 20 mg/kg q12h by 15 to 30 minutes infusion was associated with mean peak and trough plasma imipenem concentrations of 43 mcg/mL and 1.7 mcg/mL after multiple doses, respectively. However, moderate accumulation of cilastatin in neonates may occur following multiple doses of Imipenem and Cilastatin for Injection (I.V.). The safety of this accumulation is unknown.

12.4 Microbiology Mechanism of Action

Imipenem and Cilastatin for Injection (I.V.) is a combination of imipenem and cilastatin. The bactericidal activity of imipenem results from the inhibition of cell wall synthesis. Its greatest affinity is for penicillin binding proteins (PBPs) 1A, 1B, 2, 4, 5 and 6 of *Escherichia coli*, and 1A, 1B, 2, 4 and 5 of *Pseudomonas aeruginosa*. The lethal effect is related to binding to PBP 2 and PBP 1B.

Imipenem has a high degree of stability in the presence of beta-lactamases, both penicillinases and cephalosporinases produced by Gram-negative and Gram-positive bacteria. It is a potent inhibitor of beta-lactamases from certain Gram-negative bacteria which are inherently resistant to most beta-lactam antibacterials, e.g., *Pseudomonas aeruginosa, Serratia* spp., and *Enterobacter* spp.

<u>Resistanc</u>

Imipenem is inactive *in vitro* against *Enterococcus faecium*, *Stenotrophomonas maltophilia* and some isolates of *Burkholderia cepacia*. Methicillin-resistant staphylococci should be reported as resistant to imipenem.

Interaction with Other Antimicrobials

In vitro tests show imipenem to act synergistically with aminoglycoside antibacterials against some isolates of *Pseudomonas aeruginosa*.

Antimicrobial Activity

Imipenem has been shown to be active against most isolates of the following microorganisms, both *in vitro* and in clinical infections *[see Indications and Usage (1.1)]*.

Aerobic bacteria

Gram-positive bacteria Enterococcus faecalis Staphylococcus aureus Staphylococcus epidermidis Streptococcus agalactiae (Group B streptococci) Streptococcus pneumoniae Streptococcus pvogenes

Gram-negative bacteria

Acinetobacter spp. Citrobacter spp. Enterobacter spp. Escherichia coli Gardnerella vaginalis Haemophilus influenzae Haemophilus parainfluenzae Klebsiella spp. Morganella morganii Proteus vulgaris Providencia rettgeri Pseudomonas aeruginosa Serratia spp., including *S. marcescens*

Anaerobic bacteria

Gram positive bacteria Bifidobacterium spp. Clostridium spp. Eubacterium spp. Peptococcus spp. Peptostreptococcus spp. Propionibacterium spp.

<u>Gram-negative bacteria</u> Bacteroides spp., including B. fragilis Fusobacterium spp. The following *in vitro* data are available, but their clinical significance is unknown. At least 90 percent of the following bacteria exhibit an *in vitro* minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for imipenem against isolates of similar genus or organism group. However, the efficacy of imipenem in treating clinical infections due to these bacteria has not been established in adequate and well-controlled clinical trials.

Aerobic bacteria

Gram-positive bacteria

- Bacillus spp. Listeria monocytogenes Nocardia spp. Staphylococcus saprophyticus
- Group C streptococci Group G streptococci Viridans group streptococci

<u>Gram-negative bacteria</u> Aeromonas hydrophila

Alcaligenes spp. Capnocytophaga spp. Haemophilus ducreyi Neisseria gonorrhoead

Pasteurella spp. Providencia stuartii

Anaerobic bacteria

Prevotella bivia Prevotella disiens

Prevotella disfens Prevotella melaninogenica Veillonella spp.

Susceptibility Test Methods

When available, the clinical microbiology laboratory should provide cumulative reports of *in vitro* susceptibility tests results for antimicrobial drugs used in local hospitals and practice areas to the physician as periodic reports that describe the susceptibility profile of nosocomial and community-acquired pathogens. These reports should aid the physician in selecting an antibacterial drug for treatment.

Dilution Techniques:

Quantitative methods are used to determine antimicrobial MICs. These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized test method (broth and/or agar).^{1,2} The MIC values should be interpreted according to breakpoints provided in Table 10.

Diffusion Techniques:

Quantitative methods that require measurement of zone diameters can also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. The zone size should be determined using a standardized test method.^{2,3} This procedure uses paper disks impregnated with 10-mcg of imipenem to test the susceptibility of bacteria to imipenem. The disk diffusion breakpoints are provided in Table 10.

Anaerobic Techniques:

For anaerobic bacteria, the susceptibility to imipenem can be determined by a standardized test method.⁴ The MIC values obtained should be interpreted according to the breakpoints provided in Table 10.

Table 10: Susceptibility Test Interpretive Criteria for Imipenem*

	Minimum Inhibitory Concentrations MIC (mcg/mL)		Disk Diffusion (zone diameters in mm)			
Pathogen	S	I	R	S	I	R
Enterobacteriaceae	≤ 1	2†	≥ 4	≥ 23	20 to 22	≤ 19
Pseudomonas aeruginosa	≤ 2	4†	≥ 8	≥ 19	16 to 18	≤ 15
Acinetobacter spp.	≤ 2	4†	≥ 8	≥ 22	19 to 21	≤ 18
<i>Haemophilus influenza</i> and <i>H. parainfluenzae</i> ‡	≤ 4	-	-	≥ 16	-	-
Streptococcus pneumoniae§	≤ 0.12	0.25 to 0.5	≥ 1	-	-	-
Anaerobes	≤ 4	8	≥ 16	-	-	-

Interpretive criteria are based on a dosing regimen of 500 mg every 6 hours or 1,000 mg every 8 hours.

 Use 1,000 mg every 6 hours for bacteria with intermediate susceptibility in patients with creatinine clearance greater than or equal to 90 mL/min.
 The current absence of data on resistant isolates precludes defining any category other than

Susceptible'. If isolates yield MIC results other than susceptible, they should be submitted to a reference laboratory for additional testing.

[§] For non-meningitis *Ś. pneumoniae* isolates, penicillin MICs ≤ 0.06 mcg/mL (or oxacillin zones ≥ 20 mm) indicate susceptibility to imipenem.

Susceptibility of staphylococci to imipenem may be deduced from testing penicillin and either cefoxitin or oxacillin. $^{\rm 2}$

A report of "Susceptible" (S) indicates that the antimicrobial drug is likely to inhibit growth of the pathogen if the antimicrobial drug reaches the concentration usually achievable at the site of infection. A report of "Intermediate" (I) indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where a high dosage of the drug can be used. This category also provides a buffer zone that prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "Resistant" (R) indicates that the antimicrobial drug is not likely to inhibit growth of the pathogen if the antimicrobial drug be selected.

Quality Control

Standardized susceptibility test procedures require the use of laboratory controls to monitor and ensure the accuracy and precision of supplies and reagents used in the assay, and the techniques of the individuals performing the test.^{1,2,3,4} Standard imipenem powder should provide the following range of MIC values noted in Table 11. For the diffusion technique using the 10 mcg disk, the criteria in Table 11 should be achieved.

Table 11: Acceptable Quality Control Ranges for Imipenem

Microorganism	Minimum Inhibitory Concentrations (mcg/mL)	Disk Diffusion (zone diameters in mm)
<i>Bacteroides fragilis</i> ATCC 25285 [*]	0.03 to 0.125° 0.03 to 0.25 ⁺	-
<i>Bacteroides thetaiotaomicron</i> ATCC 29741	0.125 to 0.5 [°] 0.25 to 1 ⁺	-
<i>Eggerthella lenta</i> ATCC 43055	0.125 to 0.5 [*] 0.25 to 2 [†]	-
<i>Enterococcus faecalis</i> ATCC 29212	0.5 to 2	-
<i>Escherichia coli</i> ATCC 25922	0.06 to 0.25	26 to 32
<i>Haemophilus influenzae</i> ATCC 49247	-	21 to 29
<i>Haemophilus influenzae</i> ATCC 49766	0.25 to 1	-
<i>Staphylococcus aureus</i> ATCC 29213	0.015 to 0.06	-
<i>Pseudomonas aeruginosa</i> ATCC 27853	1 to 4	20 to 28
<i>Streptococcus pneumoniae</i> ATCC 49619	0.03 to 0.12	-

13 NONCLINICAL TOXICOLOGY 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long term studies in animals have not been performed to evaluate carcinogenic potential of imipenem-cilastatin. A variety of bacterial and mammalian tests were performed to evaluate genetic toxicity. The tests used were: V79 mammalian cell mutagenesis assay (cilastatin sodium alone and imipenem alone), Ames test (cilastatin sodium alone and imipenem alone), unscheduled DNA synthesis assay (imipenem-cilastatin sodium) and *in vivo* mouse cytogenetics test (imipenem-cilastatin sodium). None of these tests showed any evidence of genetic alterations.

Impairment of fertility or reproductive performance was not observed in male and female rats given imipenem-cilastatin at intravenous doses up to 80 mg/kg/day and at a subcutaneous dose of 320 mg/kg/day. In rats, a dose of 320 mg/kg/day was approximately equal to the highest recommended human dose based on body surface area.

15 REFERENCES

- Clinical and Laboratory Standards Institute (CLSI). Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically; Approved Standard - Tenth Edition. CLSI document M07-A10, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA, 2015.
- Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Susceptibility Testing; Twenty-sixth Informational Supplement, CLSI document M100-S26, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsvlvania 19087. USA. 2016.
- Clinical and Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Disk Diffusion Susceptibility Tests; Approved Standard – Twelfth Edition. CLSI document M02-A12, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA, 2015.
- Clinical and Laboratory Standards Institute (CLSI). Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria; Approved Standard – Eighth Edition. CLSI document M11-A8 Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA 2012.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied Imipenem and Cilastatin for Injection, USP (I.V.) is supplied as a sterile powder mixture in single dose vials containing imipenem (anhydrous equivalent) and cilastatin sodium as follows:

Product No.	NDC No.		
349025	63323-349-25	250 mg imipenem equivalent and 250 mg cilastatin equivalent and 10 mg sodium bicarbonate as a buffer	Packaged in carton of 25.
342025	63323-322-25	500 mg imipenem equivalent and 500 mg cilastatin equivalent and 20 mg sodium bicarbonate as a buffer	Packaged in carton of 25.

16.2 Storage and Handling Before Reconstitution:

Imipenem and Cilastatin for Injection, USP (I.V.) sterile powder should be stored at 20° to 25° C (68° to 77° F) [See USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

- Advise the patient to read the FDA-approved patient labeling (Instructions for Use). • Advise patients that allergic reactions, including serious allergic reactions, could occur
- and that serious reactions require immediate treatment. They should report any previous hypersensitivity reactions to Imipenem and Cilastatin for Injection (I.V.), other carbapenems, beta-lactams or other allergens.
- Counsel patients that antibacterial drugs including Imipenem and Cilastatin for Injection (I.V.) should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Imipenem and Cilastatin for Injection (I.V.) is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Imipenem and Cilastatin for Injection (I.V.) or other antibacterial drugs in the future.
- Counsel patients to inform their physician:
- if they have central nervous system disorders such as stroke or history of seizures. Seizures have been reported during treatment with Imipenem and Cilastatin for Injection (I.V.) and with closely related antibacterial drugs.
- if they are taking valproic acid or sodium valproate. Valproic acid concentrations in the blood may drop below the therapeutic range upon co-administration with Imipenem and Cilastatin for Injection (I.V.). If treatment with Imipenem and Cilastatin for Injection (I.V.) is necessary and continued, alternative or supplemental anti-convulsant medication to prevent and/or treat seizures may be needed.
- Advise patients that diarrhea is a common problem caused by antibacterial drugs and usually
 resolves when the drug is discontinued. Sometimes, frequent watery or bloody diarrhea may
 occur and may be a sign of a more serious intestinal infection. If severe watery or bloody
 diarrhea develops, patients should contact their healthcare provider.

Manufac	tured for:
\$	FRESENIUS

	I NABI	
_ake	Zurich, IL 60047	

Made in Italy www.fresenius-kabi.com/us

451205D Revised: October 2017

* Quality control ranges for agar dilution testing † Quality control ranges for broth microdilution testing