Naropin® (ropivacaine hydrochloride) Injection, USP

HIGHLIGHTS OF PRESCRIBING INFORMATION These highlights do not include all the information needed to use NAROPIN® safely and effectively. See full prescribing information History of hypersensitivity to local anesthetics of the amide type. (4)

NAROPIN® (ropivacaine hydrochloride) injection, for epidural, rineural, or infiltration use perineural, or minusus... Initial U.S. Approval: 1996 - INDICATIONS AND USAGE -

NAROPIN is an amide local anesthetic indicated in adults for the roduction of local or regional anesthesia for surgery and for acute

Surgical Anesthesia: epidural block for surgery including cesarean ion; major nerve block; local infiltration <u>Acute Pain Management</u>: epidural continuous infusion or intermittent bolus, e.g., postoperative or labor; local infiltration

— DOSAGE AND ADMINISTRATION -

• See Table 1 for Dosage Recommendations (2.2)

---- DOSAGE FORMS AND STRENGTHS --

Injection: 2 mg/mL (0.2%), 5 mg/mL (0.5%), 7.5 mg/mL (0.75%) or 10 mg/mL (1%) in single-dose vials (3)
 Injection: 2 mg/mL (0.2%) or 5 mg/mL (0.5%) in single-dose infusion

- DRUG INTERACTIONS -Agents structurally related to amide-type local anesthetics: Concurrent use may cause additive effects. (7)

Pediatric Use

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action 12.2 Pharmacodynamics 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

Peripheral Nerve Block Local Infiltration

16 HOW SUPPLIED/STORAGE AND HANDLING

14 CLINICAL STUDIES

up to 72 hours.

10 OVERDOSAGE

11 DESCRIPTION

Geriatric Use Hepatic Impairmer Renal Impairment

acidosis, cardiac arrest and, possibly, death, (5.1)

See 17 for PATIENT COUNSELING INFORMATION

— CONTRAINDICATIONS —

- WARNINGS AND PRECAUTIONS -

Delay in proper management of dose-related toxicity, underventilation, and/or altered sensitivity may lead to the development of

actoosis, cardiac arrest and, possibly, death. (5.1)

In performing NAROPIN blocks, unintended intravenous injection is possible and may result in cardiac arrhythmia or cardia arrest. (5.2)

Intra-articular infusions of local anesthetics may cause chondrolysis. NAROPIN is not approved for this use. (5.3).

Signs of methemoglobinemia may occur. (5.4)

----- ADVERSE REACTIONS -

Most common adverse reactions (incidence \geq 5%) are hypotension, nausea, vomiting, bradycardia, fever, pain, postoperative complications, anemia, paresthesia, headache, pruritus, and back pain. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or

Revised: 3/2024

data to support such use. Intravenous regional anesthesia (bie

block) should not be performed due to a lack of clinical experi ence and the risk of attaining toxic blood levels of ropivacaine

It is essential that aspiration for blood, or cerebrospinal fluid (where applicable), be done prior to injecting any local anesthetic, both the original dose and all subsequent doses,

to avoid intravascular or subarachnoid injection. However, a

negative aspiration does not ensure against an intravascular o

arrest. The potential for successful resuscitation has not beer

studied in humans. There have been rare reports of cardiac

arrest during the use of NAROPIN for epidural anesthesia of peripheral nerve blockade, the majority of which occurred after unintentional accidental intravascular administration in elderly patients and in patients with concomitant heart disease. In some instances, resuscitation has been difficult. Should cardiac

arrest occur, prolonged resuscitative efforts may be required to improve the probability of a successful outcome.

NAROPIN should be administered in incremental doses. It is not

recommended for emergency situations, where a fast onset of surgical anesthesia is necessary. Historically, pregnant patients were reported to have a high risk for cardiac arrhythmias, cardiac/circulatory arrest and death when 0,75% bupivacaine

(another member of the amino amide class of local anesthetics) was inadvertently rapidly injected intravenously.

Intra-articular infusions of local anesthetics following arthroscopic and other surgical procedures is an unapproved use, and there have been post-marketing reports of chondrolysis in patients receiving such infusions. The majority of reported

cases of chondrolysis have involved the shoulder joint: cases

of gleno-humeral chondrolysis have been described in pediatric

and adult patients following intra-articular infusions of local anesthetics with and without epinephrine for periods of 48 to 72 hours. There is insufficient information to determine whether

shorter infusion periods are not associated with these findings

The time of onset of symptoms, such as joint pain, stiffness and loss of motion can be variable, but may begin as early as the

2nd month after surgery. Currently, there is no effective treatment for chondrolysis; patients who experienced chondrolysis have required additional diagnostic and therapeutic procedures

Risk of Methemoglobinemia
Cases of methemoglobinemia have been reported in association with local anesthetic use. Although all patients are at risk

tion with local anesthetic use. Although all patients are at risk for methemoglobinemia, patients with glucose-6-phosphate dehydrogenase deficiency, congenital or idiopathic methemoglobinemia, cardiac or pulmonary compromise, infants under 6 months of age, and concurrent exposure to oxidizing agents or their metabolities are more susceptible to developing clinical manifestations of the condition. If local anesthetics must be used in these patients, close monitoring for symptoms and signs of methemoglobinemia is recommended.

Signs of methemoglobinemia may occur immediately or may be delayed some hours after exposure, and are characterized by a cyanotic skin discoloration and/or abnormal coloration of the

blood. Methemoglobin levels may continue to rise; therefore, immediate treatment is required to avert more serious central

nervous system and cardiovascular adverse effects, including seizures, coma, arrhythmias, and death. Discontinue NAROPIN and any other oxidizing agents. Depending on the severity of the

signs and symptoms, patients may respond to supportive care, i.e., oxygen therapy, hydration. A more severe clinical presentation may require treatment with methylene blue, exchange

anesthetic injection. It should be kept in mind at such times

that restlessness, anxiety, incoherent speech, light-headedness, numbness and tingling of the mouth and lips, metallic taste, tinnitus, dizziness, blurred vision, tremors, twitching, depression, or drowsiness may be early warning signs of central nervous

A well-known risk of epidural anesthesia may be an unintentional

A well-known risk of epidural anesthesia may be an unintentional subarachnoid injection of local anesthetic. Two clinical studies have been performed to verify the safety of NAROPIN at a volume of 3 mL injected into the subarachnoid space since this dose represents an incremental epidural volume that could be unintentionally injected. The 15 and 22.5 mg doses injected resulted in sensory levels as high as T5 and T4, respectively. Anesthesia to pinprick started in the sacral dermatomes in 2 to 3 minutes, extended to the T10 level in 10 to 13 minutes and lasted for approximately 2 hours. The results of these two clinical studies showed that a 3 mL dose did not produce any serious adverse events when spinal anesthesia blockade was achieved.

adverse events when spinal anesthesia blockade was achieved

During epidural administration, NAROPIN should be admin-

istered in incremental doses of 3 to 5 mL with sufficient time

between doses to detect toxic manifestations of unintentional intravascular or intrathecal injection. Syringe aspirations should also be performed before and during each supplemental injection.

njection in continuous (intermittent) catheter techniques. In intravascular injection is still possible even if aspirations

for blood are negative. During the administration of epidural

for blood are negative. During the administration of epidural anesthesia, it is recommended that a test dose be administered initially and the effects monitored before the full dose is given. When clinical conditions permit, the test dose should contain an appropriate dose of epinephrine to serve as a warning of unintentional intravascular injection. If injected into a blood vessel, this amount of epinephrine is likely to produce a transient "epinephrine response" within 45 seconds, consisting of an increase in heart rate and systolic blood pressure, circumoral pallor, palpitations and nervousness in the unsedated patient. The sedated patient may exhibit only a pulse rate increase of 20 or more beats per minute for 15 or more seconds. Therefore, following the test dose, the heart should be continuously

fore, following the test dose, the heart should be continuously

monitored for a heart rate increase. Patients on beta-blockers may not manifest changes in heart rate, but blood pressure monitoring can detect a rise in systolic blood pressure. A test dose of a short-acting amide anesthetic such as lidocaine is

recommended to detect an unintentional intrathecal administra-tion. This will be manifested within a few minutes by signs of spinal block (e.g., decreased sensation of the buttocks, paresis of the legs, or, in the sedated patient, absent knee jerk). An intravascular or subarachnoid injection is still possible even if results of the test dose are negative. The test dose itself may produce a systemic toxic reaction, high spinal or epinephrine-induced cardiovascular effects.

Ropivacaine plasma concentrations may approach the threshold for central nervous system toxicity after the administration of 300 mg of ropivacaine for brachial plexus block. Caution should be exercised when using the 300 mg dose [see Overdosage (10)].

The dose for a major nerve block must be adjusted according

to the site of administration and patient status. Supracla-vicular brachial plexus blocks may be associated with a higher

frequency of serious adverse reactions, regardless of the local

wald peripried in the blocks may result in the administration of a large volume of local anesthetic in highly vascularized areas, often close to large vessels where there is an increased risk of intravascular injection and/or rapid systemic absorption, which

heral nerve blocks may result in the administration of

5.5 Central Nervous System Toxicity Careful and constant monitoring of cardiovascular and respiratory vital signs (adequacy of ventilation) and the patient's state of consciousness should be performed after each local

system toxicity.

5.6 Epidural Anesthesia

and some required arthroplasty or shoulder replacement.

5.3 Intra-Articular Infusions and Risk of Chondrolysis

5.2 Unintended Intravenous Injection
In performing NAROPIN blocks, unintended intravenous injection is possible and may result in cardiac arrhythmia or cardiac

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE 2 DOSAGE AND ADMINISTRATION

Important Administration Instructions Dosage Recommendations Other Administration Considerations

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

General Warnings and Precautions Unintended Intravenous Injection Intra-Articular Infusions and Risk of Chondrolysis

Risk of Methemoglobinemia Central Nervous System Toxicity Epidural Anesthesia Use in Brachial Plexus Block Use in Peripheral Nerve Block Use in Head and Neck Area

5.9 Use in Plead and Neck Area
5.10 Use in Ophthalmic Surgery
5.11 Hepatic Disease
5.12 Cardiovascular Impairment
5.13 Risk of Additive Effects
5.14 Malignant Hyperthermia

6 ADVERSE REACTIONS 7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy 8.2 Lactation

17 PATIENT COUNSELING INFORMATION *Sections or subsections omitted from the full prescribing information

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility 13.2 Animal Toxicology and/or Pharmacology

Epidural Administration in Surgery
Epidural Administration in Cesarean Section

Epidural Administration in Labor and Delivery Epidural Administration in Postoperative Pain Management

FULL PRESCRIBING INFORMATION INDICATIONS AND USAGE

NAROPIN is indicated for the production of local or regional

anesthesia for surgery and for acute pain managen <u>Surgical Anesthesia</u>: epidural block for surgery including cesarean section; major nerve block; local infiltration

Acute Pain Management: epidural continuous infusion or interittent bolus, e.g., postoperative or labor; local infiltration

DOSAGE AND ADMINISTRATION 2.1 Important Administration Instructions

There have been adverse event reports of chondrolysis in patients receiving intra-articular infusions of local anesthetics following arthroscopic and other surgical procedures. NAROPIN is not approved for this use [see Warnings and Precautions (5.3)].

The rapid injection of a large volume of local anesthetic solution should be avoided and fractional (incremental) doses should always be used. The smallest dose and concentration required to produce the desired result should be administered.

The dose of any local anesthetic administered varies with the anesthetic procedure, the area to be anesthetized, the vascularity of the tissues, the number of neuronal segments to be blocked, the depth of anesthesia and degree of muscle relaxto be blocked, ine depin of an esthesia at a degree of muscale relaxation required, the duration of an esthesia desired, individual tolerance, and the physical condition of the patient. Patients in poor general condition due to aging or other compromising factors such as partial or complete heart conduction block, advanced liver disease or severe renal dysfunction require special attention although regional anesthesia is frequently indicated in these patients. To reduce the risk of potentially serious advance reactions attempts should be made to optimize serious adverse reactions, attempts should be made to optimiz the patient's condition before major blocks are performed, and

Use an adequate test dose (3 to 5 mL of a short acting local anesthetic solution containing epinephrine) prior to induction of complete block. This test dose should be repeated if the patient is moved in such a fashion as to have displaced the epidural catheter. Allow adequate time for onset of anesthesia following administration of each test dose.

These products are intended for single dose and are free from reservatives. Any solution remaining from an opened container hould be discarded promptly. In addition, continuous infusion ottles should not be left in place for more than 24 hours.

2.2 Dosage Recommendations

Rx only

Table 1
Dosage Recommendations

	Co mg/mL	nc. (%)	Volume mL	Dose mg	Onset min	Duration hours
SURGICAL ANESTHESIA						
Lumbar Epidural	5	(0.5%)	15 to 30	75 to 150	15 to 30	2 to 4
Administration	7.5	(0.75%)	15 to 25	113 to 188	10 to 20	3 to 5
Surgery	10	(1%)	15 to 20	150 to 200	10 to 20	4 to 6
Lumbar Epidural	5	(0.5%)	20 to 30	100 to 150	15 to 25	2 to 4
Administration	7.5	(0.75%)	15 to 20	113 to 150	10 to 20	3 to 5
Cesarean Section						
Thoracic Epidural	5	(0.5%)	5 to 15	25 to 75	10 to 20	n/a*
Administration	7.5	(0.75%)	5 to 15	38 to 113	10 to 20	n/a*
Surgery						
Major Nerve Block†	5	(0.5%)	35 to 50	175 to 250	15 to 30	5 to 8
(e.g., brachial plexus block)	7.5	(0.75%)	10 to 40	75 to 300	10 to 25	6 to 10
Field Block	5	(0.5%)	1 to 40	5 to 200	1 to 15	2 to 6
(e.g., minor nerve blocks and	infiltration	1)				
LABOR PAIN MANAGEMENT						
Lumbar Epidural Administra	tion					
Initial Dose	2	(0.2%)	10 to 20	20 to 40	10 to 15	0.5 to 1.5
Continuous infusion‡	2	(0.2%)	6 to 14 mL/h	12 to 28 mg/h	n/a*	n/a*
Incremental injections (top-up) [‡]	2	(0.2%)	10 to 15 mL/h	20 to 30 mg/h	n/a*	n/a*
POSTOPERATIVE PAIN MANA	AGEMENT					
Lumbar Epidural Administra	tion					
Continuous infusion§	2	(0.2%)	6 to 14 mL/h	12 to 28 mg/h	n/a*	n/a*
Thoracic Epidural	2	(0.2%)	6 to 14 mL/h	12 to 28 mg/h	n/a*	n/a*
Administration						

* = Not Applicable † = The dose for a major nerve block must be adjusted according to site of administratio and patient status. Supraclavicular brachial plexus blocks may be associated with and patient status. Supraclavicular brachial plexus blocks may be associated with a higher frequency of serious adverse reactions, regardless of the local anesthetic used [see Warnings and Precautions (5.7)].

1 Median dose of 21 mg per hour was administered by continuous infusion or by incremental injections (top-ups) over a median delivery time of 5.5 hours.

5 Cumulative doses up to 770 mg of NAROPIN over 24 hours (Intraoperative block plus postoperative infusion); Continuous epidural infusion at rates up to 28 mg per hour for

2 (0.2%) 1 to 100 2 to 200 1 to 5 2 to 6

5 (0.5%) 1 to 40 5 to 200 1 to 5 2 to 6

ostoperative infusion); Continuous epidural infusion at rates up و حوالية المراقعة المراقعة

The doses in the table are those considered to be necessary to produce a successful block and should be regarded as guidelines for use in adults. Individual variations in onset and duration occur. The figures reflect the expected average dose range needed. For other local anesthetic techniques standard current textbooks should be consulted.

When prolonged blocks are used, either through continuous infusion or through repeated bolus administration, the risk reaching a toxic plasma concentration or inducing local ne injury must be considered. Experience to date indicates that a cumulative dose of up to 770 mg NAROPIN administered over 24 hours is well tolerated in adults when used for postoperative pain management: i.e., 2016 mg. Caution should be exercised when administering NAROPIN for prolonged periods of time, e.g., >70 hours in debilitated patients. For treatment of postoperative pain, the following technique For treatment of postoperative pain, the following technique can be recommended: If regional anesthesia was not used intraoperatively, then an initial epidural block with 5 to 7 mL NAROPIN is induced via an epidural catheter. Analgesia is maintained with an infusion of NAROPIN, 2 mg/mL (0.2%). Clinical studies have demonstrated that infusion rates of 6 to 14 mL (12 to 28 mg) per hour provide adequate analgesia with nonprogressive motor block. With this technique a significant reduction in the need for opioids was demonstrated. Clinical experience supports the use of NAROPIN epidural infusions for

2.3 Other Administration Considerations Disinfecting agents containing heavy metals, which cause release of respective ions (mercury, zinc, copper, etc.) should not be used for skin or mucous membrane disinfection since they have been related to incidents of swelling and edema.

When chemical disinfection of the container surface is desired, either isopropyl alcohol (91%) or ethyl alcohol (70%) is recon either isopropyl alcohol (91%) or ethyl alcohol (70%) is recom-mended. It is recommended that chemical disinfection be accomplished by wiping the polypropylene ampule or vial stopper thoroughly with cotton or gauze that has been moist-ened with the recommended alcohol just prior to use. When a container is required to have a sterile outside, a Sterile-Pak

should be chosen. Glass containers may, as an alternative be autoclaved once. Stability has been demonstrated using a targeted F_0 of 7 minutes at 121°C. The solubility of ropivacaine is limited at pH above 6. Thus care must be taken as precipitation may occur if NAROPIN is mixed with alkaline solutions. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Solutions which are discolored or which contain particulate

DOSAGE FORMS AND STRENGTHS NAROPIN® (ropivacaine hydrochloride) injection is a clear, colorless, preservative-free solution available as:

matter should not be administered

NAROPIN® Single Dose Infusion Bottles

0.2%, 200 mg per 100 mL (2 mg/mL), 100 mL single-dose infusion bottle 0.2%, 400 mg per 200 mL (2 mg/mL), 200 mL single-dose

• 0.5%, 500 mg per 100 mL (5 mg/mL), 100 mL single-dose • 0.5%, 1,000 mg per 200 mL (5 mg/mL), 200 mL single-dose

CONTRAINDICATIONS

AROPIN is contraindicated in patients with a known hyper sensitivity to ropivacaine or to any local anesthetic agent of the

WARNINGS AND PRECAUTIONS

General Warnings and Precautions General Warnings and Precautions
Prior to receiving major blocks the general condition of the
patient should be optimized and the patient should have an
IV line inserted. All necessary precautions should be taken to
avoid intravascular injection. Local anesthetics should only be
administered by clinicians who are well versed in the diagnosis
and management of does related twicity and other actus memory. and management of dose-related toxicity and other acute emer-gencies which might arise from the block to be employed, and hen only after insuring the immediate (without delay) availresuscitative equipment, and the personnel resources needed for proper management of toxic reactions and related emergencies (see Adverse Reactions (6) and Overdosage (10.1)]. Delay in proper management of dose-related toxicity, underventilation from any cause, and/or altered sensitivity may lead to the development of acidosis, cardiac arrest and, possibly, death The safe and effective use of local anesthetics depends on

dose because circulatory changes compatible with epinephrine

may also serve as a warning sign of unintended intravascular

injection. An intravascular injection is still possible even if aspirations for blood are negative. Administration of higher than recommended doses of NAROPIN to achieve greater motor blockade or increased duration of sensory blockade may

esult in cardiovascular depression, particularly in the event

of inadvertent intravascular injection. Tolerance to elevated

Debilitated, elderly patients and acutely ill patients should be given reduced doses commensurate with their age and physical condition. Local anesthetics should also be used with caution

Solutions of NAROPIN should not be used for the production

of obstetrical paracervical block anesthesia, retrobulbar block or spinal anesthesia (subarachnoid block) due to insufficien

plood levels varies with the physical condition of the patient

in patients with hypotension, hypovolemia or heart block.

Use in Head and Neck Area mall doses of local anesthetics injected into the head and proper dosage, correct technique, adequate precautions and readiness for emergencies. neck area may produce adverse reactions similar to systemic toxicity seen with unintentional intravascular injections of larger doses. The injection procedures require the utmost care. Confusion, convulsions, respiratory depression, and/or respiratory arrest, and cardiovascular stimulation or depression have been reported. These reactions may be due to intra-arterial injection of Resuscitative equipment, oxygen and other resuscitative drugs should be available for immediate use [see Adverse Reactions (6)]. The lowest dosage that results in effective anesthesia should be used to avoid high plasma levels and serious adverse events. Injections should be made slowly and incrementally, with the local anesthetic with retrograde flow to the cerebral circula-tion. Patients receiving these blocks should have their circulation and respiration monitored and be constantly observed. Resus-citative equipment and personnel for treating adverse reactions obsuid by immediately usualishs. Decara recommendations requent aspirations before and during the injection to avoid ntravascular injection. When a continuous catheter technique is used, syringe aspirations should also be performed before and during each supplemental injection. During the administration of epidural anesthesia, it is recommended that a test dose of a local anesthetic with a fast onset be administered initially should be immediately available. Dosage recommendations should not be exceeded [see Dosage and Administration (2.2)]. 5.10 Use in Ophthalmic Surgery
The use of NAROPIN in retrobulbar blocks for ophthalmic surgery and that the patient be monitored for central nervous system and cardiovascular toxicity, as well as for signs of unintended has not been studied. Until appropriate experience is gained the use of NAROPIN for such surgery is not recommended. and cardiovascual toxicity, as well as for signs of difficultied intrathecal administration before proceeding. When clinical conditions permit, consideration should be given to employing local anesthetic solutions, which contain epinephrine for the test

nduced cardiovascular effects.

5.7 Use in Brachial Plexus Block

5.8 Use in Peripheral Nerve Block

can lead to high plasma concentrations.

5.11 Hepatic Disease
Because amide-type local anesthetics such as ropivacaine are

metabolized by the liver, these drugs, especially repeat doses, should be used cautiously in patients with hepatic disease. Patients with severe hepatic disease, because of their inability to metabolize local anesthetics normally, are at a greater risk of developing toxic plasma concentrations. 5.12 Cardiovascular Impairment

Local anesthetics should also be used with caution in patients with impaired cardiovascular function because they may be less able to compensate for functional changes associated with the prolongation of A-V conduction produced by these drugs. 5.13 Risk of Additive Effects
NAROPIN should be used with caution in patients receiving other local anesthetics or agents structurally related to amide-type local anesthetics, since the toxic effects of these drugs are additive. [See Drug Interactions 7.0]

5.14 Malignant Hyperthermia
Many drugs used during the conduct of anesthesia are consid

wany drugs used using the conduct of ariestics are considered potential triggering agents for malignant hyperthermia (MH). Amide-type local anesthetics are not known to trigger this reaction. However, since the need for supplemental general anesthesia cannot be predicted in advance, it is suggested that a standard protocol for MH management should be available. ADVERSE REACTIONS Reactions to ropivacaine are characteristic of those associated with other amide-type local anesthetics. A major cause of

nteractions 7.01

Patients treated with class III antiarrhythmic drugs (e.g., amiodarone) should be under close surveillance and ECG monitoring considered, since cardiac effects may be additive. [See Drug lettered]

adverse reactions to this group of drugs may be associated with excessive plasma levels, which may be due to overdosage, unin-tentional intravascular injection or slow metabolic degradation. The reported adverse events are derived from clinical studies conducted in the U.S. and other countries. The reference drug was usually bupivacaine. The studies used a variety of premedications, sedatives, and surgical procedures of varying length. A total of 3,988 patients have been exposed to NAROPIN at concentrations up to 1% in clinical trials. Each patient was counted once for each type of adverse event. Because clinical trials are conducted under widely conditions, adverse reactions rates observed in the clinical trials of a drug

cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice. Incidence ≥ 5%

For the indications of epidural administration in surgery, cesarean section, postoperative pain management, peripheral nerve block, and local infiltration, the following treatment-emergent adverse events were reported with an incidence of ≥ 5% in all clinical studies (N=3988): hypotension (37%), nausea (24.8%), vomiting (11.6%), bradycardia (9.3%), fever (9.2%), pain (8%), postoperative complications (7.1%), anemia (6.1%), paresthesia (5.6%), headache (5.1%), pruritus (5.1%), and back pain (5%). ncidence 1 to 5%

Urinary retention, dizziness, rigors, hypertension, tachycardia, anxiety, oliguria, hypoesthesia, chest pain, hypokalemia, dyspnea, cramps, and urinary tract infection. Incidence in Controlled Clinical Trials

The reported adverse events are derived from controlled clinical studies with NAROPIN (concentrations ranged from 0.125% to 1% for NAROPIN and 0.25% to 0.75% for bupivacaine) in the U.S. and other countries involving 3,094 patients. Table 2 and Table 3 list adverse events (number and percentage) that occurred in at least 1% of NAROPIN-treated patients in these studies. The majority of patients receiving concentrations highe than 5 mg/mL (0.5%) were treated with NAROPIN. Table 2

Adverse Events Reported in ≥1% of Adult Patients Receiving Regional or Local Anesthesia (Surgery, Labor, Cesarean Section, Postoperative Pain Management, Peripheral Nerve Block and Local Infiltration)

	NAR	OPIN	Bupivacaine		
	total N	=1661	total N=1433		
Adverse Reaction	N	%	N	%	
Hypotension	536	(32.3)	408	(28.5)	
Nausea	283	(17)	207	(14.4)	
Vomiting	117	(7)	88	(6.1)	
Bradycardia	96	(5.8)	73	(5.1)	
Headache	84	(5.1)	68	(4.7)	
Paresthesia	82	(4.9)	57	(4)	
Back pain	73	(4.4)	75	(5.2)	
Pain	71	(4.3)	71	(5)	
Pruritus	63	(3.8)	40	(2.8)	
Fever	61	(3.7)	37	(2.6)	
Dizziness	42	(2.5)	23	(1.6)	
Rigors (Chills)	42	(2.5)	24	(1.7)	
Postoperative complications	41	(2.5)	44	(3.1)	
Hypoesthesia	27	(1.6)	24	(1.7)	
Urinary retention	23	(1.4)	20	(1.4)	
Progression of labor poor/failed	23	(1.4)	22	(1.5)	
Anxiety	21	(1.3)	11	(0.8)	
Breast disorder, breast-feeding	21	(1.3)	12	(0.8)	
Rhinitis	18	(1.1)	13	(0.9)	

Table 3 Adverse Events Reported in ≥1% of Fetuses or

Neonates of Mothers Who Received Regional Anesthesia (Cesarean Section and Labor Studies)										
	NAF	ROPIN	Bupiv	racaine						
	total	N=639	total	N=573						
Adverse Reaction	N	%	N	%						
Fetal bradycardia	77	(12.1)	68	(11.9)						
Neonatal jaundice	49	(7.7)	47	(8.2)						
Neonatal complication-NOS	42	(6.6)	38	(6.6)						
Apgar score low	18	(2.8)	14	(2.4)						
Neonatal respiratory disorder	17	(2.7)	18	(3.1)						
Neonatal tachypnea	14	(2.2)	15	(2.6)						
Neonatal fever	13	(2)	14	(2.4)						
Fetal tachycardia	13	(2)	12	(2.1)						
Fetal distress	11	(1.7)	10	(1.7)						
Neonatal infection	10	(1.6)	8	(1.4)						
Neonatal hypoglycemia	8	(1.3)	16	(2.8)						

Incidence <1%
The following adverse events were reported during the NAROPIN clinical program in more than one patient (N=3988), occurred at an overall incidence of <1%, and were considered relevant: Application Site Reactions - injection site pain

Cardiovascular System - vasovagal reaction, syncope, postural hypotension, non-specific ECG abnormalities Female Reproductive - poor progression of labor, uterine atony

Gastrointestinal System - fecal incontinence, tenesmus, neonatal General and Other Disorders - hypothermia, malaise, asthenia,

Hearing and Vestibular - tinnitus, hearing abnormalities

Heart Rate and Rhythm - extrasystoles, non-specific arrhythmias

Liver and Biliary System - jaundice Metabolic Disorders - hypomagnesemia

Musculoskeletal System - myalgia Myo/Endo/Pericardium - ST segment changes, myocardial

Nervous System - tremor, Horner's syndrome, paresis, dyski hypotonia, ptosis, stupor

Psychiatric Disorders - agitation, confusion, somnolence, nervousness, amnesia, hallucination, emotional lability, Respiratory System - bronchospasm, coughing

Skin Disorders - rash, urticaria Urinary System Disorders - urinary incontinence, micturition

Vascular - deep vein thrombosis, phlebitis, pulmonary embolism Vision - vision abnormalities For the indication epidural anesthesia for surgery, the 15 most common adverse events were compared between different concentrations of NAROPIN and bupivacaine. Table 4 is based

NAROPIN was administered as an epidural anesthetic for Table 4
Common Events (Epidural Administration)

n data from trials in the U.S. and other countries where

NAROPIN Bupivacaine

	5 m	g/mL	7.5 n	ng/mL	10 m	ng/mL	5 m	g/mL	7.5 n	ng/mL
Adverse	total	N=256	total N=297		total N=207		total N=236		total N=174	
Reaction	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)
hypotension	99	(38.7)	146	(49.2)	113	(54.6)	91	(38.6)	89	(51.1)
nausea	34	(13.3)	68	(22.9)			41	(17.4)	36	(20.7)
bradycardia	29	(11.3)	58	(19.5)	40	(19.3)	32	(13.6)	25	(14.4)
back pain	18	(7)	23	(7.7)	34	(16.4)	21	(8.9)	23	(13.2)
vomiting	18	(7)	33	(11.1)	23	(11.1)	19	(8.1)	14	(8)
headache	12	(4.7)	20	(6.7)	16	(7.7)	13	(5.5)	9	(5.2)
fever	8	(3.1)	5	(1.7)	18	(8.7)	11	(4.7)		
chills	6	(2.3)	7	(2.4)	6	(2.9)	4	(1.7)	3	(1.7)
urinary retention	5	(2)	8	(2.7)	10	(4.8)	10	(4.2)		
paresthesia	5	(2)	10	(3.4)	5	(2.4)	7	(3)		
nn iritus			14	(4.7)	3	(1.4)			7	(4)

Using data from the same studies, the number (%) of patients iencing hypotension is displayed by patient age, drug oncentration in Table 5. In Table 6, the adverse events for NAROPIN are broken down by gender.

Table 5 Effects of Age on Hypotension (Epidural Administration) Total N: NAROPIN = 760, Bupivacaine = 410

	NAROPIN							Bupiva	caine	
GE	5 mg/mL		7.5 r	ng/mL	10 n	ng/mL	5 m	g/mL	7.5 r	ng/mL
	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)
65	68	(32.2)	99	(43.2)	87	(51.5)	64	(33.5)	73	(48.3)
65	31	(68.9)	47	(69.1)	26	(68.4)	27	(60)	16	(69.6)
Table 6_										

Most Common Adverse Events by G Total N: Fer

Adverse Reaction	Fer	nale	M	ale
Adverse Reaction	N	(%)	N	(%)
hypotension	220	(54.3)	138	(38.9)
nausea	119	(29.4)	23	(6.5)
bradycardia	65	(16)	56	(15.8)
vomiting	59	(14.6)	8	(2.3)
back pain	41	(10.1)	23	(6.5)
headache	33	(8.1)	17	(4.8)
chills	18	(4.4)	5	(1.4)
fever	16	(4)	3	(0.8)
pruritus	16	(4)	1	(0.3)
pain	12	(3)	4	(1.1)
urinary retention	11	(2.7)	7	(2)
dizziness	9	(2.2)	4	(1.1)
hypoesthesia	8	(2)	2	(0.6)
paresthesia	8	(2)	10	(2.8)
Systemic Reactions				

The most commonly encountered acute adverse experiences that demand immediate countermeasures are related to the central nervous system and the cardiovascular system. These adverse experiences are generally dose-related and due to high plasma levels that may result from overdosage, rapid absorption from the injection site, diminished tolerance or from unintentional intravascular injection of the local anesthetic solu uninterluoria intravascular injection of the local anesthetic solu-tion. In addition to systemic dose-related toxicity, unintentional subarachnoid injection of drug during the intended performance of lumbar epidural block or nerve blocks near the vertebral column (especially in the head and neck region) may result in underventilation or apnea ("Total or High Spinal"). Also, hypotension due to loss of sympathetic tone and resp nypotension due to loss of sympathetic tone and respiratory paralysis or underventilation due to cephalad extension of the motor level of anesthesia may occur. This may lead to secondary cardiac arrest if untreated. Factors influencing plasma protein binding, such as acidosis, systemic diseases that alter protein production or competition with other drugs for protein binding sites, may diminish individual tolerance.

Epidural administration of NAROPIN has, in some cases, as with other local anesthetics, been associated with transient increases in temperature to > 38.5°C. This occurred more frequently at doses of NAROPIN > 16 mg/h.

Neurologic Reactions These are characterized by excitation and/or depression. Restlessness, anxiety, dizziness, tinnitus, blurred vision or tremors may occur, possibly proceeding to convulsions. However, excitement may be transient or absent, with depression being the first manifestation of an adverse reaction. This may quickly

be followed by drowsiness merging into unconsciousness and respiratory arrest. Other central nervous system effects may be nausea, vomiting, chills, and constriction of the pupils. The incidence of convulsions associated with the use of local anesthetics varies with the route of administration and the total dose administered. In a survey of studies of epidural anesthesia,

overt toxicity progressing to convulsions occurred in approxi-mately 0.1% of local anesthetic administrations.

The incidence of adverse neurological reactions associated with the use of local anesthetics may be related to the total dose and concentration of local anesthetic administered and are also dependent upon the particular drug used, the route of administration, and the physical status of the patient. Many of these observations may be related to local anesthetic techniques, with or without a contribution from the drug. During lumbar epidural block, occasional unintentional penetration of the subarachnoid space by the catheter or needle may occur. Subsequent adverse effects may depend partially on the amount of drug administered intrathecally as well as the physiological and physical effects of a dural puncture. These observations may include spinal block of varying magnitude (including high or total spinal block), hypotension secondary to spinal block, urinary retention, loss of bladder and bowel control (fecal and urinary incontinence), and loss of perineal sensation and sexual function. Signs and symptoms of subarachnoid block typically start within 2 to 3 migrates of the spinal sense of 15 and 22 5 mg. start within 2 to 3 minutes of injection. Doses of 15 and 22.5 mg of NAROPIN resulted in sensory levels as high as T5 and T4 respectively. Analgesia started in the sacral dermatomes in 2 to 3 minutes and extended to the T10 level in 10 to 13 minutes and lasted for approximately 2 hours. Other neurological effects following unintentional subarachnoid administration during epidural anesthesia may include persistent anesthesia, pares-thesia, weakness, paralysis of the lower extremities, and loss of sphincter control; all of which may have slow, incomplete or no recovery. Headache, septic meningitis, meningismus, slowing of labor, increased incidence of forceps delivery, or cranial nerve palsies due to traction on nerves from loss of cerebrospinal fluid have been reported [see Dosage and Administration (2.1)]. A high spinal is characterized by paralysis of the arms, loss of consciousness, respiratory paralysis and bradycardia.

Cardiovascular System Reactions

High doses or unintentional intravascular injection may lead to high plasma levels and related depression of the myocardium, decreased cardiac output, heart block, hypotension, bradycardia, ventricular arrhythmias, including ventricular tachycardia and ventricular fibrillation, and possibly cardiac arrest [see Warnings and Precautions (5.2) and Overdosage (10)].

Allergic Reactions
Allergic type reactions are rare and may occur as a result of sensitivity to the local anesthetic [see Warnings and Precautions (5.1)]. These reactions are characterized by signs such as urticaria, pruritus, erythema, angioneurotic edema (including laryngeal edema), tachycardia, sneezing, nausea, vomiting, dizziness, syncope, excessive sweating, elevated temperature, and possibly, anaphylactoid symptomatology (including severe hypotension). Cross-sensitivity among members of the amidetype local anesthetic group has been reported. The usefulness of screening for sensitivity has not been definitively established. DRUG INTERACTIONS

Patients who are administered local anesthetics are at increased risk of developing methemoglobinemia when concurrently exposed to the following drugs, which could include other local anesthetics [see Warnings and Precautions (5.4)]: Examples of Drugs Associated with Mether

Class	Examples				
Nitrates/Nitrites	nitric oxide, nitroglycerin, nitroprusside, nitrous oxide				
Local anesthetics	articaine, benzocaine, bupivacaine, lidocaine, mepivacaine, prilocaine, procaine, ropivacaine, tetracaine				
Antineoplastic agents cyclophosphamide, flutamide, hydroxyurea, ifosfamide, rasburicase					
Antibiotics	dapsone, nitrofurantoin, para-aminosalicylic acid, sulfonamides				
Antimalarials	chloroquine, primaquine				
Anticonvulsants	Phenobarbital, phenytoin, sodium valproate				
Other drugs	acetaminophen, metoclopramide, quinine, sulfasalazine				
NAROPIN should be used with caution in patients receiving					

other local anesthetics or agents structurally related to amidetype local anesthetics, since the toxic effects of these drugs are additive. Cytochrome P4501A2 is involved in the formation of 3-hydroxy ropivacaine, the major metabolite. In vivo, the plasma clearance of ropivacaine was reduced by 70% during coadministration of fluvoxamine (25 mg bid for 2 days), a selective and potent CYP1A2 inhibitor. Thus strong inhibitors of cytochrome P4501A2, such as fluvoxamine, given concomitantly during administration of NAROPIN, can interact with NAROPIN leading to increased ropivacaine plasma levels. Caution should be exercised when CYP1A2 inhibitors are coadministered. Possible interactions with drugs known to be metabolized by CYP1A2 via competitive inhibition such as theophylline and imipramine may also occur. Coadministration of a selective and potent inhibitor of CYP3A4, ketoconazole (100 mg bid for 2 days with ropivacaine infusion administered 1 hour after ketoconazole) caused a 15% reduction in in vivo plasma clearance of ropivacaine. Specific trials studying the interaction between ropivacaine

and class III antiarrhythmic drugs (e.g., amiodarone) have not been performed, but caution is advised [see Warnings and Precautions (5.13)]. 8 USE IN SPECIFIC POPULATIONS

There are no available human data on use of Naropin (ropiya-

8.1 Pregnancy

Risk Summary

fural Administration) males = 405, Males = 355									
Female Male									
	N	(%)	N	(%)					
	220	(54.3)	138	(38.9)					
	119	(29.4)	23	(6.5)					
	65	(16)	56	(15.8)					
	59	(14.6)	8	(2.3)					

cardiac function. Maternal Adverse reactions

Clinical Considerations

Labor or Delivery

Maternal hypotension has resulted from regional anesthesia. Local anesthetics produce vasodilation by blocking sympathetic nerves. Therefore, during treatment of systemic toxicity, materna hypotension or fetal bradycardia following regional block, the parturient should be maintained in the left lateral decubitus position if possible, or manual displacement of the uterus off the great vessels be accomplished. Elevating the patient's legs will also help prevent decreases in blood pressure. The fetal heart rate also should be monitored continuously, and electronic fetal monitoring in blobby to displace. nonitoring is highly advisable.

maternal or fetal outcomes. Local anesthetics may cause

varying degrees of toxicity to the mother and fetus and adverse reactions include alterations of the central nervous system peripheral vascular tone, and cardiac function (see Clinica

Considerations). No teratogenicity was observed at doses up to 0.3 times the maximum recommended human dose of 770 mg/24 hours for epidural use, and equal to the MRHD of

250 mg for nerve block use, based on body surface area (BSA) comparisons and a 60 kg human weight (see Animal data).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U. S. general population, the estimated background risk of major birth defects and miscar-

riage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Local anesthetics, including ropivacaine, rapidly cross the placenta, and when used for epidural block can cause varying

degrees of maternal, fetal, and neonatal toxicity [see Clinical Pharmacology (12)]. The incidence and degree of toxicity

depend upon the procedure performed, the type and amount

of drug used, and the technique of drug administration. Adverse reactions in the parturient, fetus and neonate involve alterations of the central nervous system, peripheral vascular tone and

No malformations were reported in embryo-fetal development toxicity studies conducted in pregnant New Zealand white rabbits and Sprague-Dawley rats. During gestation days 6 to rabbits received daily subcutaneous doses of ropivacain at 1.3, 4.2, or 13 mg/kg/day (equivalent to 0.03, 0.10, and 0.33 times the maximum recommended human dose (MRHD) of 770 mg/24 hours, respectively, and 0.10, 0.32, and 1.0 times the MRHD of 250 mg for nerve block use, respectively based on body surface area (BSA) comparisons and a 60 kg human weight). Rats received daily subcutaneous doses of 5.3, 11, and 26 mg/kg/day (equivalent to 0.07, 0.14, and 0.33 times the MRHD for epidural use, respectively, and 0.21, 0.43, and 1.0 times the MRHD for nerve block use, respectively, based on BSA comparisons) during GD 6 to 15. No treatment-related effects on late fetal development, partur

tion, litter size, lactation, neonatal viability, or growth of the offspring were reported in a prenatal and postnatal reproductive onspring were reported in a prenatial and postnatial reproductive and development toxicity study; however functional endpoints were not evaluated. Female rats were dosed daily subcutaneously from GD 15 to Lactation Day 20 at doses of 5.3, 11, and 26 mg/kg/day (equivalent to 0.07, 0.1, and 0.3 times the MRHD for epidural use, respectively, and 0.21, 0.43, and 1.0 times the MRHD for nerve block use, respectively), with maternal toxicity exhibited at the high dose. No adverse effects in physical developmental milestones or in

behavioral tests were reported in a 2-generational reprodustudy, in which rats received daily subcutaneous doses of 12, and 23 mg/kg/day (equivalent to 0.08, 0.15, and 0.29 to 1.00 t the MRHD for epidural use, respectively, and 0.24, 0.45, and 0.88 times the MRHD for nerve block use, respectively, based or 0.88 times the MIHID for nerve block use, respectively, based on BSA comparisons) for 9 weeks before mating and during mating for males, and for 2 weeks before mating and during mating, pregnancy, and lactation, up to day 42 post coitus for females. Significant pup loss was observed in the high dose group during the first 3 days postpartum, from a few hours up to 3 days after delivery compared to the control group, which was considered secondary to impaired maternal care due to maternal toxicity. No differences were observed in littler parameters, or fertility, mean gestation time, or number of live births were observed between the control (saline) and treatment groups [see Carcinogenesis, Mutagenesis, Impairment of Fertility (13.1)].

8.2 Lactation Risk Summary

One publication reported that ropivacaine is present in human milk at low levels following administration of ropivacaine in women undergoing cesarean section. No adverse reactions were reported in the infants. There is no available information on the drug's effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for NAROPIN and any potential adverse effects on the breastfed child from NAROPÍN or from

Pediatric Use The safety and efficacy of NAROPIN in pediatric patients have not been established

Geriatric Use
Of the 2,978 subjects that were administered NAROPIN Injection in of the 2,976 subjects that were administrated NAM-PIN injection in 71 controlled and uncontrolled clinical studies, 803 patients (27%) were 65 years of age or older which includes 127 patients (4%) 75 years of age and over. NAROPIN Injection was found to be safe and effective in the patients in these studies. Clinical data in one published article indicate that differences in various pharmacodynamic measures were observed with increasing age. In one study, the upper level of analgesia increased with age, the maximum decrease of mean arterial pressure (MAP) declined with age during the first hour after epidural administration, and the intensity of motor blockade increased with age. This drug and its metabolites are known to be excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Elderly patients are more likely to have decreased hepatic, renal, or cardiac function, as well as concomitant disease. Therefore, care should be taken in dose selection, starting at the low end of the dosage range, and it may be useful to monitor renal function [see Clinical Pharmacology (12.3)] Pharmacology (12.3)].

Hepatic Impairment Because amide-type local anesthetics such as ropivacaine are metabolized by the liver, these drugs, especially repeat doses, should be used cautiously in patients with hepatic disease. Patients with severe hepatic disease, because of their inability to metabolize local anesthetics normally, are at a greater risk of developing toxic plasma concentrations [see Warning and Preparties [5.11]].

This drug and its metabolites are known to be excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Therefore, care should be taken in dose selection, starting at the low end of the dosage range, and it may be useful to monitor renal function [see Clinical Pharmacology (12.3)].

Renal Impairment

OVERDOSAGE acute emergencies from local anesthetics are generally related to high plasma levels encountered, or large doses administered, during therapeutic use of local anesthetics or to unintended subarachnoid or intravascular injection of local anesthetic soluion [see Adverse Reactions (6) and Warnings and Precautions

10.1 Treatment
Therapy with NAROPIN should be discontinued at the first sign of toxicity. No specific information is available for the treatment of toxicity with NAROPIN; therefore, treatment should be sympomatic and supportive. The first consideration is preven tomatic and supportive. The lifst consideration is prevention, best accomplished by incremental injection of NAROPIN, careful and constant monitoring of cardiovascular and respiratory vital signs and the patient's state of consciousness after each local anesthetic and during continuous infusion. At the first sign of change in mental status, oxygen should be administered.

The first step in the management of systemic toxic reactions, as well as underventiation or apnea due to unintentional subarach-noid injection of drug solution, consists of immediate attention to the establishment and maintenance of a patent airway and effective assisted or controlled ventilation with 100% oxygen with a delivery system capable of permitting immediate positive airway pressure by mask. Circulation should be assisted as necessary. This may prevent convulsions if they have not already occurred

If necessary, use drugs to control convulsions. Intravenou barbiturates, anticonvulsant agents, or muscle relaxants should only be administered by those familiar with their use. Immediately after the institution of these ventilatory measures, the adequacy of the circulation should be evaluated. Supportive treatment of circulatory depression may require administration of intravenous fluids, and, when appropriate, a vasopressol dictated by the clinical situation (such as ephedrine or epinephrine to enhance myocardial contractile force).

Should cardiac arrest occur, prolonged resuscitative efforts may be required to improve the probability of a successful outcome. The mean dosages of ropivacaine producing seizures, after intravenous infusion in dogs, nonpregnant and pregnant sheep were 4.9, 6.1 and 5.9 mg/kg, respectively. These doses were associated with peak arterial total plasma concentrations of 11.4, 4.3 and 5 mcg/mL, respectively.

convulsions demonstrated rapid development of hypoxia, hypercarbia and acidosis within a minute of the onset of convulsions. These observations suggest that oxygen consumption and carbon dioxide production are greatly increased during local anesthetic convulsions and emphasize the importance of immediate and effective ventilation with oxygen which may avoid cardiac arrest. If difficulty is encountered in the maintenance of a patent airway

twitching) were noted.

or if prolonged ventilatory support (assisted or controlled) is indicated, endotracheal intubation, employing drugs and techniques familiar to the clinician, may be indicated after initial administration of oxygen by mask.

In human volunteers given intravenous NAROPIN, the mean


(min-max) maximum tolerated total and free arterial plasma concentrations were 4.3 (3.4 to 5.3) and 0.6 (0.3 to 0.9) mcg/mL respectively, at which time moderate CNS symptoms (muscle

Clinical data from patients experiencing local anesthetic induced

The supine position is dangerous in pregnant women at term because of aortocaval compression by the gravid uterus. Therefore, during treatment of systemic toxicity, maternal hypotension or fetal bradycardia following regional block, the parturient should be maintained in the left lateral decubitus position if possible, or manual displacement of the uterus off the great vessels should be accomplished. Resuscitation of obstetrical patients may take longer than resuscitation of non-pregnant patients and closed-chest cardiac compression may be ineffective. Rapid delivery of the fetus may improve the response to resuscitative efforts.

DESCRIPTION

NAROPIN® Injection is a sterile, isotonic solution that contains ropivacaine hydrochloride as the active pharmaceutical ingredient. Ropivacaine hydrochloride is a member of the amino amide class of local anesthetics. NAROPIN® injection is administered parenterally by for infiltration, epidural, and nerve block. Ropivacaine hydrochloride is chemically described as S-(-)-1-propyl-2',6'-pipecoloxylidide hydrochloride monohydrate. The drug substance is a white crystalline powder, with the following structural formula:



At 25 °C ropivacaine hydrochloride has a solubility o 53.8 mg/mL in water, a distribution ratio between *n*-octanol and phosphate buffer at pH 7.4 of 14:1 and a pKa of 8.07 in 0.1 M KCI solution. The pKa of ropivacaine is approximately the same as bupivacaine (8.1) and is similar to that of mepivacaine 7.7). However, ropivacaine has an intermediate degree of lipic solubility compared to bupivacaine and mepivacaine.

NAROPIN (ropivacaine hydrochloride) injection is a clear, color-less, and preservative-free solution. Each mL contains 2.1 mg, 5.3 mg, 7.9 mg or 10.6 mg ropivacaine hydrochloride monohydrate (equivalent to 2.0 mg, 5.0 mg, 7.5 mg or 10 mg of ropivacaine hydrochloride anhydrous), and 8.6 mg, 8.0 mg, 7.5 mg or 7.5 7.1 mg of sodium chloride; respectively, and sodium hydroxide and hydrochloric acid as pH adjusters, in water for injection The pH is adjusted between 4.0 to 6.0. The specific gravity o NAROPIN Injection solutions range from 1.002 to 1.005 at 25°C 12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action Ropivacaine is a member of the amino amide class of local

Ropivacaine is a member of the amino amide class of local anesthetics and is supplied as the pure S-()-enantinmer. Local anesthetics block the generation and the conduction of nerve impulses, presumably by increasing the threshold for electrical excitation in the nerve, by slowing the propagation of the nerve impulse, and by reducing the rate of rise of the action potential. In general, the progression of anesthesia is related to the diameter, myelination and conduction velocity of affected nerve fibers. Clinically, the order of loss of nerve function is as follows: (1) pain. (2) temperature. (3) touch. (4) proprioception, and (1) pain, (2) temperature, (3) touch, (4) proprioception, and (5) skeletal muscle tone. 12.2 Pharmacodynamics
Studies in humans have demonstrated that, unlike most other

local anesthetics, the presence of epinephrine has no major effect on either the time of onset or the duration of action of ropivacaine. Likewise, addition of epinephrine to ropivacaine has no effect on limiting systemic absorption of ropivacaine. Systemic absorption of local anesthetics can produce effects on the central nervous and cardiovascular systems. At blood concentrations achieved with therapeutic doses, changes in cardiac conduction, excitability, refractoriness, contractility, and peripheral vascular resistance have been reported. Toxic blood concentrations depress cardiac conduction and excitability, which may lead to atrioventricular block, ventricular arrhythmas and to cardiac arrest, sometimes resulting in fatallities. In addition, myocardial contractility is depressed and peripheral

vasodilation occurs, leading to decreased cardiac output and arterial blood pressure. Following systemic absorption, local anesthetics can produce central nervous system stimulation, depression or both Apparent central stimulation is usually manifested as restless

respiratory arrest. However, the local anesthetics have a primary depressant effect on the medula and on higher centers. The depressed stage may occur without a prior excited stage. In 2 clinical pharmacology studies (total n=24) ropivacaine and bupivacaine were infused (10 mg/min) in human volunteers until the appearance of CNS symptoms, e.g., visual or hearing disturbances, perioral numbness, tingling and others. Similar symptoms were seen with both drugs. In 1 study, the mean ± SD maximum tolerated intravenous dose of ropivacaine infused (124 ± 38 mg) was significantly higher than that of human significantly (90 + 30 mg) while in the other study the doses bupivacaine (99 ± 30 mg) while in the other study the doses were not different (115 ± 29 mg of ropivacaine and 103 ± 30 mg of bupivacaine). In the latter study, the number of subjects reporting each symptom was similar for both drugs with the exception of muscle twitching which was reported by more subjects with bupivacaine than ropivacaine at comparable intravenous doses. At the end of the infusion, ropivacaine in both studies caused significantly less depression of cardiac

both studies caused significantly less depression of cardiac conductivity (less QRS widening) than bupivacaine. Ropivacaine and bupivacaine caused evidence of depression of cardiac contractility, but there were no changes in cardiac output. Clinical data in one published article indicate that differences in various pharmacodynamic measures were observed with increasing age. In one study, the upper level of analgesia increased with age, the maximum decrease of mean arterial pressure (MAP) declined with age during the first hour after epidural administration, and the intensity of motor blockade increased with age. However, no pharmacokinetic differences were observed between elderly and younger patients.

In non-clinical pharmacology studies comparing ropivacaine and bupivacaine in several animal species, the cardiac toxicity of ropivacaine was less than that of bupivacaine, although both were considerably more toxic than lidocaine animals at significantly higher doses of ropivacaine than bupiva-

12.3 Pharmacokinetics

he systemic concentration of ropivacaine is dependent on the total dose and concentration of drug administered, the route of

biphasic absorption. The half-lives of the 2 phases, (mean \pm SD) are 14 \pm 7 minutes and 4.2 \pm 0.9 h, respectively. The slow absorption is the rate limiting factor in the elimination of ropivacaine that explains why the terminal half-life is longer after epidural than after intravenous administration.

Table 7 Pharmacokinetic (Plasma Concentration-Time) Data From Clinical Trials

	1		IIIIusioii	DIOUK	DIOCK	DIOCK	iiiiusioii
Dose (mg)	1493 ± 10	2075 ± 206	1217 ± 277	150	187.5	300	40
N	12	12	11	8	8	10	12
C _{max} (mg/L)	2.4 ± 1 ¹	2.8 ± 0.5^{1}	2.3 ± 1.1 ¹	1.1 ± 0.2	1.6 ± 0.6	2.3 ± 0.8	1.2 ± 0.2#
T _{max} (min)	n/a*	n/a	n/a	43 ± 14	34 ± 9	54 ± 22	n/a
AUC ₀ - (mg.h/L)	135.5 ± 50	145 ± 34	161 ± 90	7.2 ± 2	11.3 ± 4	13 ± 3.3	1.8 ± 0.6
CL (L/h)	11.03	13.7	n/a	5.5 ± 2	5 ± 2.6	n/a	21.2 ± 7
t _{1/2} (hr)♥	5 ± 2.5	5.7 ± 3	6 ± 3	5.7 ± 2	7.1 ± 3	6.8 ± 3.2	1.9 ± 0.5
* Continuou	s 72 hour e	pidural infus	ion after an	epidural b	lock with 5	or 10 mg	/mL.

n/a=not applicable t/b is the true terminal elimination half-life. On the other hand, t/b follows absorption dependent elimination (flip-flop) after non-intravenous administration.

Committee 2 Problems and Committee (1,75%) for cessarean delivery. Production and Committee (1,75%) for cessarean delivery. Prachial plexus block with 7.5 mg/mL (0,75%) for powacaine. O minute IV influsion to volunteers (40 mg. at 72 hr). Chax measured at the end of influsion (i.e., at 20 minutes).

caine. The incidence of successful resuscitation was not signifi cantly different between the ropivacaine and bupivacaine groups

administration, the patient's hemodynamic/circulatory condition and the vascularity of the administration site. From the epidural space, ropivacaine shows complete and

Ropivacaine shows dose-proportionality up to the highest intravenous dose studied, 80 mg, corresponding to a mean \pm SD peak plasma concentration of 1.9 \pm 0.3 mcg/mL.

Route Epidural Infusion* Epidural Epidural Epidural Plexus IV

In some patients after a 300 mg dose for brachial plexus block. free plasma concentrations of ropivacaine may approach the threshold for CNS toxicity (see Warnings and Precautions (5.7)). At a dose of greater than 300 mg, for local infiltration,

the terminal half-life may be longer (>30 hours).

DistributionAfter intravascular infusion, ropivacaine has a steady-state volume of distribution of 41 \pm 7 liters. Ropivacaine is 94% protein bound, mainly to α 1-acid glycoprotein. An increase in total plasma concentrations during continuous epidural infusion has been observed, related to a postoperative increase of α1-acid glycoprotein. Variations in unbound, i.e., pharmacologically active, concentrations have been less than in total plasma concentration. Ropivacaine readily crosses the placenta and equilibrium in regard to unbound concentration will be rapidly reached [see Warnings and Precautions (5) and Use in Specific Population (8.1)].

Metabolism

Ropivacaine is extensively metabolized in the liver, predominantly by aromatic hydroxylation mediated by cytochrome P4501A to 3-hydroxy ropivacaine. After a single IV dose approximately 37% of the total dose is excreted in the urine as both free and conjugated 3-hydroxy ropivacaine. Low concentrations of 3-hydroxy ropivacaine have been found in the plasma. Urinary S-hydroxy robivacanie have been found in the plasma. Offinally excretion of the 4 hydroxy ropivacaine, and both the 3-hydroxy N de alkylated (3-OH-PPX) and 4-hydroxy N-de-alkylated (4-OH-PPX) metabolites account for less than 3% of the dose. An additional metabolite, 2-hydroxy-methyl-ropivacaine, has been identified but not quantified in the urine. The N-de-alkylated metabolite of ropivacaine (PPX) and 3-OH-ropivacaine are the major metabolite accepted in the urine during poidural in fusion. aior metabolites excreted in the urine during epidural infusion Total PPX concentration in the plasma was about half as that of total ropivacaine; however, mean unbound concentrations of PPX were about 7 to 9 times higher than that of unbound ropivacaine following continuous epidural infusion up to 72 hours. Unbound PPX, 3-hydroxy and 4-hydroxy ropivacaine, have a pharmacological activity in animal models less than that of ropivacaine. There is no evidence of in vivo racemization in urine of ropivacaine.

Elimination
The kidney is the main excretory organ for most local anesthetic metabolites. In total, 86% of the ropivacaine dose is excreted in the urine after intravenous administration of which only 1% relates to unchanged drug. After intravenous administration ropivacaine has a mean ± SD total plasma clearance of 387 ± 107 mL/min, an unbound plasma clearance of 7.2 ± 1.6 L/min, and a renal clearance of 1 mL/min. The mean \pm SD terminal half-life is 1.8 \pm 0.7 h after intravascular administration and 4.2 \pm 1 h after epidural administration.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility Carcinogenesis

Long-term studies in animals to evaluate the carcinogenic potential of ropivacaine have not been conducted. Mutagenesis

Weak mutagenic activity was seen in the mouse lymphoma test. However, ropivacaine was negative in an in vitro Ames assay and an in vivo mouse micronucleus assay.

Impairment of Fertility No adverse effects on fertility or early embryonic development

were reported in a 2-generational reproduction study in which female rats (F₀) were administered subcutaneous doses of 6.3, 12, and 23 mg/kg/day (equivalent to 0.08, 0.15, and 0.29 times the maximum recommended human dose (MRHD) of 770 mg/24 hours for epidural use, respectively, and 0.24, 0.45, and 0.88 times the MRHD of 250 mg for nerve block use, respectivel based on BSA comparisons and a 60 kg human) throughout the nating period and pregnancy, partus, and lactation.

13.2 Animal Toxicology and/or Pharmacology
The mean dosages of ropivacaine producing seizures, after intravenous infusion in dogs, nonpregnant and pregnant sheep were 4.9, 6.1 and 5.9 mg/kg (HED: 5.3, 6.6, and 6.4 mg/kg, based on 75 kg sheep weight and 60 kg human weight), respectively. These doses were associated with peak arterial total plasma concentrations of 11.4, 4.3 and 5 mcg/mL, respectively.

Ropivacaine was studied as a local anesthetic both for surgical anesthesia and for acute pain management [see Dosage and

The onset, depth and duration of sensory block are, in general, similar to bupivacaine. However, the depth and duration of motor block, in general, are less than that with bupivacaine.

14.1 Epidural Administration in Surgery
There were 25 clinical studies performed in 900 patients to evaluate NAROPIN epidural injection for general surgery. NAROPIN was used in doses ranging from 75 to 250 mg. In doses of 100 to 200 mg, the median (1st to 3rd quartile) onset time to achieve a T10 sensory block was 10 (5 to 13) minutes and the median (1st to 3rd quartile) duration at the T10 level was

14.2 Epidural Administration in Cesarean Section

A total of 12 studies were performed with epidural administra-tion of NAROPIN for cesarean section. Eight of these studies involved 218 patients using the concentration of 5 mg/mL (0.5%) in doses up to 150 mg. Median onset measured at T6 ranged from 11 to 26 minutes. Median duration of sensory block at T6 ranged from 1.7 to 3.2 h, and duration of motor block ranged from 1.4 to 2.9 h. NAROPIN provided adequate muscle relaxation for nursery in all cooper.

4 (3 to 5) hours [see Dosage and Administration (2.2)]. Higher oses produced a more profound block with a greater duration

ation for surgery in all cases. In addition, 4 active controlled studies for cesarean section were performed in 264 patients at a concentration of 7.5 mg/mL (0.75%) in doses up to 187.5 mg. Median onset measured at T6 ranged from 4 to 15 minutes. Seventy-seven to 96% of NAROPIN-exposed patients reported no pain at delivery. Some patients received other anesthetic, analgesic, or sedative modalities during the course of the operative procedure.

14.3 Epidural Administration in Labor and Delivery
A total of 9 double-blind clinical studies, involving 240 patients
were performed to evaluate NAROPIN for epidural block for
management of labor pain. When administered in doses up to
278 mg as intermittent injections or as a continuous infusion,
NAROPIN produced adequate pain relief.

A prospective meta-analysis on 6 of these studies provided detailed evaluation of the delivered newborns and showed no difference in clinical outcomes compared to bupivacaine. There were significantly fewer instrumental deliveries in mothers receiving ropivacaine as compared to bupivacaine

Table 8
Labor and Delivery Meta-analysis: Mode of Delivery

Delivery Mode	NAR	OPIN	Bupivacaine n = 188					
Delivery Wode	n =	199						
	n	%	n	%				
Spontaneous Vertex	116	58	92	49				
Vacuum Extractor	26		33					
		}27*		}40				
Forceps	28		42					
Cesarean Section	29	15	21	11				
*p=0.004 versus bupivacaine								

14.4 Epidural Administration in Postoperative Pain Management There were 8 clinical studies performed in 382 patients to evaluate NAROPIN 2 mg/mL (0.2%) for postoperative pain management after upper and lower abdominal surgery and after orthopedic surgery. The studies utilized intravascular morphine via PCA as a rescue medication and quantified as an efficacy

variable.

Epidural anesthesia with NAROPIN 5 mg/mL, (0.5%) was used intraoperatively for each of these procedures prior to initiation of postoperative NAROPIN. The incidence and intensity of the motor block were dependent on the dose rate of NAROPIN and the site of injection. Cumulative doses of up to 770 mg of ropivacaine were administered over 24 hours (intraoperative block plus postoperative continuous infusion). The overall quality of pain relief, as judged by the patients, in the ropivacaine groups was rated as good or excellent (73% to 100%). The frequency of motor block was greatest at 4 hours and decreased during the infusion period in all groups. At least 80% of patients in the upper and lower abdominal studies and 42% in the orthopedic upper and lower abdominal studies and 42% in the orthopedic studies had no motor block at the end of the 21-hour infusion period. Sensory block was also dose rate dependent and a decrease in spread was observed during the infusion period.

A double-blind, randomized, clinical trial compared lumbar epidural infusion of NAROPIN (n=26) and bupivacaine (n=26) at 2 mg/mL (8 mL/h), for 24 hours after knee replacement. In this study, the pain scores were higher in the NAROPIN group but the incidence and the intensity of motor block were lower. Continuous epidural infusion of NAROPIN 2 mg/mL (0.2%) during up to 72 hours for postoperative pain management after major abdominal surgery was studied in 2 multicenter, doubleblind studies. A total of 391 patients received a low thoracic epidural catheter, and NAROPIN 7.5 mg/L (0.75%) was given for surgery, in combination with GA. Postoperatively, NAROPIN 2 mg/mL (0.2%), 4 to 14 mL/h, alone or with fentanyl 1, 2, or 4 mcg/mL was infused through the epidural catheter and

adjusted according to the patient's needs. These studies support the use of NAROPIN 2 mg/mL (0.2%) for epidural infusion at 6 to 14 mL/h (12 to 28 mg) for up to 72 hours and demonstrated adequate analgesia with only slight and nonprogressive motor block in cases of moderate to severe

postoperative pain. Clinical studies with 2 mg/mL (0.2%) NAROPIN have demonstrated that infusion rates of 6 to 14 mL (12 to 28 mg) per hour provide adequate analgesia with nonprogressive motor block in cases of moderate to severe postoperative pain. In these studies, this technique resulted in a significant reduction in patients' morphine rescue dose requirement. Clinical experience supports the use of NAROPIN epidural infusions for up to

14.5 Peripheral Nerve Block NAROPIN, 5 mg/mL (0.5%), was evaluated for its ability to provide anesthesia for surgery using the techniques of Peripheral Nerve Block. There were 13 studies performed including a series of 4 pharmacodynamic and pharmacokinetic studie performed on minor nerve blocks. From these, 235 NAROPIN-treated patients were evaluable for efficacy.

NAROPIN was used in doses up to 275 mg. When used for brachial plexus block, onset depended on technique used. Supraclavicular blocks were consistently more successful than axillary blocks. The median onset of sensory block (anesthesia produced by ropivacaine 0.5% via axillary block ranged from 10 minutes (medial brachial cutaneous nerve) to 45 minutes (musculocutaneous nerve). Median duration ranged from 3.7 hours (medial brachial cutaneous nerve) to 8.7 hours (ulnar nerve). The 5 mg/mL (0.5%) NAROPIN solution gave success rates from 56% to 86% for axillary blocks, compared with 92%

In addition, NAROPIN, 7.5 mg/mL (0.75%), was evaluated in 99 NAROPIN-treated patients, in 2 double-blind studies, performed to provide anesthesia for surgery using the techniques of Brachial Plexus Block. NAROPIN 7.5 mg/mL was compared to bupivacaine 5 mg/mL. In 1 study, patients underwent axillary brachial plexus block using injections of 40 mL (300 mg) of NAROPIN, 7.5 mg/mL (0.75%) or 40 mL injections of bupivacaine, 5 mg/mL (200 mg). In a second study, patients underwent subclavian perivascular brachial plexus block using 30 mL (225 mg) of NAROPIN, 7.5 mg/mL (0.75%) or 30 mL of bupivacaine 5 mg/mL (150 mg). There was no significant difference between the NAROPIN and bupivacaine groups in either study with regard to onset of anesthesia, duration of sensory blockade, or duration of anesthesia.

The median duration of anesthesia varied between 11.4 and 14.4 hours with both techniques. In one study, using the axillary 14.4 nours with both techniques. In one study, using the axiliary technique, the quality of analgesia and muscle relaxation in the NAROPIN group was judged to be significantly superior to bupivacaine by both investigator and surgeon. However, using the subclavian perivascular technique, no statistically significant difference was found in the quality of analgesia and muscle relaxities and was deal with the surgicians are accounted. ation as judged by both the investigator and surgeon. The use of NAROPIN 7.5 mg/mL for block of the brachial plexus via either the subclavian perivascular approach using 30 mL (225 mg) or via the axillary approach using 40 mL (300 mg) both provided effective and reliable anesthesia.

14.6 Local Infiltration

A total of 7 clinical studies were performed to evaluate the local infiltration of NAROPIN to produce anesthesia for surgery and analgesia in postoperative pain management. In these studies 297 patients who received NAROPIN in doses up to 200 mg (concentrations up to 5 mg/mL, 0.5%) were evaluable for efficacy. With infiltration of 100 to 200 mg NAROPIN, the time to first request for analgesic was 2 to 6 hours. When compared to placebo, NAROPIN produced lower pain scores and a reduction of analgesic consumption.

How Supplied/Storage and Handling NAROPIN (ropivacaine hydrochloride) Injection is a clear colorless, and preservative-free solution, available in single-dose containers in 2 mg/mL (0.2%), 5 mg/mL (0.5%), 7.5 mg/mL (0.75%) and 10 mg/mL (1%) concentrations.

Solutions should be stored at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

NAROPIN® Single-Dose Vials

Product Code	Unit of Sale	Strength	Each
278513	NDC 63323-285-13	0.2%, 20 mg per 10 mL	NDC 63323-285-03
	Unit of 25	(2 mg per mL)	10 mL Single-Dose V
278523	NDC 63323-285-23	0.2%, 40 mg per 20 mL	NDC 63323-285-07
	Unit of 25	(2 mg per mL)	20 mL Single-Dose V
278623	NDC 63323-286-23	0.5%,100 mg per 20 mL	NDC 63323-286-05
	Unit of 25	(5 mg per mL)	20 mL Single-Dose V
278630	NDC 63323-286-30	0.5%, 150 mg per 30 mL	NDC 63323-286-30
	Packaged Individually	(5 mg per mL)	30 mL Single-Dose V
278631	NDC 63323-286-31	0.5%, 150 mg per 30 mL	NDC 63323-286-09
	Unit of 5	(5 mg per mL)	30 mL Single-Dose V
278635	NDC 63323-286-35	0.5%, 150 mg per 30 mL	NDC 63323-286-11
	Unit of 25	(5 mg per mL)	30 mL Single-Dose V
278721	NDC 63323-287-21	0.75%, 150 mg per 20 mL	NDC 63323-287-03
	Unit of 25	(7.5 mg per mL)	20 mL Single-Dose V
278811	NDC 63323-288-11	1%, 100 mg per 10 mL	NDC 63323-288-03
	Unit of 25	(10 mg per mL)	10 mL Single-Dose V
278821	NDC 63323-288-21	1%, 200 mg per 20 mL	NDC 63323-288-07
	Unit of 25	(10 mg per mL)	20 mL Single-Dose V

NAROPIN® Single-Dose Infusion Bottles

Product Code	Unit of Sale	Strength	Each
278565	NDC 63323-285-65 Individual bottles sold as a case of 12	0.2%, 200 mg per 100 mL (2 mg per mL)	NDC 63323-285-51 100 mL Single-Dose Infusion Bottle
278564	NDC 63323-285-64 Individual bottles sold as a case of 12	0.2%, 400 mg per 200 mL (2 mg per mL)	NDC 63323-285-57 200 mL Single-Dose Infusion Bottle
278600	NDC 63323-286-00 Unit of 12	0.5%, 500 mg per 100 mL (5 mg per mL)	NDC 63323-286-03 100 mL Single-Dose Infusion Bottle
278663	NDC 63323-286-63 Unit of 12	0.5%, 1,000 mg per 200 mL (5 mg per mL)	NDC 63323-286-33 200 mL Single-Dose Infusion Bottle

For single-dose vials: Discard unused portion.

NAROPIN container closure is not made with natural rubber

17 PATIENT COUNSELING INFORMATION 17.1 Information for Patients and Caregivers

When appropriate, patients should be informed in advance that they may experience temporary loss of sensation and motor activity in the anesthetized part of the body following proper administration of lumbar epidural anesthesia. Also, when appropriate, the physician should discuss other information including adverse reactions in the NAROPIN package insert.

Inform patients that use of local anesthetics may cause methemoglobinemia, a serious condition that must be treated promptly. Advise patients or caregivers to seek immediate medical attention if they or someone in their care experience the following signs or symptoms: pale, gray, or blue colored skin (cyanosis); headache; rapid heart rate; shortness of breath; lightheadedness; or fatigue.

NAROPIN is a trademark of Fresenius Kabi USA, LLC.



451112L