HIGHLIGHTS OF PRESCRIBING INFORMATION These highlights do not include all the information needed to use ONDANSETRON INJECTION safely and effectively. See full prescribing information for ONDANSETRON INJECTION, USP.	 Patients known to h Concomitant use of 		(e.g., anaphylaxis) to this product or any of	f its components. (4)	tramadol, and intrav	enous methylene blue). Some o	f the reported cases we	bitors, mirtazapine, fentanyl, lithium, re fatal. Serotonin syndrome occurring reports of serotonin syndrome related
ONDANSETRON injection for intravenous or intramuscular use			WARNINGS AND PRECAUTIONS		to 5-HT ₃ receptor an	tagonist use occurred in a post-a	anesthesia care unit or a	an infusion center.
Initial U.S. Approval: 1991			vity reactions, including anaphylaxis and broating to other selective 5-HT ₃ receptor antage					combination of signs and symptoms: autonomic instability (e.g., tachycardia,
			<u>s: QT prolongation occurs in a dose-depen</u>		labile blood pressure	, dizziness, diaphoresis, flushing,	, hyperthermia), neurom	uscular symptoms (e.g., tremor, rigidity,
Ondansetron Injection is a 5-HT ₃ receptor antagonist indicated for the prevention of: nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy. (1.1) postoperative nausea and/or vomiting. (1.2) DOSAGE AND ADMINISTRATION	Pointes have been re <u>Serotonin Syndrome</u> concomitant use of <u>Masking of Progress</u>	eported. Avoid Onda <u>::</u> Serotonin syndrom serotonergic drugs. (<u>ive lleus and/or Gas</u> t	nsetron in patients with congenital long QT ne has been reported with 5-HT ₃ receptor a	T syndrome. (5.2) agonists alone but particularly with y or Chemotherapy-Induced Nausea	myoclonus, hyperrefl diarrhea). Patients sl of Ondansetron and and initiate supporti	exia, incoordination), seizures, wi nould be monitored for the emer other serotonergic drugs. If sym re treatment. Patients should be i	ith or without gastrointes rgence of serotonin synd ptoms of serotonin synd informed of the increase	stinal symptoms (e.g., nausea, vomiting, lrome, especially with concomitant use drome occur, discontinue Ondansetron d risk of serotonin syndrome, especially <i>ig Interactions (7.5), Overdosage (10)</i>].
 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Cancer Chemotherapy (2.1): Dilution of Ondansetron Injection in 50 mL of 5% Dextrose Injection or 0.9% Sodium Chloride Injection is required before 	obstruction. (5.4)			-	5.4 Masking of Progre	ssive lleus and Gastric Dister	nsion	
administration to adult and pediatric patients. • Adults and pediatric patients 6 months of age and older: The recommended dosage is 0.15 mg/kg per dose for 3 doses	<u>Chemotherapy-Induced</u>	d Nausea and Vomiti	ADVERSE REACTIONS		and vomiting may m	ron in patients following abdom ask a progressive ileus and gastr factors for gastrointestinal obstr	ric distention. Monitor fo	its with chemotherapy-induced nausea or decreased bowel activity, particularly
(maximum of 16 mg per dose), infused intravenously over 15 minutes.Administer the first dose 30 minutes before the start of chemotherapy and subsequent doses 4 and 8 hours after the	The most common a	dverse reactions (\geq	7%) in adults are diarrhea, headache, and f	fever. (6.1)	Ondansetron Injection	•		stalsis. It should not be used instead of
first dose.	 Postoperative Nausea a The most common a 	and vomiting: adverse reaction (≥`	10%) which occurs at a higher frequency c	compared with placebo in adults is	nasogastric suction.			
 Prevention of Postoperative Nausea and/or Vomiting (2.2): Do not administer a full prefilled syringe (4 mg dose) to pediatric patients less than 40 kg as this exceeds the recommended dose. 	headache. (6.1)	adverse reaction (\geq	2%) which occurs at a higher frequency co		 Hypersensitivity 	Ily significant adverse reactions reactions [see Warnings and Pre-	cautions (5.1)]	e in the labeling:
 Dilution of Ondansetron Injection is not required before administration to adult and pediatric patients. See full prescribing information for the recommended dosage and administration instructions for adult and pediatric patients 1 month of age and older. 	1-800-FDA-1088 or v	vww.fda.gov/medv		C at 1-800-551-7176 or FDA at	 Serotonin syndro 	[see Warnings and Precautions (ome [see Warnings and Precaution ressive ileus and gastric distention	ons (5.3)]	recautions (5.4)]
Patients With Severe Hepatic Impairment (2.3): Do not exceed a total daily dose of 8 mg. 	See 17 for PATIENT C	OUNSELING INFO	RMATION.		6.1 Clinical Trials Expe Because clinical trial trials of a drug cann	s are conducted under widely va	rying conditions, adverse	e reaction rates observed in the clinical f another drug and may not reflect the
DOSAGE FORMS AND STRENGTHS					rates observed in cli			another drug and may not reliect the
Ondansetron Injection, USP (2 mg/mL): 2 mL Prefilled disposable single-use syringe (3)				Revised: 10/2021	the active ingredien	of intravenous Ondansetron ac		ult patients treated with ondansetron, 5. A causal relationship to therapy with
FULL PRESCRIBING INFORMATION: CONTENTS*	8 USE IN SPECIFIC	POPULATIONS			Ondansetron was ur	eed Nausea and Vomiting		
 INDICATIONS AND USAGE Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Cancer Chemotherapy 	8.1 Pregnancy 8.2 Lactation 8.4 Pediatric Use	2				e Reactions Reported in > 5%		
						at a Dosage of Thr	ree 0.15-mg/kg Doses	
1.2 Prevention of Postoperative Nausea and/or Vomiting	8.5 Geriatric Use 8.6 Hepatic Impa	2					ree 0.15-mg/kg Doses er of Adult Patients W	
1.2 Prevention of Postoperative Nausea and/or Vomiting DOSAGE AND ADMINISTRATION	8.5 Geriatric Use	e airment			Adverse Reaction		er of Adult Patients W	ith Reaction
1.2 Prevention of Postoperative Nausea and/or Vomiting	 8.5 Geriatric Use 8.6 Hepatic Impa 8.7 Renal Impair 9 DRUG ABUSE AN 	e airment ment			Adverse Reaction	Numbe		ith Reaction
 Prevention of Postoperative Nausea and/or Vomiting DOSAGE AND ADMINISTRATION Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy Prevention of Postoperative Nausea and Vomiting Dosage Adjustment for Patients With Hepatic Impairment DOSAGE FORMS AND STRENGTHS 	 8.5 Geriatric Use 8.6 Hepatic Impa 8.7 Renal Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 	e airment ment			Adverse Reaction	Numbe Ondansetron Injection 0.15 mg/kg x 3	er of Adult Patients W Metoclopramide	ith Reaction e Placebo
 Prevention of Postoperative Nausea and/or Vomiting DOSAGE AND ADMINISTRATION Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy Prevention of Postoperative Nausea and Vomiting Dosage Adjustment for Patients With Hepatic Impairment DOSAGE FORMS AND STRENGTHS CONTRAINDICATIONS 	 8.5 Geriatric Use 8.6 Hepatic Impair 8.7 Renal Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 	e airment ment D DEPENDENCE				Numbe Ondansetron Injection 0.15 mg/kg x 3 (n = 419)	er of Adult Patients W Metoclopramido (n = 156)	e Placebo (n = 34)
 Prevention of Postoperative Nausea and/or Vomiting DOSAGE AND ADMINISTRATION Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy Prevention of Postoperative Nausea and Vomiting Dosage Adjustment for Patients With Hepatic Impairment DOSAGE FORMS AND STRENGTHS CONTRAINDICATIONS WARNINGS AND PRECAUTIONS Hypersensitivity Reactions 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism 	e airment D DEPENDENCE MACOLOGY of Action			Diarrhea	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16%	er of Adult Patients W Metoclopramide (n = 156) 44%	ith Reaction e Placebo (n = 34) 18%
 Prevention of Postoperative Nausea and/or Vomiting DOSAGE AND ADMINISTRATION Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy Prevention of Postoperative Nausea and Vomiting Dosage Adjustment for Patients With Hepatic Impairment DOSAGE FORMS AND STRENGTHS CONTRAINDICATIONS WARNINGS AND PRECAUTIONS Nupersensitivity Reactions Q T Prolongation Serotonin Syndrome 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism 12.2 Pharmacody 12.3 Pharmacody 	e airment D DEPENDENCE MACOLOGY of Action namics netics			Diarrhea Headache Fever	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8%	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5%	Placebo (n = 34) 18% 15%
 Prevention of Postoperative Nausea and/or Vomiting DOSAGE AND ADMINISTRATION Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy Prevention of Postoperative Nausea and Vomiting Dosage Adjustment for Patients With Hepatic Impairment DOSAGE FORMS AND STRENGTHS CONTRAINDICATIONS WARNINGS AND PRECAUTIONS Hypersensitivity Reactions QT Prolongation 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism 12.2 Pharmacody 12.3 Pharmacokir 13 NONCLINICAL TC 	airment ment D DEPENDENCE MACOLOGY of Action namics netics DXICOLOGY	pairment of Fertility		Diarrhea Headache Fever <i>Cardiovascular:</i> Rare have been reported.	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% electrocardiographic alte	Placebo e Placebo (n = 34) 18% 15% 3%
 Prevention of Postoperative Nausea and/or Vomiting DOSAGE AND ADMINISTRATION Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy Prevention of Postoperative Nausea and Vomiting Dosage Adjustment for Patients With Hepatic Impairment DOSAGE FORMS AND STRENGTHS CONTRAINDICATIONS WARNINGS AND PRECAUTIONS I Hypersensitivity Reactions Q T Prolongation Serotonin Syndrome Masking of Progressive Ileus and Gastric Distension ADVERSE REACTIONS Clinical Trials Experience 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism 12.2 Pharmacody 12.3 Pharmacokir 13 NONCLINICAL TO 13.1 Carcinogene 14 CLINICAL STUDIE 	e airment ment D DEPENDENCE MACOLOGY of Action namics netics DXICOLOGY sis, Mutagenesis, Im ES			Diarrhea Headache Fever <i>Cardiovascular:</i> Rare have been reported. <i>Gastrointestinal:</i> Cor <i>Hepatic:</i> In compara	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e stipation has been reported in 1 tive trials in cisplatin chemothe	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% electrocardiographic alter 1% of chemotherapy pa rapy patients with norm	ith Reaction e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia attents receiving multiday ondansetron. nal baseline values of aspartate trans-
 Prevention of Postoperative Nausea and/or Vomiting DOSAGE AND ADMINISTRATION Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy Prevention of Postoperative Nausea and Vomiting Dosage Adjustment for Patients With Hepatic Impairment DOSAGE FORMS AND STRENGTHS CONTRAINDICATIONS WARNINGS AND PRECAUTIONS Itypersensitivity Reactions Q T Prolongation Serotonin Syndrome Masking of Progressive Ileus and Gastric Distension ADVERSE REACTIONS Clinical Trials Experience Postmarketing Experience DRUG INTERACTIONS 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARN 12.1 Mechanism 12.2 Pharmacody 12.3 Pharmacokir 13 NONCLINICAL TO 13.1 Carcinogene 14.1 Chemothera 14.2 Prevention of 	e airment ment D DEPENDENCE MACOLOGY of Action namics netics DXICOLOGY sis, Mutagenesis, Im ES py-induced Nausea <i>i</i> f Postoperative Nau	and Vomiting sea and/or Vomiting		Diarrhea Headache Fever <i>Cardiovascular:</i> Rare have been reported. <i>Gastrointestinal:</i> Cor <i>Hepatic:</i> In compara aminase (AST) and a of normal in approx	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e istipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The increase	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% electrocardiographic alte 1% of chemotherapy pa rrapy patients with norm e enzymes have been re eases were transient an	e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia attents receiving multiday ondansetron. nal baseline values of aspartate transported to exceed twice the upper limit d id not appear to be related to dose
 1.2 Prevention of Postoperative Nausea and/or Vomiting 2 DOSAGE AND ADMINISTRATION 2.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy 2.2 Prevention of Postoperative Nausea and Vomiting 2.3 Dosage Adjustment for Patients With Hepatic Impairment 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS 5 WARNINGS AND PRECAUTIONS 5.1 Hypersensitivity Reactions 5.2 QT Prolongation 5.3 Serotonin Syndrome 5.4 Masking of Progressive Ileus and Gastric Distension 6 ADVERSE REACTIONS 6.1 Clinical Trials Experience 6.2 Postmarketing Experience 7 DRUG INTERACTIONS 7.1 Drugs Affecting Cytochrome P-450 Enzymes 7.2 Apomorphine 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism 12.2 Pharmacody 13.1 Carcinogene 14 CLINICAL STUDIE 14.1 Chemothera 14.2 Prevention o 14.3 Prevention o 	e airment ment D DEPENDENCE MACOLOGY of Action namics hetics DXICOLOGY sis, Mutagenesis, Im S py-induced Nausea a f Postoperative Nau f Further Postoperat	and Vomiting sea and/or Vomiting ive Nausea and Vomiting		Diarrhea Headache Fever <i>Cardiovascular:</i> Rare have been reported. <i>Gastrointestinal:</i> Cor <i>Hepatic:</i> In compara aminase (AST) and a of normal in approx or duration of thera	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e istipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The increase	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% electrocardiographic alte 1% of chemotherapy patients with norm e enzymes have been re eases were transient an transient elevations in f	ith Reaction e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia attents receiving multiday ondansetron. nal baseline values of aspartate trans- ported to exceed twice the upper limit
 1.2 Prevention of Postoperative Nausea and/or Vomiting 2 DOSAGE AND ADMINISTRATION 2.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy 2.2 Prevention of Postoperative Nausea and Vomiting 2.3 Dosage Adjustment for Patients With Hepatic Impairment 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS 5 WARNINGS AND PRECAUTIONS 5.1 Hypersensitivity Reactions 5.2 QT Prolongation 5.3 Serotonin Syndrome 5.4 Masking of Progressive Ileus and Gastric Distension 6 ADVERSE REACTIONS 6.1 Clinical Trials Experience 6.2 Postmarketing Experience 7 DRUG INTERACTIONS 7.1 Drugs Affecting Cytochrome P-450 Enzymes 7.2 Apomorphine 7.3 Phenytoin, Carbamazepine, and Rifampin 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism (12.2 Pharmacody) 12.3 Pharmacody 13.1 Carcinogene 14 CLINICAL STUDII 14.1 Chemothera 14.2 Prevention on 14.3 Prevention on 14.1 Chemothera 14.2 Prevention on 14.3 Prevention on 14.1 Chemothera 14.2 Prevention on 14.3 Prevention on 	airment ment D DEPENDENCE MACOLOGY of Action namics netics DXICOLOGY sis, Mutagenesis, Im ES py-induced Nausea a f Postoperative Nau f Further Postoperat STORAGE AND HAN	and Vomiting sea and/or Vomiting ive Nausea and Vomiting IDLING		Diarrhea Headache Fever Cardiovascular: Rare have been reported. Gastrointestinal: Cor Hepatic: In compara aminase (AST) and a of normal in approx or duration of thera courses, but symptor	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e stipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The incre py. On repeat exposure, similar	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% electrocardiographic alter 1% of chemotherapy patients with norm e enzymes have been re eases were transient and transient elevations in to cur.	e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia stients receiving multiday ondansetron. nal baseline values of aspartate transported to exceed twice the upper limit d id not appear to be related to dose transaminase values occurred in some
 1.2 Prevention of Postoperative Nausea and/or Vomiting 2 DOSAGE AND ADMINISTRATION 2.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy 2.2 Prevention of Postoperative Nausea and Vomiting 2.3 Dosage Adjustment for Patients With Hepatic Impairment 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS 5 WARNINGS AND PRECAUTIONS 5.1 Hypersensitivity Reactions 5.2 QT Prolongation 5.3 Serotonin Syndrome 5.4 Masking of Progressive Ileus and Gastric Distension 6 ADVERSE REACTIONS 6.1 Clinical Trials Experience 6.2 Postmarketing Experience 6.2 Postmarketing Experience 7 DRUG INTERACTIONS 7.1 Drugs Affecting Cytochrome P-450 Enzymes 7.2 Apomorphine 7.3 Phenytoin, Carbamazepine, and Rifampin 7.4 Tramadol 7.5 Serotonergic Drugs 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism 12.2 Pharmacody 13.1 Carcinogene 14 CLINICAL STUDIE 14.1 Chemothera 14.2 Prevention o 14.3 Prevention o 	airment ment D DEPENDENCE MACOLOGY of Action namics netics DXICOLOGY sis, Mutagenesis, Im ES py-induced Nausea a f Postoperative Nau f Further Postoperat STORAGE AND HAN	and Vomiting sea and/or Vomiting ive Nausea and Vomiting IDLING		Diarrhea Headache Fever Cardiovascular: Rare have been reported. Gastrointestinal: Cor Hepatic: In compara aminase (AST) and a of normal in approx or duration of thera courses, but sympto Integumentary: Rash Neurological: There H	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e stipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The increpy. On repeat exposure, similar natic hepatic disease did not oco has occurred in approximately	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% electrocardiographic alte 1% of chemotherapy pa arapy patients with norm e enzymes have been re eases were transient and transient elevations in ta cur. 1% of patients receiving with, but not diagnostic	e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia stients receiving multiday ondansetron. nal baseline values of aspartate transported to exceed twice the upper limit d id not appear to be related to dose transaminase values occurred in some
 1.2 Prevention of Postoperative Nausea and/or Vomiting 2 DOSAGE AND ADMINISTRATION 2.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy 2.2 Prevention of Postoperative Nausea and Vomiting 2.3 Dosage Adjustment for Patients With Hepatic Impairment 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS 5 WARNINGS AND PRECAUTIONS 5.1 Hypersensitivity Reactions 5.2 QT Prolongation 5.3 Serotonin Syndrome 5.4 Masking of Progressive Ileus and Gastric Distension 6 ADVERSE REACTIONS 6.1 Clinical Trials Experience 6.2 Postmarketing Experience 6.2 Postmarketing Experience 7 DRUG INTERACTIONS 7.1 Drugs Affecting Cytochrome P-450 Enzymes 7.2 Apomorphine 7.3 Phenytoin, Carbamazepine, and Rifampin 7.4 Tramadol 7.5 Serotomergic Drugs 7.6 Chemotherapy 7.7 Temazepam 	 8.5 Geriatric Use 8.6 Hepatic Imp. 8.7 Renal Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism (12.2 Pharmacody) 12.3 Pharmacody 12.3 Pharmacokin 13 NONCLINICAL TC 13.1 Carcinogene 14 CLINICAL STUDII 14.1 Chemothera 14.2 Prevention on 14.3 Prevention on 14.1 Chemothera 14.2 Prevention on 14.3 Prevention on 14.1 Chemothera 14.2 Prevention on 14.3 Prevention on 14.1 PATIENT COUNSIDE 	airment ment D DEPENDENCE MACOLOGY of Action namics netics DXICOLOGY sis, Mutagenesis, Im ES py-induced Nausea f Postoperative Nau f Forther Postoperat STORAGE AND HAN ELING INFORMATIO	and Vomiting sea and/or Vomiting ive Nausea and Vomiting NDLING DN		Diarrhea Headache Fever Cardiovascular: Rare have been reported. Gastrointestinal: Cor Hepatic: In compara aminase (AST) and a of normal in approx or duration of thera courses, but symptor Integumentary: Rash Neurological: There h receiving Ondansetr	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e stipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The increported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The increported in 1 natic hepatic disease did not occup thas occurred in approximately ave been rare reports consistent	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% dectrocardiographic alte 1% of chemotherapy pa rapy patients with norm e enzymes have been re eases were transient and transient elevations in for cur. 1% of patients receiving with, but not diagnostic rand mal seizure.	e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia attents receiving multiday ondansetron. nal baseline values of aspartate transported to exceed twice the upper limit d id not appear to be related to dose transaminase values occurred in some g ondansetron.
 Prevention of Postoperative Nausea and/or Vomiting DOSAGE AND ADMINISTRATION Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy Prevention of Postoperative Nausea and Vomiting Dosage Adjustment for Patients With Hepatic Impairment DOSAGE FORMS AND STRENGTHS CONTRAINDICATIONS WARNINGS AND PRECAUTIONS Hypersensitivity Reactions Q T Prolongation Serotonin Syndrome Masking of Progressive Ileus and Gastric Distension ADVERSE REACTIONS Clinical Trials Experience Postmarketing Experience Postmarketing Experience Dordy Affecting Cytochrome P-450 Enzymes Apomorphine Serotonergic Drugs Chemotherapy Serotonergic Drugs Chemotherapy Serotonergic Drugs Alfentanil and Atracurium 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARN 12.1 Mechanism (12.2 Pharmacody) 12.3 Pharmacody 13 NONCLINICAL TO 13.1 Carcinogene 14 CLINICAL STUDIE 14.1 Chemothera 14.2 Prevention on 14.3 Prevention on 14.3 Prevention on 15 PATIENT COUNSI *Sections or subsection 	airment ment D DEPENDENCE MACOLOGY of Action namics netics DXICOLOGY sis, Mutagenesis, Im S py-induced Nausea f Postoperative Nau f Further Postoperat STORAGE AND HAN ELING INFORMATION as omitted from the f	and Vomiting sea and/or Vomiting ive Nausea and Vomiting IDLING		Diarrhea Headache Fever Cardiovascular: Rare have been reported. Gastrointestinal: Cor Hepatic: In compara aminase (AST) and a of normal in approx or duration of thera courses, but symptor Integumentary: Rasl Neurological: There h receiving Ondansetr Other: Rare cases of Postoperative Nause The adverse reactior	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e stipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these py. On repeat exposure, similar natic hepatic disease did not oc has occurred in approximately ave been rare reports consistent on Injection, and rare cases of gr hypokalemia have been reported and Vomiting s in Table 3 have been reported	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% electrocardiographic alter 1% of chemotherapy patients with norm e enzymes have been re eases were transient and transient elevations in for cur. 1% of patients receiving with, but not diagnostic rand mal seizure. ed.	e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia attents receiving multiday ondansetron. nal baseline values of aspartate transported to exceed twice the upper limit d id not appear to be related to dose transaminase values occurred in some g ondansetron.
 1.2 Prevention of Postoperative Nausea and/or Vomiting 2 DOSAGE AND ADMINISTRATION 2.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy 2.2 Prevention of Postoperative Nausea and Vomiting 2.3 Dosage Adjustment for Patients With Hepatic Impairment 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS 5 WARNINGS AND PRECAUTIONS 5.1 Hypersensitivity Reactions 5.2 QT Prolongation 5.3 Serotonin Syndrome 5.4 Masking of Progressive Ileus and Gastric Distension 6 ADVERSE REACTIONS 6.1 Clinical Trials Experience 6.2 Postmarketing Experience 6.2 Postmarketing Experience 7 DRUG INTERACTIONS 7.1 Drugs Affecting Cytochrome P-450 Enzymes 7.2 Apomorphine 7.3 Phenytoin, Carbamazepine, and Rifampin 7.4 Tramadol 7.5 Serotomergic Drugs 7.6 Chemotherapy 7.7 Temazepam 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12. Pharmacody 12. Pharmacody 13 NONCLINICAL TO 13.1 Carcinogene 14 CLINICAL STUDIE 14.1 Chemothera 14.2 Prevention o 14.3 Prevention o 14.3 Prevention o 15 HOW SUPPLIED/S 17 PATIENT COUNSI *Sections or subsection 	A airment ment D DEPENDENCE MACOLOGY of Action namics hetics DXICOLOGY sis, Mutagenesis, Im S py-induced Nausea f Postoperative Nau f Further Postoperat STORAGE AND HAN ELING INFORMATION as omitted from the f	and Vomiting sea and/or Vomiting ive Nausea and Vomiting NDLING DN full prescribing information are not listed.	natients 1 month of are and older	Diarrhea Headache Fever Cardiovascular: Rare have been reported. Gastrointestinal: Cor Hepatic: In compara aminase (AST) and a of normal in approx or duration of thera courses, but symptor Integumentary: Rash Neurological: There H receiving Ondansetr Other: Rare cases of Postoperative Nause The adverse reactior intravenous over 2 t	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e stipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The increpy. On repeat exposure, similar natic hepatic disease did not occl has occurred in approximately ave been rare reports consistent on Injection, and rare cases of gr hypokalemia have been reported and Vomiting is in Table 3 have been reported o 5 minutes in clinical trials.	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% dectrocardiographic alter 1% of chemotherapy patients with norm e enzymes have been re eases were transient and transient elevations in for cur. 1% of patients receiving with, but not diagnostic rand mal seizure. ed. in ≥ 2% of adults receiving	ith Reaction e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia attents receiving multiday ondansetron. nal baseline values of aspartate transported to exceed twice the upper limit d did not appear to be related to dose transaminase values occurred in some g ondansetron. of, extrapyramidal reactions in patients iving ondansetron at a dosage of 4 mg
 1.2 Prevention of Postoperative Nausea and/or Vomiting 2 DOSAGE AND ADMINISTRATION 2.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy 2.2 Prevention of Postoperative Nausea and Vomiting 2.3 Dosage Adjustment for Patients With Hepatic Impairment 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS 5.1 Hypersensitivity Reactions 5.2 QT Prolongation 5.3 Sectonin Syndrome 5.4 Masking of Progressive Ileus and Gastric Distension 6 ADVERSE REACTIONS 6.1 Clinical Trials Experience 6.2 Postmarketing Experience 7 DRUG INTERACTIONS 7.1 Drugs Affecting Cytochrome P-450 Enzymes 7.2 Apomorphine 7.3 Phenytoin, Carbamazepine, and Rifampin 7.4 Tramadol 7.5 Serotonergic Drugs 7.6 Chemotherapy 7.7 Temazepam 7.8 Alfentanil and Atracurium FULL PRESCRIBING INFORMATION 1 INDICATIONS AND USAGE 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism (12.2 Pharmacody) 12.3 Pharmacody 13 NONCLINICAL TO 13.1 Carcinogene 14 CLINICAL STUDIE 14.1 Chemothera 14.2 Prevention o 14.3 Prevention o 14.3 Prevention o 15 PATIENT COUNSI *Sections or subsection 	A airment ment D DEPENDENCE MACOLOGY of Action namics tetics DXICOLOGY sis, Mutagenesis, Im S py-induced Nausea f Postoperative Nau f Further Postoperat STORAGE AND HAN ELING INFORMATION as omitted from the f <u>ninistration</u> ed dose and administ f postoperative naus	and Vomiting sea and/or Vomiting ive Nausea and Vomiting NDLING DN full prescribing information are not listed.		Diarrhea Headache Fever Cardiovascular: Rare have been reported. Gastrointestinal: Cor Hepatic: In compara aminase (AST) and a of normal in approx or duration of thera courses, but symptor Integumentary: Rash Neurological: There H receiving Ondansetr Other: Rare cases of <u>Postoperative Nause</u> The adverse reactior intravenous over 2 tt Table 3. Adverse Rea	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e stipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The increpy. On repeat exposure, similar natic hepatic disease did not occl has occurred in approximately ave been rare reports consistent on Injection, and rare cases of gr hypokalemia have been reported and Vomiting is in Table 3 have been reported o 5 minutes in clinical trials.	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% dectrocardiographic alte 1% of chemotherapy pa rapy patients with norm e enzymes have been re eases were transient an transient elevations in t cur. 1% of patients receiving with, but not diagnostic rand mal seizure. ed. in ≥ 2% of adults recei d with greater freque	ith Reaction e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia attents receiving multiday ondansetron. nal baseline values of aspartate transported to exceed twice the upper limit d did not appear to be related to dose transaminase values occurred in some g ondansetron. of, extrapyramidal reactions in patients iving ondansetron at a dosage of 4 mg ency than the placebo group) of
 1.2 Prevention of Postoperative Nausea and/or Vomiting 2 DOSAGE AND ADMINISTRATION 2.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy 2.2 Prevention of Postoperative Nausea and Vomiting 2.3 Dosage Adjustment for Patients With Hepatic Impairment 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS 5 WARNINGS AND PRECAUTIONS 5.1 Hypersensitivity Reactions 5.2 QT Prolongation 5.3 Serotonin Syndrome 5.4 Masking of Progressive Ileus and Gastric Distension 6 ADVERSE REACTIONS 6.1 Clinical Trials Experience 6.2 Postmarketing Experience 7 DRUG INTERACTIONS 7.1 Drugs Affecting Cytochrome P-450 Enzymes 7.2 Apomorphine 7.3 Phenytoin, Carbamazepine, and Rifampin 7.4 Tramadol 7.5 Serotonergic Drugs 7.6 Chemotherapy 7.7 Ternazepam 7.8 Alfentanil and Atracurium FULL PRESCRIBING INFORMATION 1 INDICATIONS AND USAGE 1.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Cancer Chemotherapy	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism (12.2 Pharmacody) 12.3 Pharmacody 13 NONCLINICAL TO 13.1 Carcinogene 14 CLINICAL STUDIE 14.1 Chemothera 14.2 Prevention o 14.3 Prevention o 14.3 Prevention o 15 PATIENT COUNSI *Sections or subsection 	Airment ment D DEPENDENCE MACOLOGY of Action namics netics DXICOLOGY sis, Mutagenesis, Im S py-induced Nausea f Postoperative Nau f Further Postoperat STORAGE AND HAN ELING INFORMATION as omitted from the f <u>ninistration</u> ed dose and administ f postoperative naus a 1. Recommended Prevention	and Vomiting sea and/or Vomiting ive Nausea and Vomiting NDLING DN full prescribing information are not listed.	ron Injection for	Diarrhea Headache Fever Cardiovascular: Rare have been reported. Gastrointestinal: Cor Hepatic: In compara aminase (AST) and a of normal in approx or duration of thera courses, but symptor Integumentary: Rash Neurological: There H receiving Ondansetr Other: Rare cases of <u>Postoperative Nause</u> The adverse reactior intravenous over 2 tt Table 3. Adverse Rea	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e stipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The increpy. On repeat exposure, similar natic hepatic disease did not occl has occurred in approximately ave been rare reports consistent on Injection, and rare cases of gr hypokalemia have been reported a and Vomiting is in Table 3 have been reported o 5 minutes in clinical trials. ctions Reported in ≥ 2% (an Receiving Ondansetron at a D	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% electrocardiographic alte 1% of chemotherapy patients with norm e enzymes have been re eases were transient and transient elevations in the cur. 1% of patients receiving with, but not diagnostic rand mal seizure. ed. in ≥ 2% of adults receiving d with greater freque bosage of 4 mg Intravenous	ith Reaction e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia attents receiving multiday ondansetron. nal baseline values of aspartate transported to exceed twice the upper limit d did not appear to be related to dose transaminase values occurred in some g ondansetron. of, extrapyramidal reactions in patients iving ondansetron at a dosage of 4 mg ency than the placebo group) of
 1.2 Prevention of Postoperative Nausea and/or Vomiting 2 DOSAGE AND ADMINISTRATION 2.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy 2.2 Prevention of Postoperative Nausea and Vomiting 2.3 Dosage Adjustment for Patients With Hepatic Impairment 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS 5.1 Hypersensitivity Reactions 5.2 QT Prolongation 5.3 Serotonin Syndrome 5.4 Masking of Progressive Ileus and Gastric Distension 6 ADVERSE REACTIONS 6.1 Clinical Trials Experience 6.2 Postmarketing Experience 7 DRUG INTERACTIONS 7.1 Drugs Affecting Cytochrome P-450 Enzymes 7.2 Apomorphine 7.3 Phenytoin, Carbamazepine, and Rifampin 7.4 Tramadol 7.5 Serotonergic Drugs 7.6 Chemotherapy 7.7 Temazepam 7.8 Alfentanil and Atracurium FULL PRESCRIBING INFORMATION 1 INDICATIONS AND USAGE 1.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Cancer 	 8.5 Geriatric Use 8.6 Hepatic Impair 9 DRUG ABUSE AN 10 OVERDOSAGE 11 DESCRIPTION 12 CLINICAL PHARM 12.1 Mechanism (12.2 Pharmacody) 12.3 Pharmacody 13 NONCLINICAL TO 13.1 Carcinogene 14 CLINICAL STUDIE 14.1 Chemothera 14.2 Prevention o 14.3 Prevention o 14.3 Prevention o 15 PATIENT COUNSI *Sections or subsection 	A airment ment D DEPENDENCE MACOLOGY of Action namics hetics DXICOLOGY sis, Mutagenesis, Im S py-induced Nausea f Postoperative Nau f Further Postoperat STORAGE AND HAN ELING INFORMATION as omitted from the f <u>hinistration</u> ed dose and administ f postoperative nause a 1. Recommended	and Vomiting sea and/or Vomiting ive Nausea and Vomiting NDLING DN full prescribing information are not listed. stration instructions for adult and pediatric ea and vomiting are shown in Table 1.	ron Injection for	Diarrhea Headache Fever Cardiovascular: Rare have been reported. Gastrointestinal: Cor Hepatic: In compara aminase (AST) and a of normal in approx or duration of thera courses, but sympto Integumentary: Rash Neurological: There h receiving Ondansetr Other: Rare cases of <u>Postoperative Nause</u> The adverse reactior intravenous over 2 t Table 3. Adverse Rea Adult Patients I	Number Ondansetron Injection 0.15 mg/kg x 3 (n = 419) 16% 17% 8% cases of angina (chest pain), e stipation has been reported in 1 tive trials in cisplatin chemothe lanine transaminase (ALT), these mately 5% of patients. The incomposition of the patic disease did not occur has occurred in approximately ave been rare reports consistent on Injection, and rare cases of gin hypokalemia have been reported a and Vomiting s in Table 3 have been reported in 2 2% (an Receiving Ondansetron at a D	er of Adult Patients W Metoclopramide (n = 156) 44% 7% 5% dectrocardiographic alte 1% of chemotherapy pa rapy patients with norm e enzymes have been re eases were transient an transient elevations in 1 cur. 1% of patients receiving with, but not diagnostic rand mal seizure. ed. in ≥ 2% of adults received d with greater freque bosage of 4 mg Intravenous 547)	ith Reaction e Placebo (n = 34) 18% 15% 3% erations, hypotension, and tachycardia attents receiving multiday ondansetron. nal baseline values of aspartate transported to exceed twice the upper limit d did not appear to be related to dose transaminase values occurred in some g ondansetron. of, extrapyramidal reactions in patients iving ondansetron at a dosage of 4 mg incy than the placebo group) of enous Over 2 to 5 Minutes Placebo

1.2 Prevention of Postoperative Nausea and/or Vomiting

Ondansetron Injection is indicated for the prevention of postoperative nausea and/or vomiting. As with other antiemetics, routine prophylaxis is not recommended for patients in whom there is little expectation that nausea and/or vomiting will occur postoperatively. In patients in whom nausea and/or vomiting must be avoided postoper atively. Ondansetron Injection is recommended even when the incidence of postoperative nausea and/or vomiting is low. For patients who do not receive prophylactic Ondansetron Injection and experience nausea and/or vomiting ostoperatively, Ondansetron Injection may be given to prevent further episodes.

Ondansetron is approved for patients aged 1 month and older.

2 DOSAGE AND ADMINISTRATION

2.1 Prevention of Nausea and Vomiting Associated With Initial and Repeat Courses of Emetogenic Chemotherapy

Important Preparation Instructions

Dilution of Ondansetron Injection in 50 mL of 5% Dextrose Injection or 0.9% Sodium Chloride Injection is required before administration to adult and pediatric patients for the prevention of nausea and vomiting associated with emetogenic chemotherapy

For pediatric patients between 6 months and 1 year of age and/or 10 kg or less: Depending on the fluid needs of the patient, Ondansetron Injection may be diluted in 10 to 50 mL of 5% Dextrose Injection or 0.9% Sodium Chloride 2.3 Dosage Adjustment for Patients With Hepatic Impairment

- Do not mix Ondansetron Injection with solutions for which physical and chemical compatibility has not been established. In particular, this applies to alkaline solutions as a precipitate may form.
- Inspect the diluted Ondansetron Injection solution for particulate matter and discoloration before administration; discard if present. • Storage: After dilution, do not use beyond 24 hours. Although Ondansetron Injection is chemically and physically
- stable when diluted as recommended, sterile precautions should be observed because diluents generally do not contain preservative.
- Compatibility: Ondansetron Injection is compatible and stable at room temperature under normal lighting conditions for 48 hours after dilution with the following intravenous fluids: 0.9% Sodium Chloride Injection, 5% Dextrose Injection, 5% Dextrose and 0.9% Sodium Chloride Injection, 5% Dextrose and 0.45% Sodium Chloride Injection, and 3% Sodium Chloride Injection.

Dosage and Administration

The recommended dosage for adult and pediatric patients 6 months of age and older for prevention of nausea and vomiting associated with emetogenic chemotherapy is 0.15-mg/kg per dose for 3 doses (maximum of 16 mg per Caution: Dilution of Ondansetron Injection is required in adult and pediatric patients prior to administration

Infuse intravenously over 15 minutes beginning 30 minutes before the start of emetogenic chemotherapy and then repeat 4 and 8 hours after the first dose.

2.2 Prevention of Postoperative Nausea and Vomiting

Caution: Do not administer a full prefilled syringe (4 mg dose) to pediatric patients less than 40 kg as this exceeds the recommended dose.

Important Preparation Instructions

Dilution of Ondansetron Injection is not required before administration to adult and pediatric patients. • Inspect Ondansetron Injection visually for particulate matter and discoloration before administration; discard if

Prevention of Postoperative Nausea and Vomiting						
Population	Recommended Single Dose	Administration Instructions	Timing of Administration			
Adults and pediatric patients older than 12 years of age	4 mg ^a	 May be administered intravenously or intramuscularly: Intravenously: infuse undiluted syringe contents (4 mg) over at least 30 seconds and preferably longer (over 2 to 5 minutes). Intramuscularly: inject undiluted syringe contents (4 mg) 	Administer immediately before induction of anesthesia, or postoperatively if the patient did not receive			
Pediatric patients 1 month to 12 years and more than 40 kg	4 mg	Infuse intravenously over at least 30 seconds and preferably longer (over 2 to 5 minutes).	prophylactic antiemetics and experiences nausea and/or vomiting occurring within 2 hours after surgery ^{b,c}			
Pediatric patients 1 month to 12 years and 40 kg or less	0.1 mg/kg	Infuse intravenously over at least 30 seconds and preferably longer (over 2 to 5 minutes).				

Few patients above 80 kg have been studied

6.2 additional control of a second intravenous dose of 4 mg ondansetron postoperatively in adult patients who received a 4 mg prophylactic dose does not provide additional control of nausea and vomiting [see *Clinical Studies* (14.3)].

patients with severe hepatic impairment (Child-Pugh score of 10 or greater), a single maximal daily dose of 8 mg nfused over 15 minutes beginning 30 minutes before the start of the emetogenic chemotherapy is recommend There is no experience beyond first-day administration of ondansetron in these patients [see Use in Specific Populations

DOSAGE FORMS AND STRENGTHS

etron Injection, USP 2 mg/mL is a clear, colorless, nonpyrogenic, sterile solution available as a 2 mL Prefilled disposable single-use syringe

CONTRAINDICATIONS Ondansetron Injection is contraindicated for patients known to have hypersensitivity (e.g., anaphylaxis) to this product or any of its components. Anaphylactic reactions have been reported in patients taking ondansetron [see Adverse Reactions (6.2)]

The concomitant use of apomorphine with ondansetron is contraindicated based on reports of profound hypotension and loss of consciousness when apomorphine was administered with ondansetron.

WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions Hypersensitivity reactions, including anaphylaxis and bronchospasm, have been reported in patients who have exhibited hypersensitivity to other selective 5-HT₃ receptor antagonists.

5.2 QT Prolongation

Ondansetron prolongs the QT interval in a dose-dependent manner [see Clinical Pharmacology (12.2)]. In addition, postmarketing cases of Torsade de Pointes have been reported in patients using ondansetron. Avoid Ondansetron in patients with congenital long QT syndrome. Electrocardiogram (ECG) monitoring is recommended in patients with electrolyte abnormalities (e.g., hypokalemia or hypomagnesemia), congestive heart failure, bradyarrhythmias, or patients taking other medicinal products that lead to QT prolongation.

5.3 Serotonin Syndrome

The development of serotonin syndrome has been reported with 5-HT₃ receptor antagonists. Most reports have been associated with concomitant use of serotonergic drugs (e.g., selective serotonin reuptake inhibitors (SSRIs), serotonin

Postmarketing Experience

Cardiovascula

Iniection-site reaction

Cold sensation

Fever

Pruritus

Paresthesia

Arrhythmias (including ventricular and supraventricular tachycardia, premature ventricular contractions, and atrial fibrillation), bradycardia, electrocardiographic alterations (including second-degree heart block, QT/QTc interval 8.2 Lactation prolongation, and ST segment depression), palpitations, and syncope. Rarely and predominantly with intravenous ndansetron, transient ECG changes, including QT/QTc interval prolongation have been reported [see Warnings and Precautions (5.2)].

Flushing: Rare cases of hypersensitivity reactions, sometimes severe (e.g., anaphylactic reactions, angioedema, bronchospasm, cardiopulmonary arrest, hypotension, laryngeal edema, laryngospasm, shock, shortness of breath, stridor) have also been reported. A positive lymphocyte transformation test to ondansetron has been reported, which suggests immunologic sensitivity to ondansetron.

Hepatobiliary

Local Reactions

Pain, redness, and burning at site of injection. Lower Respiratory

<u>Neurological</u>

intravenous infusion.

Urticaria, Stevens-Johnson syndrome, and toxic epidermal necrolysis.

setron Injection 4 mg Intravenous (n = 547)	Placebo (n = 547)
92 (17%)	77 (14%)
44 (8%)	37 (7%)
21 (4%)	18 (3%)
10 (2%)	6 (1%)
9 (2%)	8 (1%)
9 (2%)	3 (< 1%)
9 (2%)	2 (< 1%)
a 10 and a difference in the standard statement of the sector	

rse reactions: Rates of these reactions were not significantly different in the ondansetron and placebo group Patients were receiving multiple concomitant perioperative and postoperative medication

Pediatric Use: Rates of adverse reactions were similar in both the ondansetron and placebo groups in pediatric patients receiving ondansetron (a single 0.1-mg/kg dose for pediatric patients weighing 40 kg or less, or 4 mg for pediatric patients weighing more than 40 kg) administered intravenously over at least 30 seconds. Diarrhea was seen more frequently in patients taking Ondansetron (2%) compared with placebo (< 1%) in the 1-month to 24-month age-group. These patients were receiving multiple concomitant perioperative and postoperative medications.

The following adverse reactions have been identified during post-approval use of ondansetron. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. The reactions have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to ondansetron.

Liver enzyme abnormalities have been reported. Liver failure and death have been reported in patients with cancer receiving concurrent medications, including potentially hepatotoxic cytotoxic chemotherapy and antibiotics.

Oculogyric crisis, appearing alone, as well as with other dystonic reactions. Transient dizziness during or shortly after

Eve Disorders

Cases of transient blindness, predominantly during intravenous administration, have been reported. These cases of transient blindness were reported to resolve within a few minutes up to 48 hours. Transient blurred vision, in some cases associated with abnormalities of accommodation, has also been reported

DRUG INTERACTIONS

7.1 Drugs Affecting Cytochrome P-450 Enzymes Ondansetron does not appear to induce or inhibit the cytochrome P-450 drug-metabolizing enzyme system of the liver. Because ondansetron is metabolized by hepatic cytochrome P-450 drug-metabolizing enzymes (CYP3A4, CYP2D6, 8.7 CYP1A2), inducers or inhibitors of these enzymes may change the clearance and, hence, the half-life of ondansetron [see Clinical Pharmacology (12.3)]. On the basis of limited available data, no dosage adjustment is recommended for patients on these drugs.

7.2 Apomorphine

Based on reports of profound hypotension and loss of consciousness when apomorphine was administered with ondansetron, the concomitant use of apomorphine with ondansetron is contraindicated [see Contraindications (4)]. 10 7.3 Phenytoin, Carbamazepine, and Rifampin

In patients treated with potent inducers of CYP3A4 (i.e., phenytoin, carbamazepine, and rifampin), the clearance of ondansetron was significantly increased and ondansetron blood concentrations were decreased. However, on the basis of available data, no dosage adjustment for ondansetron is recommended for patients on these drugs [see Clinical Pharmacology (12.3)].

7.4 Tramadol

Although there are no data on pharmacokinetic drug interactions between ondansetron and tramadol, data from two small trials indicate that concomitant use of ondansetron may result in reduced analgesic activity of tramadol. Patients on concomitant ondansetron self-administered tramadol more frequently in these trials, leading to an increased cumulative dose in patient-controlled administration of tramadol.

7.5 Serotonergic Drugs

Serotonin syndrome (including altered mental status, autonomic instability, and neuromuscular symptoms) has been described following the concomitant use of 5-HT₃ receptor antagonists and other serotonergic drugs, including SSRIs and SNRIs [see Warnings and Precautions (5.3)].

7.6 Chemotherapy

In humans, carmustine, etoposide, and cisplatin do not affect the pharmacokinetics of ondansetron. In a crossover trial in 76 pediatric patients, intravenous ondansetron did not increase blood levels of high-dose

7.7 Temazepam

The coadministration of ondansetron had no effect on the pharmacokinetics and pharmacodynamics of temazepam. 7.8 Alfentanil and Atracurium

Ondansetron does not alter the respiratory depressant effects produced by alfentanil or the degree of neuromuscular blockade produced by atracurium. Interactions with general or local anesthetics have not been studied. 8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

<u>Risk Summary</u>

Published epidemiological studies on the association between ondansetron use and major birth defects have reported inconsistent findings and have important methodological limitations that preclude conclusions about the safety of ondansetron use in pregnancy (see Data). Available postmarketing data have not identified a drug-associated risk of miscarriage or adverse maternal outcomes. Reproductive studies in rats and rabbits did not show evidence of harm to the fetus when ondansetron was administered intravenously during organogenesis at approximately 3.6 and 2.9 times the maximum recommended human intravenous dose of 0.15 mg/kg given three times a day, based on body surface area, respectively (see Data).

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, miscarriages, or other adverse outcomes. In the US general population, the estimated background risk of major birth defects and miscarriages in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Human Data

Available data on ondansetron use in pregnant women from several published epidemiological studies preclude an assessment of a drug-associated risk of adverse fetal outcomes due to important methodological limitations. including the uncertainty of whether women who filled a prescription actually took the medication, the concomitant use of other medications or treatments, recall bias, and other unadjusted confounder

Ondansetron exposure in utero has not been associated with overall major congenital malformations in aggregate analyses. One large retrospective cohort study examined 1970 women who received a prescription for ondansetron during pregnancy and reported no association between ondansetron exposure and major congenital malformations. miscarriage, stillbirth, preterm delivery, infants of low birth weight, or infants small for gestational age.

Two large retrospective cohort studies and one case-control study have assessed ondansetron exposure in the first trimester and risk of cardiovascular defects with inconsistent findings. Relative risks (RR) ranged from 0.97 (95% CI 0.86 to 1.10) to 1.62 (95% CI 1.04, 2.54). A subset analysis in one of the cohort studies observed that ondansetron was specifically associated with cardiac septal defects (RR 2.05, 95% CI 1.19, 3.28); however this association was not confirmed in other studies.

Several studies have assessed ondansetron and the risk of oral clefts with inconsistent findings. A retrospective cohort study of 1.8 million pregnancies in the US Medicaid Database showed an increased risk of oral clefts among 88.467 pregnancies in which oral ondansetron was prescribed in the first trimester (RR 1.24, 95% CI 1.03, 1.48). but no such association was reported with intravenous ondansetron in 23.866 pregnancies (RR 0.95, 95% CI 0.63, 1.43). In the subgroup of women who received both forms of administration, the RR was 1.07 (95% CI 0.59, 1.93). Two case-control studies, using data from birth defects surveillance programs, reported conflicting associations between maternal use of ondansetron and isolated cleft palate (OR 1.6 [95% CI 1.1, 2.3] and 0.5 [95% CI 0.3, 1.0]). It is unknown whether ondansetron exposure in utero in the cases of cleft palate occurred during the time of palate formation (the palate is formed between the 6th and 9th weeks of pregnancy).

Animal Data

In embryo-fetal development studies in rats and rabbits, pregnant animals received intravenous doses of ondansetron up to 10 mg/kg/day and 4 mg/kg/day, respectively, during the period of organogenesis. With the exception of short periods of maternal weight loss and a slight increase in the incidence of early uterine deaths at the high dose level in rabbits, there were no significant effects of ondansetron on the maternal animals or the development of the offspring At doses of 10 mg/kg/day in rats and 4 mg/kg/day in rabbits, the maternal exposure margin was approximately 3 and 2.9 times the maximum recommended human oral dose of 0.15 mg/kg given three times a day, respectively based on body surface area.

No intravenous pre- and post-natal developmental toxicity study was performed with ondansetron. In an oral pre- and post-natal development study pregnant rats received oral doses of ondansetron up to 15 mg/kg/day from Day 17 of pregnancy to litter Day 21. With the exception of a slight reduction in maternal body weight gain, there were no effects upon the pregnant rats and the pre- and postnatal development of their offspring, including reproductive performance of the mated F1 generation.

Risk Summar

It is not known whether ondansetron is present in human milk. There are no data on the effects of Ondansetron Injection on the breastfed infant or the effects on milk production. However, it has been demonstrated that ondansetron is present in the milk of rats. When a drug is present in animal milk, it is likely that the drug will be present in human milk.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Ondansetron Injection and any potential adverse effects on the breast-fed infant from Ondansetron Injection or from the underlying maternal condition.

8.4 Pediatric Use

Little information is available about the use of ondansetron in pediatric surgical patients younger than 1 month [see Clinical Studies (14.2)]. Little information is available about the use of ondansetron in pediatric cancer patients younger than 6 months [see Clinical Studies (14.1), Dosage and Administration (2)].

The clearance of ondansetron in pediatric patients aged 1 month to 4 months is slower and the half-life is ~2.5- fold longer than patients who are aged > 4 to 24 months. As a precaution, it is recommended that patients younger than 4 months receiving this drug be closely monitored [see Clinical Pharmacology (12.3)].

8.5 Geriatric Use

Of the total number of subjects enrolled in cancer chemotherapy-induced and postoperative nausea and vomiting US- and foreign-controlled clinical trials, 862 were aged 65 years and older. No overall differences in safety or effectiveness were observed between subjects 65 years and older and younger subjects. A reduction in clearance and increase in elimination half-life were seen in patients older than 75 years compared with younger subjects [see Clinical Pharmacology (12.3)]. There were an insufficient number of patients older than 75 years of age and older in the clinical trials to permit safety or efficacy conclusions in this age-group. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. Dosage adjustment is not needed in patients over the age of 65.

8.6 Hepatic Impairment

In patients with severe hepatic impairment (Child-Pugh score of 10 or greater), clearance is reduced and apparent volume of distribution is increased with a resultant increase in plasma half-life [see Clinical Pharmacology (12.3)]. In such patients, a total daily dose of 8 mg should not be exceeded [see Dosage and Administration (2.3)].

Renal Impairment

Although plasma clearance is reduced in patients with severe renal impairment (creatinine clearance < 30 mL/min), no dosage adjustment is recommended [see Clinical Pharmacology (12.3)].

DRUG ABUSE AND DEPENDENCE

Animal studies have shown that ondansetron is not discriminated as a benzodiazepine nor does it substitute for benzodiazepines in direct addiction studies.

OVERDOSAG


There is no specific antidote for ondansetron overdose. Patients should be managed with appropriate supportive therapy. Individual intravenous doses as large as 150 mg and total daily intravenous doses as large as 252 mg have been inadvertently administered without significant adverse events. These doses are more than 10 times the recommended daily dose.

In addition to the adverse reactions listed above, the following events have been described in the setting of ondansetron overdose: "Sudden blindness" (amaurosis) of 2 to 3 minutes' duration plus severe constipation occurred in one patient that was administered 72 mg of ondansetron intravenously as a single dose. Hypotension (and faintness) occurred in another patient that took 48 mg of ondansetron hydrochloride tablets. Following infusion of 32 mg over only a 4-minute period, a vasovagal episode with transient second-degree heart block was observed. In all instances, the events resolved completely.

Pediatric cases consistent with serotonin syndrome have been reported after inadvertent oral overdoses of ondansetron (exceeding estimated ingestion of 5 mg/kg) in young children. Reported symptoms included somnolence. agitation, tachycardia, tachypnea, hypertension, flushing, mydriasis, diaphoresis, myoclonic movements, horizontal nystagmus, hyperreflexia, and seizure. Patients required supportive care, including intubation in some cases, with complete recovery without sequelae within 1 to 2 days.

11 DESCRIPTION

The active ingredient in Ondansetron Injection, USP is ondansetron hydrochloride, a selective blocking agent of the serotonin 5-HT₃ receptor type. Its chemical name is (±) 1, 2, 3, 9-tetrahydro-9-methyl-3-[(2-methyl-1H-imidazol-1-yl) methyll-4H-carbazol-4-one, monohydrochloride, dihydrate. It has the following structural formula:



The empirical formula is $C_{18}H_{19}N_3O \bullet HCI \bullet 2H_2O$, representing a molecular weight of 365.9 g/mol. Ondansetron HCl is a white to off-white powder that is soluble in water and normal saline Each 1 mL of aqueous solution contains 2 mg of ondansetron as the hydrochloride dihydrate: 9 mg of sodium chloride. USP: and 0.5 mg of citric acid monohydrate, USP and 0.25 mg of sodium citrate dihydrate, USP as buffers in Water for Iniection, USP.

ndansetron Injection, USP is a clear, colorless, nonpyrogenic, sterile solution for intravenous or intramuscular use. The pH of the injection solution is 3.3 to 4.0.

CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Ondansetron is a selective 5-HT₂ receptor antagonist. While ondansetron's mechanism of action has not been fully characterized, it is not a dopamine-receptor antagonist

12.2 Pharmacodynamics

In normal volunteers, single intravenous doses of 0.15 mg/kg of ondansetron had no effect on esophageal motility, gastric motility, lower esophageal sphincter pressure, or small intestinal transit time. In another trial in six normal male volunteers, a 16 mg dose infused over 5 minutes showed no effect of the drug on cardiac output, heart rate stroke volume, blood pressure, or ECG. Multiday administration of ondansetron has been shown to slow colonic transit in normal volunteers. Ondansetron has no effect on plasma prolactin concentrations. In a gender balanced pharmacodynamic trial (n = 56), ondansetron 4 mg administered intravenously or intramuscularly was dynamically similar in the prevention of nausea and vomiting using the ipecacuanha model of emesis. Cardiac Electrophysiology

QTc interval prolongation was studied in a double-blind, single intravenous dose, placebo- and positive-controlled, crossover trial in 58 healthy subjects. The maximum mean (95% upper confidence bound) difference in OTcF from placebo after baseline correction was 19.5 (21.8) ms and 5.6 (7.4) ms after 15-minute intravenous infusions of 32 mg and 8 mg Ondansetron, respectively. A significant exposure-response relationship was identified between ondansetron concentration and $\Delta\Delta$ OTcF. Using the established exposure-response relationship, 24 mg infused intravenously over 15 minutes had a mean predicted (95% upper prediction interval) $\Delta\Delta$ OTcF of 14.0 (16.3) ms. In contrast, 16 mg infused intravenously over 15 minutes using the same model had a mean predicted (95% upper prediction interva $\Delta\Delta\Delta$ QTcF of 9.1 (11.2) ms. In this study, the 8-mg dose infused over 15 minutes did not prolong the QT interval to any clinically relevant extent.

12.3 Pharmacokinetics

In normal adult volunteers, the following mean pharmacokinetic data have been determined following a single 0.15-mg/kg intravenous dose

Table 4. Pharmacokinetics in Normal Adult Volunteers

Age-group (years)	n	Peak Plasma Concentration (ng/mL)	Mean Elimination Half-life (h)	Plasma Clearance (L/h/kg)
19-40	11	102	3.5	0.381
61-74	12	106	4.7	0.319
≥ 75	11	170	5.5	0.262

A trial was performed in normal volunteers (n = 56) to evaluate the pharmacokinetics of a single 4-mg dose administered as a 5-minute infusion compared with a single intramuscular injection. Systemic exposure as measured b mean area under curve (AUC) were equivalent, with values of 156 [95% CI: 136, 180] and 161 [95% CI: 137, 190 ng•h/mL for intravenous and intramuscular groups, respectively. Mean peak plasma concentrations were 42.9 [95% CI: 33.8, 54.4] ng/mL at 10 minutes after intravenous infusion and 31.9 [95% CI: 26.3, 38.6] ng/mL at 41 minutes after intramuscular injection.

Distribution

ondansetron as measured in vitro was 70% to 76%, over the pharmacologic concentration range of 10 to 500 ng/mL. Circulating drug also distributes into erythrocytes.

Metabolism: Ondansetron is extensively metabolized in humans, with approximately 5% of a radiolabeled dose recovered as the parent compound from the urine. The primary metabolic pathway is hydroxylation on the indole ring followed by subsequent glucuronide or sulfate conjugation

CYP2D6 in ondansetron *in vivo* metabolism is relatively minor.

ondansetron disposition in vivo.

Although some nonconjugated metabolites have pharmacologic activity, these are not found in plasma at concentrations likely to significantly contribute to the biological activity of ondansetron. The metabolites are observed in the urine.

HCI+2H_C

In vitro metabolism studies have shown that ondansetron is a substrate for multiple human hepatic cytochrome P-450 enzymes, including CYP1A2, CYP2D6, and CYP3A4. In terms of overall ondansetron turnover, CYP3A4 plays a predominant role while formation of the major *in vivo* metabolites is apparently mediated by CYP1A2. The role of

The pharmacokinetics of intravenous ondansetron did not differ between subjects who were poor metabolizers of CYP2D6 and those who were extensive metabolizers of CYP2D6, further supporting the limited role of CYP2D6 in

Excretion: In adult cancer patients, the mean ondansetron elimination half-life was 4.0 hours, and there was no difference in the multidose pharmacokinetics over a 4-day period. In a dose-proportionality trial, systemic exposure to 32 mg of ondansetron was not proportional to dose as measured by comparing dose-normalized AUC values with an 8-mg dose. This is consistent with a small decrease in systemic clearance with increasing plasma concentrations.



920 ΝΟΙΤΟΞΙΝΙ ΝΟΑΤΞΣΝΑΟΝΟ

ONDANSETRON INJECTION USP



Specific Populations

Geriatric Patients: A reduction in clearance and increase in elimination half-life are seen in patients older than 75 years of age [see Use in Specific Populations (8.5)].

Pediatric Patients: Pharmacokinetic samples were collected from 74 cancer patients aged 6 to 48 months, who received a dose of 0.15 mg/kg of intravenous ondansetron every 4 hours for 3 doses during a safety and efficacy trial. These Median number of emetic episodes data were combined with sequential pharmacokinetics data from 41 surgery patients aged 1 month to 24 months, who received a single dose of 0.1 mg/kg of intravenous ondansetron prior to surgery with general anesthesia, and a population pharmacokinetic analysis was performed on the combined data set. The results of this analysis are ncluded in Table 5 and are compared with the pharmacokinetic results in cancer patients aged 4 to 18 years.

Table 5. Pharmacokinetics in Pediatric Cancer Patients Aged 1 Month to 18 Years

Subjects and Age-group	Ν	CL (L/h/kg)	Vd _{ss} (L/kg)	t _½ (h)
		Geon	netric Mean	Mean
Pediatric Cancer Patients 4 to 18 years	N = 21	0.599	1.9	2.8
Population PK Patients ^a 1 month to 48 months	N = 115	0.582	3.65	4.9

pulation PK (Pharmacokinetic) Patients: 64% cancer patients and 36% surgery patient

Based on the population pharmacokinetic analysis, cancer patients aged 6 to 48 months who receive a dose of 0.15 mg/kg of intravenous ondansetron every 4 hours for 3 doses would be expected to achieve a systemic exposure (AUC) consistent with the exposure achieved in previous pediatric trials in cancer patients (4 to 18 years) at similar

In a trial of 21 pediatric patients (3 to 12 years) who were undergoing surgery requiring anesthesia for a duration of 45 minutes to 2 hours, a single intravenous dose of ondansetron, 2 mg (3 to 7 years) or 4 mg (8 to 12 years), was administered immediately prior to anesthesia induction. Mean weight-normalized clearance and volume of distribution values in these pediatric surgical patients were similar to those previously reported for young adults. Mean terminal half-life was slightly reduced in pediatric patients (range, 2.5 to 3 hours) in comparison with adults (range, 3 to 3.5 hours).

In a trial of 51 pediatric patients (aged 1 month to 24 months) who were undergoing surgery requiring general anesthesia, a single intravenous dose of ondansetron, 0.1 or 0.2 mg/kg, was administered prior to surgery. As shown in Table 6, the 41 patients with pharmacokinetic data were divided into 2 groups, patients aged 1 month to 4 months and patients aged 5 to 24 months, and are compared with pediatric patients aged 3 to 12 years.

Table 6. Pharmacokinetics in Pediatric Surgery Patients Aged 1 Month to 12 Years

Subjects and Age-group	N	CL (L/h/kg)	Vd _{ss} (L/kg)	t _½ (h)
		Geome	tric Mean	Mean
Pediatric Surgery Patients 3 to 12 years	N = 21	0.439	1.65	2.9
Pediatric Surgery Patients 5 to 24 months	N = 22	0.581	2.3	2.9
Pediatric Surgery Patients 1 month to 4 months	N = 19	0.401	3.5	6.7

In general, surgical and cancer pediatric patients younger than 18 years tend to have a higher ondansetron clearance compared with adults leading to a shorter half-life in most pediatric patients. In patients aged 1 month to 4 months, a longer half-life was observed due to the higher volume of distribution in this age-group.

In a trial of 21 pediatric cancer patients (aged 4 to 18 years) who received three intravenous doses of 0.15 mg/kg of ondansetron at 4-hour intervals, patients older than 15 years exhibited ondansetron pharmacokinetic parameters similar to those of adults.

Patients with Renal Impairment: Due to the very small contribution (5%) of renal clearance to the overall clearance, renal impairment was not expected to significantly influence the total clearance of ondansetron. However, ondan setron mean plasma clearance was reduced by about 41% in patients with severe renal impairment (creatinine clearance < 30 mL/min). This reduction in clearance is variable and was not consistent with an increase in half-life [see Use in Specific Populations (8.7)].

Patients with Hepatic Impairment: In patients with mild-to-moderate hepatic impairment, clearance is reduced 2-fold and mean half-life is increased to 11.6 hours compared with 5.7 hours in those without hepatic impairment. In patients with severe hepatic impairment (Child-Pugh score of 10 or greater), clearance is reduced 2-fold to 3-fold and apparent volume of distribution is increased with a resultant increase in half-life to 20 hours [see Dosage and Administration (2.3), Use in Specific Populations (8.6)].

Drug Interaction Studies

CYP 3A4 Inducers: Ondansetron elimination may be affected by cytochrome P-450 inducers. In a pharmacokinetic trial of 16 epileptic patients maintained chronically on CYP3A4 inducers, carbamazepine, or phenytoin, a reduction in AUC. C_{max}, and t₁₂₂ of ondansetron was observed. This resulted in a significant increase in the clearance of ondansetron. In a pharmacokinetic study of 10 healthy subjects receiving a single-dose intravenous dose of ondansetron 8 mg after 600 mg rifampin once daily for five days, the AUC and the t_{y_0} of ondansetron were reduced by 48% and 46%, respectively. These changes in ondansetron exposure with CYP3A4 inducers are not thought to be clinically relevant [see Drug Interactions (7.3)]

Chemotherapeutic Agents: Carmustine, etoposide, and cisplatin do not affect the pharmacokinetics of ondansetron [see Drug Interactions (7.6)].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenic effects were not seen in 2-year studies in rats and mice with oral ondansetron doses up to 10 and 30 mg/kg per day, respectively (approximately 3.6 and 5.4 times the recommended human intravenous dose of 0.15 mg/kg given three times a day, based on body surface area). Ondansetron was not mutagenic in standard tests for mutagenicity

Oral administration of ondansetron up to 15 mg/kg per day (approximately 3.8 times the recommended human intravenous dose, based on body surface area) did not affect fertility or general reproductive performance of male "Efficacy based on "all-patients-treated" analysis. and female rats

14 CLINICAL STUDIES

The clinical efficacy of ondansetron hydrochloride, the active ingredient of Ondansetron Injection, was assessed in clinical trials as described below.

14.1 Chemotherapy-induced Nausea and Vomiting

Adults

In a double-blind trial of three different dosing regimens of Ondansetron Injection, 0.015 mg/kg, 0.15 mg/kg, and 0.30 mg/kg, each given three times during the course of cancer chemotherapy, the 0.15-mg/kg dosing regimen was more effective than the 0.015-mg/kg dosing regimen. The 0.30-mg/kg dosing regimen was not shown to be more effective than the 0.15-mg/kg dosing regimen.

Cisplatin-based Chemotherapy: In a double-blind trial in 28 patients, Ondansetron Injection (three 0.15-mg/kg doses) was significantly more effective than placebo in preventing nausea and vomiting induced by cisplatin-based chemotherapy. Therapeutic response was as shown in Table 7.

Table 7. Therapeutic Response in Prevention of Chemotherapy-Induced Nausea and Vomiting in Single-day Cisplatin Therapy^a in Adults

	Ondansetron Injection (0.15 mg/kg x 3)	Placebo	<i>P</i> Value [♭]
Number of patients	14	14	
Treatment response 0 Emetic episodes 1-2 Emetic episodes 3-5 Emetic episodes More than 5 emetic episodes/rescued	2 (14%) 8 (57%) 2 (14%) 2 (14%)	0 (0%) 0 (0%) 1 (7%) 13 (93%)	0.001

Median time to first emetic episod

Median nausea scores (0-100)^d Global satisfaction with control or

nausea and vomiting (0-100)^e

^b Efficacy based on "all-patients-treated" analysis. ^aVisual analog scale assessment of nausea: 0 = no nausea, 100 = nausea as bad as it can be. ^eVisual analog scale assessment of satisfaction: 0 = not at all satisfied, 100 = totally satisfied

Ondansetron injection (0.15-mg/kg x 3 doses) was compared with metoclopramide (2 mg/kg x 6 doses) in a singleblind trial in 307 patients receiving cisplatin \geq 100 mg/m² with or without other chemotherapeutic agents. Patients received the first dose of ondansetron or metoclopramide 30 minutes before cisplatin. Two additional ondansetron doses were administered 4 and 8 hours later, or five additional metoclopramide doses were administered 2, 4, 7, 10, and 13 hours later. Cisplatin was administered over a period of 3 hours or less. Episodes of vomiting and retching were tabulated over the period of 24 hours after cisplatin. The results of this trial are summarized in Table 8.

Table 8. Therapeutic Response in Prevention of Vomiting Induced by Cisplatin (\geq 100 mg/m²)

	Ondansetron Injection	Metoclopramide	P Value
	0.15 mg/kg x 3	2 mg/kg x 6	
Number of patients in efficacy population	136	138	
Treatment response 0 Emetic episodes 1-2 Emetic episodes 3-5 Emetic episodes More than 5 emetic episodes/rescued	54 (40%) 34 (25%) 19 (14%) 29 (21%)	41 (30%) 30 (22%) 18 (13%) 49 (36%)	
Comparison of treatments with respect to 0 Emetic episodes More than 5 emetic episodes/rescued	54/136 29/136	41/138 49/138	0.083 0.009
Median number of emetic episodes	1	2	0.005
Median time to first emetic episode (h)	20.5	4.3	< 0.001
Global satisfaction with control of nausea and vomiting (0-100) ^b	85	63	0.001
Acute dystonic reactions	0	8	0.005
Akathisia	0	10	0.002

n-	0.15-mg/kg doses) in 20 patients receiv	ing cyclophosphamide (500	to 600 mg/m ²) chem	otherapy, Ondansetron	n Pediatric Patients Aged 2 to 12 Years			
ne fe	Injection was significantly more effective t in Table 9.		-		Treatment Response Over 24 Hours	Ondansetron n (%)	Placebo n (%)	P Value
ed		Gable 9. Therapeutic Response in Prevention of Chemotherapy-Induced Nausea and Vomiting in Single-day Cyclophosphamide Therapy ^a in Adults			Study 1 Number of patients	205	210	
nt. Id nd		Ondansetron Injection	Placebo	<i>P</i> Value⁵	0 Emetic episodes Failure ^a	140 (68%) 65 (32%)	82 (39%) 128 (61%)	≤ 0.001
	Number of patients	(0.15 mg/kg x 3) 10	10		Study 2 Number of patients	112	110	≤ 0.001
ial C,	Treatment response 0 Emetic episodes	7 (70%)	0 (0%)	0.001	0 Emetic episodes Failureª	68 (61%) 44 (39%)	38 (35%) 72 (65%)	_ 0.001
n. ng %, nt	1-2 Emetic episodes 3-5 Emetic episodes More than 5 emetic episodes/rescued	0 (0%) 2 (20%) 1 (10%)	2 (20%) 4 (40%) 4 (40%)	0.131	Study 3 Number of patients 0 Emetic episodes Failure ^a	206 123 (60%) 83 (40%)	206 96 (47%) 110 (53%)	≤ 0.01
on	Median number of emetic episodes	0	4	0.008	Nausea assessments ^b Number of patients	185	191	≤ 0.01
	Median time to first emetic episode (h)	Undefined ^c	8.79		^a Failure was one or more emetic episodes, rescued	119 (64%)	99 (52%)	
nd	Median nausea scores (0-100) ^d	0	60	0.001	^b Nausea measured as none, mild, or severe.			
of	Global satisfaction with control of nausea and vomiting (0-100) ^e	100	52	0.008	A double-blind, multicenter, pla 24 months who were undergo males; 64% were white, 15% we	ing routine surgery under ere black, 13% were Ameri	general anesthesia. Sever can Hispanic, 2% were Asia	nty-five percent (75%) were n, and 6% were "other race"
'n	^a Chemotherapy consisted of cyclophosphamide in all patients,	plus other agents, including fluorour	acil, doxorubicin, methotrexat	e, and vincristine. There was no	patients. A single 0.1-mg/kg intr			

difference between treatments in the type of chemotherapy that would account for differences in response Median undefined since at least 50% of patients did not have any emetic episodes. ⁴Visual analog scale assessment of nausea: 0 = no nausea, 100 = nausea as bad as it can be.
^eVisual analog scale assessment of satisfaction: 0 = not at all satisfied, 100 = totally satisfied

Re-treatment: In uncontrolled trials, 127 patients receiving cisplatin (median dose, 100 mg/m²) and ondansetron who had two or fewer emetic episodes were re-treated with ondansetron and chemotherapy, mainly cisplatin, for a total of 269 re-treatment courses (median: 2; range, 1 to 10). No emetic episodes occurred in 160 (59%), and two or fewer emetic episodes occurred in 217 (81%) re-treatment courses.

Four open-label, noncomparative (one US, three foreign) trials have been performed with 209 pediatric cancer patients aged 4 to 18 years given a variety of cisplatin or noncisplatin regimens. In the three foreign trials, the initial dose of Ondansetron Injection ranged from 0.04 to 0.87 mg/kg for a total dose of 2.16 to 12 mg. This was followed by the oral administration of ondansetron ranging from 4 to 24 mg daily for 3 days. In the US trial, Ondansetron njection was administered intravenously (only) in three doses of 0.15 mg/kg each for a total daily dose of 7.2 to 39 mg. In these trials, 58% of the 196 evaluable patients had a complete response (no emetic episodes) on Day 1. Thus, prevention of vomiting in these pediatric patients was essentially the same as for patients older than 18 years.

An open-label, multicenter, noncomparative trial has been performed in 75 pediatric cancer patients aged 6 to 48 months receiving at least one moderately or highly emetogenic chemotherapeutic agent. Fifty-seven percent (57%) were females; 67% were white, 18% were American Hispanic, and 15% were black patients. Ondansetron Injection was administered intravenously over 15 minutes in three doses of 0.15 mg/kg. The first dose was administered 30 minutes before the start of chemotherapy; the second and third doses were administered 4 and 8 hours after the first dose, respectively. Eighteen patients (25%) received routine prophylactic dexamethasone (i.e., not given as rescue). Of the 75 evaluable patients, 56% had a complete response (no emetic episodes) on Day 1. Thus, prevention of vomiting in these pediatric patients was comparable to the prevention of vomiting in patients aged 4 years and older.

14.2 Prevention of Postoperative Nausea and/or Vomiting

Adult surgical patients who received ondansetron immediately before the induction of general balanced anesthesia (barbiturate: thiopental, methohexital, or thiamylal; opioid: alfentanil or fentanyl; nitrous oxide; neuromuscular

Table 7. Therapeutic Response in Prevention of Chemotherapy-Induced Nausea and Vomiting in Single-day Cisplatin Therapy^a in Adults (Continued)

	Ondansetron Injection (0.15 mg/kg x 3)	Placebo	<i>P</i> Value ^b					
es	1.5	Undefined ^c						
de (h)	11.6	2.8	0.001					
	3	59	0.034					
f	96	10.5	0.009					

Chemotherapy was high dose (100 and 120 mg/m²; Ondansetron Injection n = 6, placebo n = 5) or moderate dose (50 and 80 mg/m²; Ondansetron Injection r = 8, placed on r = 9. Other chemotherapeutic agents included fluorunaril, doorubicin, and cyclophosphamide. There was no difference betw the types of chemotherapy that would account for differences in response.

blockade: succinvlcholine/curare and/or vecuronium or atracurium; and supplemental isoflurane) were evaluated in two double-blind US trials involving 554 patients. Ondansetron Injection (4 mg) intravenous given over 2 to 5 minutes was significantly more effective than placebo. The results of these trials are summarized in Table 10.

Table 10. Therapeutic Response in Prevention of Postoperative Nausea and Vomiting in Adult Patients					
	Ondansetron 4 mg Intravenous	Placebo	P Value		
Study 1					
Emetic episodes: Number of patients Treatment response over 24-h postoperative period 0 Emetic episodes 1 Emetic episode More than 1 emetic episode/rescued	136 103 (76%) 13 (10%) 20 (15%)	139 64 (46%) 17 (12%) 58 (42%)	< 0.001		
Nausea assessments: Number of patients No nausea over 24-h postoperative period	134 56 (42%)	136 39 (29%)			
Study 2					
Emetic episodes: Number of patients Treatment response over 24-h postoperative period 0 Emetic episodes 1 Emetic episode More than 1 emetic episode/rescued	136 85 (63%) 16 (12%) 35 (26%)	143 63 (44%) 29 (20%) 51 (36%)	0.002		
Nausea assessments: Number of patients No nausea over 24-h postoperative period	125 48 (38%)	133 42 (32%)			

The populations in Table 10 consisted mainly of females undergoing laparoscopic procedures.

In a placebo-controlled trial conducted in 468 males undergoing outpatient procedures, a single 4-mg intravenous ondansetron dose prevented postoperative vomiting over a 24-hour period in 79% of males receiving drug compared with 63% of males receiving placebo (P < 0.001).

Two other placebo-controlled trials were conducted in 2,792 patients undergoing major abdominal or gynecological surgeries to evaluate a single 4-mg or 8-mg intravenous ondansetron dose for prevention of postoperative nausea and vomiting over a 24-hour period. At the 4-mg dosage, 59% of patients receiving ondansetron versus 45% receiving placebo in the first trial

(P < 0.001) and 41% of patients receiving ondansetron versus 30% receiving placebo in the second 0.001) experienced no emetic episodes. No additional benefit was observed in patients who received ondansetron 8 mg compared with patients who received intravenous ondansetron 4 mg.

Three double-blind, placebo-controlled trials have been performed (one US, two foreign) in 1,049 male and female * Failure was one or more emetic episodes, rescued, or withdrawn. patients (aged 2 to 12 years) undergoing general anesthesia with nitrous oxide. The surgical procedures included 16 HOW SUPPLIED/STORAGE AND HANDLING tonsillectomy with or without adenoidectomy, strabismus surgery, herniorrhaphy, and orchidopexy. Patients were randomized to either single intravenous doses of ondansetron (0.1 mg/kg for pediatric patients weighing 40 kg or less, 4 mg for pediatric patients weighing more than 40 kg) or placebo. Study drug was administered over at least 30 seconds, immediately prior to or following anesthesia induction. Ondansetron was significantly more effective than placebo in preventing nausea and vomiting. The results of these trials are summarized in Table 11.

Table 11. Therapeutic Response in Prevention of Postoperative Nausea and Vomiting in

anesthesia was statistically significantly more effective than placebo in preventing vomiting. In the placebo group, 28% of patients experienced vomiting compared with 11% of subjects who received ondansetron ($P \le 0.01$). Overall, 32 (10%) of placebo patients and 18 (5%) of patients who received ondansetron received antiemetic rescue medication(s) or prematurely withdrew from the trial.

14.3 Prevention of Further Postoperative Nausea and Vomiting

Adult surgical patients receiving general balanced anesthesia (barbiturate: thiopental, methohexital, or thiamylal; opioid: alfentanil or fentanyl: nitrous oxide: neuromuscular blockade: succinvlcholine/curare and/or vecuronium or atracurium; and supplemental isoflurane) who received no prophylactic antiemetics and who experienced nausea and/or vomiting within 2 hours postoperatively were evaluated in two double-blind US trials involving 441 patients Patients who experienced an episode of postoperative nausea and/or vomiting were given Ondansetron Injection (4 mg) intravenously over 2 to 5 minutes, and this was significantly more effective than placebo. The results of these trials are summarized in Table 12.

Table 12. Therapeutic Response in Prevention of Further P	ostoperative
Neurop and Vensiting in Adult Detionte	

Nausea and Vomiting in Adult Patients			
	Ondansetron 4 mg Intravenous	Placebo	P Value
Study 1			
Emetic episodes:			
Number of patients	104	117	
Treatment response 24 h after study drug			
0 Emetic episodes	49 (47%)	19 (16%)	< 0.001
1 Emetic episode	12 (12%)	9 (8%)	
More than 1 emetic episode/rescued	43 (41%)	89 (76%)	
Median time to first emetic episode (min) ^a	55.0	43.0	

Table 12. Therapeutic Response in Prevention of Further Postoperative a in Adult Patients (C

Nausea and Vomiting in Adult Patients (Continued)				
	Ondansetron 4 mg Intravenous	Placebo	P Value	
Study 1				
Nausea assessments: Number of patients Mean nausea score over 24-h postoperative period ^b	98 1.7	102 3.1		
Study 2				
Emetic episodes: Number of patients Treatment response 24 h after study drug 0 Emetic episodes 1 Emetic episode More than 1 emetic episode/rescued Median time to first emetic episode (min) ^a	112 49 (44%) 14 (13%) 49 (44%) 60.5	108 28 (26%) 3 (3%) 77 (71%) 34.0	0.006	
Nausea assessments: Number of patients Mean nausea score over 24-h postoperative period ^b	105 1.9	85 2.9		

After administration of study drug.

Nausea measured on a scale of 0-10 with 0 = no nausea, 10 = nausea as bad as it can be.

The populations in Table 12 consisted mainly of women undergoing laparoscopic procedures. Repeat Dosing in Adults: In patients who do not achieve adequate control of postoperative nausea and vomiting following a single, prophylactic, preinduction, intravenous dose of ondansetron 4 mg, administration of a second intravenous dose of ondansetron 4 mg postoperatively does not provide additional control of nausea and vomiting.

One double-blind, placebo-controlled, US trial was performed in 351 male and female outpatients (aged 2 to 12 years) who received general anesthesia with nitrous oxide and no prophylactic antiemetics. Surgical procedures were unrestricted. Patients who experienced two or more emetic episodes within 2 hours following discontinuation of nitrous oxide were randomized to either single intravenous doses of ondansetron (0.1 mg/kg for pediatric patients weighing 40 kg or less, 4 mg for pediatric patients weighing more than 40 kg) or placebo administered over at least 30 seconds

Ondansetron was significantly more effective than placebo in preventing further episodes of nausea and vomiting. The results of the trial are summarized in Table 13.

Table 13. Therapeutic Response in Prevention of Further Postoperative Nausea and Vomiting in Pediatric Patients Aged 2 to 12 Years

cond trial ($P =$	Treatment Response Over 24 Hours	Ondansetron n (%)	Placebo n (%)	P Value
ed intravenous	Number of patients 0 Emetic episodes Failure ^a	180 96 (53%) 84 (47%)	171 29 (17%) 142 (83%)	≤ 0.001

Ondansetron Injection, USP 2 mg/mL is available as

Product Code	Unit of Sale	Strength	Each		
RF796320	NDC 76045-216-20 Unit of 24	4 mg/2 mL (2 mg/mL)	NDC 76045-216-00 2 mL Prefilled Disposable Single Use Syringe This product contains an RFID.		
796320	NDC 76045-103-20 Unit of 24	4 mg/2 mL (2 mg/mL)	NDC 76045-103-00 2 mL Prefilled Disposable Single Use Syringe		

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature.] Product may also be stored in a refrigerator 2°C to 8°C (36°F to 46°F).

Protect from light.


DO NOT DILUTE FOR IV PUSH.

Do NOT place syringe on a Sterile Field.

INSTRUCTIONS FOR USE

Figure 1: Outer Packaging and Prefilled Syringe

Outer Package




NOTES:

- Do not introduce any other fluid into the syringe at any time.
- Do not dilute for IV push.
- Do not re-sterilize the syringe
- Do not use this product on a sterile field.
- This product is for single dose only.
- 1. Inspect the outer packaging (blister pack) to confirm the integrity of the packaging. Do not use if the blister pack or the prefilled syringe has been damaged
- Remove the syringe from the outer packaging. (See Figure 2)

Figure 2



3. Visually inspect the syringe. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. 4. Twist off the syringe tip cap. Do not remove the label around the luer lock collar. (See Figure 3)



5. Expel air bubble(s). Adjust the dose (if applicable).

- 6. Administer the dose ensuring that pressure is maintained on the plunger rod during the entire administration.
- 7. Discard the used syringe into an appropriate receptacle.

For more information concerning this drug, please call Fresenius Kabi USA, LLC at 1-800-551-7176.

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. 17 PATIENT COUNSELING INFORMATION

QT Prolongation Patients should be informed that Ondansetron Injection may cause serious cardiac arrhythmias, such as QT prolongation. Patients should be instructed to tell their healthcare provider right away if they perceive a change in their heart rate, if they feel lightheaded, or if they have a syncopal episode.

Patients should be informed that the chances of developing severe cardiac arrhythmias, such as QT prolongation and Torsade de Pointes are higher in the following people:

- Patients with a personal or family history of abnormal heart rhythms, such as congenital long QT syndrome; Patients who take medications, such as diuretics, which may cause electrolyte abnormalities;
- Patients with hypokalemia or hypomagnesemia

Ondansetron Injection should be avoided in these patients, since they may be more at risk for cardiac arrhythmias, such as QT prolongation and Torsade de Pointes [see Warnings and Precautions (5.2)]. Hypersensitivity Reactions

Inform patients that Ondansetron Injection may cause hypersensitivity reactions, some as severe as anaphylaxis and bronchospasm. The patient should report any signs and symptoms of hypersensitivity reactions, including fever, chills, rash, or breathing problems [see Warnings and Precautions (5.1)].

Masking of Progressive Ileus and Gastric Distension

Inform patients following abdominal surgery or those with chemotherapy-induced nausea and vomiting that Ondansetron Injection may mask signs and symptoms of bowel obstruction. Instruct patients to immediately report any signs or symptoms consistent with a potential bowel obstruction to their healthcare provider [see Warnings and Precautions (5.4)1.

Drug Interactions

- Instruct the patient to report the use of all medications, especially apomorphine, to their healthcare provider. Concomitant use of apomorphine and Ondansetron may cause a significant drop in blood pressure and loss of
- Advise patients of the possibility of serotonin syndrome with concomitant use of Ondansetron and another serotonergic agent, such as medications to treat depression and migraines. Advise patients to seek immediate medical attention if the following symptoms occur: changes in mental status, autonomic instability, neuromuscular symptoms with or without gastrointestinal symptoms [see Warnings and Precautions (5.3)].

The brand names mentioned in this document are the trademarks of their respective owners.



U.S. Patents 9,731,082 and 10,661,018

Lake Zurich, IL 60047

www.fresenius-kabi.com/us

451608A