

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use ROCURONIUM BROMIDE INJECTION safely and effectively. See full prescribing information for ROCURONIUM BROMIDE INJECTION.

ROCURONIUM BROMIDE injection, for intravenous use Initial U.S. Approval: 1994

RECENT MAJOR CHANGES

Dosage and Administration Important Dosing and Administration Information (2.1) 07/2018
Warnings and Precautions Risk of Death due to Medication Errors (5.3) 07/2018

INDICATIONS AND USAGE

Rocuronium Bromide Injection is a nondepolarizing neuromuscular blocking agent indicated as an adjunct to general anesthesia to facilitate both rapid sequence and routine tracheal intubation, and to provide skeletal muscle relaxation during surgery or mechanical ventilation. (1)

DOSAGE AND ADMINISTRATION

To be administered only by experienced clinicians or adequately trained individuals supervised by an experienced clinician familiar with the use, actions, characteristics, and complications of neuromuscular blocking agents. (2.1)

- Individualize the dose for each patient. (2.1)
- Peripheral nerve stimulator recommended for determination of drug response and need for additional doses, and to evaluate recovery. (2.1)
- Store Rocuronium Bromide Injection with cap and ferrule intact and in a manner that minimizes the possibility of selecting the wrong product. (2.1)
- Tracheal intubation: Recommended initial dose is 0.6 mg/kg. (2.2)
- Rapid sequence intubation: 0.6 to 1.2 mg/kg. (2.3)
- Maintenance doses: Guided by response to prior dose, not administered until recovery is evident. (2.4)
- Continuous infusion: Initial rate of 10 to 12 mcg/kg/min. Start only after early evidence of spontaneous recovery from an intubating dose. (2.5)

DOSAGE FORMS AND STRENGTHS

- 5 mL multiple dose vials containing 50 mg rocuronium bromide injection (10 mg/mL). (3)
- 10 mL multiple dose vials containing 100 mg rocuronium bromide injection (10 mg/mL). (3)

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE	7.3
2 DOSAGE AND ADMINISTRATION	7.4
2.1 Important Dosing and Administration Information	7.5
2.2 Dose for Tracheal Intubation	7.6
2.3 Rapid Sequence Intubation	7.7
2.4 Maintenance Dosing	7.8
2.5 Use by Continuous Infusion	7.9
2.6 Dosage in Specific Populations	7.10
2.7 Preparation for Administration of Rocuronium Bromide Injection	7.11
3 DOSAGE FORMS AND STRENGTHS	8
4 CONTRAINDICATIONS	8.1
5 WARNINGS AND PRECAUTIONS	8.2
5.1 Appropriate Administration and Monitoring	8.3
5.2 Anaphylaxis	8.4
5.3 Risk of Death due to Medication Errors	8.5
5.4 Need for Adequate Anesthesia	8.6
5.5 Residual Paralysis	8.7
5.6 Long-Term Use in an Intensive Care Unit	8.8
5.7 Malignant Hyperthermia (MH)	8.9
5.8 Prolonged Circulation Time	8.6
5.9 QT Interval Prolongation	8.7
5.10 Conditions/Drugs Causing Potentiation of, or Resistance to, Neuromuscular Block	8.8
5.11 Incompatibility with Alkaline Solutions	8.1
5.12 Increase in Pulmonary Vascular Resistance	8.2
5.13 Use In Patients with Myasthenia	8.3
5.14 Extravasation	8.7
6 ADVERSE REACTIONS	8
6.1 Clinical Trials Experience	8.1
6.2 Postmarketing Experience	8.2
7 DRUG INTERACTIONS	8
7.1 Antibiotics	8.1
7.2 Anticonvulsants	8.2

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Rocuronium Bromide Injection is indicated for inpatients and outpatients as an adjunct to general anesthesia to facilitate both rapid sequence and routine tracheal intubation, and to provide skeletal muscle relaxation during surgery or mechanical ventilation.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosing and Administration Information
Rocuronium Bromide Injection is for intravenous use only. This drug should only be administered by experienced clinicians or trained individuals supervised by an experienced clinician familiar with the use, actions, characteristics, and complications of neuromuscular blocking agents. Doses of Rocuronium Bromide Injection should be individualized and a peripheral nerve stimulator should be used to monitor drug effect, need for additional doses, adequacy of spontaneous recovery or antagonism, and to decrease the complications of overdosage if additional doses are administered.

The dosage information which follows is derived from studies based upon units of drug per unit of body weight. It is intended to serve as an initial guide to clinicians familiar with other neuromuscular blocking agents to acquire experience with Rocuronium Bromide Injection.

In patients in whom potentiation of, or resistance to, neuromuscular block is anticipated, a dose adjustment should be considered. [see *Dosage and Administration (2.6), Warnings and Precautions (5.10, 5.13), Drug Interactions (7.2, 7.3, 7.4, 7.5, 7.6, 7.8, 7.10), and Use in Specific Populations (8.6)*].

Risk of Medication Errors: Accidental administration of neuromuscular blocking agents may be fatal. Store Rocuronium Bromide Injection with the cap and ferrule intact and in a manner that minimizes the possibility of selecting the wrong product [see *Warnings and Precautions (5.3)*].

2.2 Dose for Tracheal Intubation

The recommended initial dose of Rocuronium Bromide Injection, regardless of anesthetic technique, is 0.6 mg/kg. Neuromuscular block sufficient for intubation (80% block or greater) is attained in a median (range) time of 1 (0.4 to 6) minute(s) and most patients have intubation completed within 2 minutes. Maximum blockade is

CONTRAINDICATIONS

- Hypersensitivity (e.g., anaphylaxis) to rocuronium bromide or other neuromuscular blocking agents. (4)

WARNINGS AND PRECAUTIONS

- Appropriate Administration and Monitoring:** Use only if facilities for intubation, mechanical ventilation, oxygen therapy, and an antagonist are immediately available. (5.1)
- Anaphylaxis:** Severe anaphylaxis has been reported. Consider cross-reactivity among neuromuscular blocking agents. (5.2)
- Risk of Death due to Medication Errors:** Accidental administration can cause death. (5.3)
- Need for Adequate Anesthesia:** Must be accompanied by adequate anesthesia or sedation. (5.4)
- Residual Paralysis:** Consider using a reversal agent in cases where residual paralysis is more likely to occur. (5.5)

ADVERSE REACTIONS

Most common adverse reactions (2%) are transient hypotension and hypertension. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Succinylcholine:** Use before succinylcholine has not been studied. (7.11)
- Nondepolarizing muscle relaxants:** Interactions have been observed. (7.7)
- Enhanced Rocuronium Bromide Injection activity possible:** Inhalation anesthetics (7.3), certain antibiotics (7.1), quinidine (7.10), magnesium (7.6), lithium (7.4), local anesthetics (7.5), procainamide. (7.8)
- Reduced Rocuronium Bromide Injection activity possible:** Anticonvulsants. (7.2)

USE IN SPECIFIC POPULATIONS

- Labor and Delivery:** Not recommended for rapid sequence induction in patients undergoing Cesarean section. (8.2)
- Pediatric Use:** Onset time and duration will vary with dose, age, and anesthetic technique. Not recommended for rapid sequence intubation in pediatric patients. (8.4)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 9/2021

7.3	Inhalation Anesthetics
7.4	Lithium Carbonate
7.5	Local Anesthetics
7.6	Magnesium
7.7	Nondepolarizing Muscle Relaxants
7.8	Procainamide
7.9	Propofol
7.10	Quinidine
7.11	Succinylcholine
8 USE IN SPECIFIC POPULATIONS	
8.1	Pregnancy
8.2	Labor and Delivery
8.4	Pediatric Use
8.5	Geriatric Use
8.6	Patients with Hepatic Impairment
8.7	Patients with Renal Impairment
10 OVERDOSAGE	
11 DESCRIPTION	
12 CLINICAL PHARMACOLOGY	
12.1	Mechanism of Action
12.2	Pharmacodynamics
12.3	Pharmacokinetics
13 NONCLINICAL TOXICOLOGY	
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
14 CLINICAL STUDIES	
14.1	Adult Patients
14.2	Geriatric Patients
14.3	Pediatric Patients
16 HOW SUPPLIED/STORAGE AND HANDLING	
17 PATIENT COUNSELING INFORMATION	

*Sections or subsections omitted from the full prescribing information are not listed.

achieved in most patients in less than 3 minutes. This dose may be expected to provide 31 (15 to 95) minutes of clinical relaxation under opioid/nitrous oxide/oxygen anesthesia. Under halothane, isoflurane, and enflurane anesthesia, some extension of the period of clinical relaxation should be expected [see *Drug Interactions (7.3)*].

A lower dose of Rocuronium Bromide Injection (0.45 mg/kg) may be used. Neuromuscular block sufficient for intubation (80% block or greater) is attained in a median (range) time of 1.3 (0.8 to 6.2) minute(s), and most patients have intubation completed within 2 minutes. Maximum blockade is achieved in most patients in less than 4 minutes. This dose may be expected to provide 22 (12 to 31) minutes of clinical relaxation under opioid/nitrous oxide/oxygen anesthesia. Patients receiving this low dose of 0.45 mg/kg who achieve less than 90% block (about 16% of these patients) may have a more rapid time to 25% recovery, 12 to 15 minutes.

A large bolus dose of 0.9 or 1.2 mg/kg can be administered under opioid/nitrous oxide/oxygen anesthesia without adverse effects to the cardiovascular system [see *Clinical Pharmacology (12.2)*].

2.3 Rapid Sequence Intubation

In appropriately premedicated and adequately anesthetized patients, Rocuronium Bromide Injection 0.6 to 1.2 mg/kg will provide excellent or good intubating conditions in most patients in less than 2 minutes [see *Clinical Studies (14.1)*].

2.4 Maintenance Dosing

Maintenance doses of 0.1, 0.15, and 0.2 mg/kg Rocuronium Bromide Injection, administered at 25% recovery of control T₁ (defined as 3 twitches of train-of-four), provide a median (range) of 12 (2 to 31), 17 (6 to 50), and 24 (7 to 69) minutes of clinical duration under opioid/nitrous oxide/oxygen anesthesia [see *Clinical Pharmacology (12.2)*]. In all cases, dosing should be guided based on the clinical duration following initial dose or prior maintenance dose and not administered until recovery of neuromuscular function is evident. A clinically insignificant cumulation of effect with repetitive maintenance dosing has been observed [see *Clinical Pharmacology (12.2)*].

2.5 Use by Continuous Infusion

Infusion at an initial rate of 10 to 12 mcg/kg/min of Rocuronium Bromide Injection should be initiated only after early evidence of spontaneous recovery from an intubating dose. Due to rapid redistribution [see

Clinical Pharmacology (12.3)] and the associated rapid spontaneous recovery, initiation of the infusion after substantial return of neuromuscular function (more than 10% of control T₁) may necessitate additional bolus doses to maintain adequate block for surgery.

Upon reaching the desired level of neuromuscular block, the infusion of Rocuronium Bromide Injection must be individualized for each patient. The rate of administration should be adjusted according to the patient's twitch response as monitored with the use of a peripheral nerve stimulator. In clinical trials, infusion rates have ranged from 4 to 16 mcg/kg/min.

Inhalation anesthetics, particularly enflurane and isoflurane, may enhance the neuromuscular blocking action of nondepolarizing muscle relaxants. In the presence of steady-state concentrations of enflurane or isoflurane, it may be necessary to reduce the rate of infusion by 30% to 50%, at 45 to 60 minutes after the intubating dose.

Spontaneous recovery and reversal of neuromuscular blockade following discontinuation of Rocuronium Bromide Injection infusion may be expected to proceed at rates comparable to that following comparable total doses administered by repetitive bolus injections [see *Clinical Pharmacology (12.2)*].

Infusion solutions of Rocuronium Bromide Injection can be prepared by mixing Rocuronium Bromide Injection with an appropriate infusion solution such as 5% glucose in water or lactated Ringers [see *Dosage and Administration (2.7)*]. These infusion solutions should be used within 24 hours of mixing. Unused portions of infusion solutions should be discarded.

Infusion rates of Rocuronium Bromide Injection can be individualized for each patient using the following tables for 3 different concentrations of rocuronium bromide solution as guidelines:

TABLE 1: Infusion Rates Using Rocuronium Bromide Injection (0.5 mg/mL)*

Patient Weight	Drug Delivery Rate (mcg/kg/min)															
	4	5	6	7	8	9	10	12	14	16	18	20	24	28		
(kg)	(lbs)	Infusion Delivery Rate (mL/hr)														
10	22	4.8	6	7.2	8.4	9.6	10.8	12	14.4	16.8	19.2	21.6	24	28.8	33.6	38.4
15	33	7.2	9	10.8	12.6	14.4	16.2	18	21.6	25.2	28.8	32.4	36	43.2	50.4	57.6
20	44	9.6	12	14.4	16.8	19.2	21.6	24	28.8	33.6	38.4	43.2	48	57.6	67.2	76.8
25	55	12	15	18	21	24	27	30	36	42	48	54	60	72	84	96
35	77	16.8	21	25.2	29.4	33.6	37.8	42	50.4	58.8	67.2	75.6	84	100.8	115.2	130
50	110	24	30	36	42	48	54	60	72	84	96	108	120	144	168	192
60	132	28.8	36	43.2	50.4	57.6	64.8	72	86.4	100.8	115.2	130	144	172.8	198	226
70	154	33.6	42	50.4	58.8	67.2	75.6	84	100.8	117.6	134.4	151.2	168	201.6	230.4	260
80	176	38.4	48	57.6	67.2	76.8	86.4	96	115.2	134.4	153.6	172.8	192	225.6	259.2	293
90	198	43.2	54	64.8	75.6	86.4	97.2	108	129.6	151.2	172.8	194.4	216	252	288	324
100	220	48	60	72	84	96	108	120	144	168	192	216	240	288	336	384

* 50 mg Rocuronium Bromide Injection in 100 mL solution.

TABLE 2: Infusion Rates Using Rocuronium Bromide Injection (1 mg/mL)*

Patient Weight	Drug Delivery Rate (mcg/kg/min)															
	4	5	6	7	8	9	10	12	14	16	18	20	24	28		
(kg)	(lbs)	Infusion Delivery Rate (mL/hr)														
10	22	2.4	3	3.6	4.2	4.8	5.4	6	7.2	8.4	9.6	10.8	12	14.4	16.8	19.2
15	33	3.6	4.5	5.4	6.3	7.2	8.1	9	10.8	12.6	14.4	16.2	18	21.6	25.2	28.8
20	44	4.8	6	7.2	8.4	9.6	10.8	12	14.4	16.8	19.2	21.6	24	28.8	33.6	38.4
25	55	6	7.5	9	10.5	12	13.5	15	18	21	24	27	30	36	42	48
35	77	8.4	10.5	12.6	14.7	16.8	18.9	21	25.2	29.4	33.6	37.8	42	50.4	57.6	64.8
50	110	12	15	18	21	24	27	30	36	42	48	54	60	72	84	96
60	132	14.4	18	21.6	25.2	28.8	32.4	36	43.2	50.4	57.6	64.8	72	86.4	100.8	115.2
70	154	16.8	21	25.2	29.4	33.6	37.8	42	50.4	58.8	67.2	75.6	84	100.8	115.2	130
80	176	19.2	24	28.8	33.6	38.4	43.2	48	57.6	67.2	76.8	86.4	96	115.2	134.4	153.6
90	198	21.6	27	32.4	37.8	43.2	48.6	54	64.8	75.6	86.4	97.2	108	129.6	151.2	172.8
100	220	24	30	36	42	48	54	60	72	84	96	108	120	144	168	192

* 100 mg Rocuronium Bromide Injection in 100 mL solution.

TABLE 3: Infusion Rates Using Rocuronium Bromide Injection (5 mg/mL)*

Patient Weight	Drug Delivery Rate (mcg/kg/min)															
	4	5	6	7	8	9	10	12	14	16	18	20	24	28		
(kg)	(lbs)	Infusion Delivery Rate (mL/hr)														
10	22	0.5	0.6	0.7	0.8	1	1.1	1.2	1.4	1.7	1.9	2.1	2.4	2.8	3.3	3.8
15	33	0.7	0.9	1.1	1.3	1.4	1.6	1.8	2.2	2.5	2.9	3.2	3.6	4.3	5.0	5.7
20	44	1	1.2	1.4	1.7	1.9	2.2	2.4	2.9	3.4	3.8	4.2	4.8	5.7	6.7	7.6
25	55	1.2	1.5	1.8	2.1	2.4	2.7	3	3.6	4.2	4.8	5.4	6.2	7.4	8.6	9.8
35	77	1.7	2.1	2.5	2.9	3.4	3.8	4.2	5	5.9	6.7	7.6	8.7	10.4	12.1	13.8
50	110	2.4	3	3.6	4.2	4.8	5.4	6	7.2	8.4	9.6	10.8	12	14.4	16.8	19.2
60	132	2.9	3.6	4.3	5	5.8	6.5	7.2	8.6	10.1	11.5	13	14.4	17.2	20	22.8
70	154	3.4	4.2	5	5.9	6.7	7.6	8.4	10.1	11.8	13.4	15.1	17.3	20.4	23.6	27
80	176	3.8	4.8	5.8	6.7	7.7	8.6	9.6	11.5	13.4	15.4	17.3	19.2	22.8	26.4	30
90	198	4.3	5.4	6.5	7.6	8.6	9.7	10.8	13	15.1	17.3	19.2	21.6	25.2	29.4	34.2
100	220	4.8	6	7.2	8.4	9.6	10.8	12	14.4	16.8	19.2	21.6	24	28.8	33.6	38.4

* 500 mg Rocuronium Bromide Injection in 100 mL solution.

2.6 Dosage in Specific Populations

Pediatric Patients:

The recommended initial intubation dose of Rocuronium Bromide Injection is 0.6 mg/kg; however, a lower dose of 0.45 mg/kg may be used depending on anesthetic technique and the age of the patient.

For sevoflurane (induction) Rocuronium Bromide Injection doses of 0.45 mg/kg and 0.6 mg/kg in general produce excellent to good intubating conditions within 75 seconds. When halothane is used, a 0.6 mg/kg dose of Rocuronium Bromide Injection resulted in excellent to good intubating conditions within 60 seconds.

The time to maximum block for an intubating dose was shortest in infants (28 days up to 3 months) and longest in neonates (birth to less than 28 days). The duration of clinical relaxation following an intubating dose is shortest in children (greater than 2 years up to 11 years) and longest in infants.

When sevoflurane is used for induction and isoflurane/nitrous oxide for maintenance of general anesthesia, maintenance dosing of Rocuronium Bromide Injection can be administered as bolus doses of 0.15 mg/kg at reappearance of T₃ in all pediatric age groups. Maintenance dosing can also be administered at the reappearance of T₂ at a rate of 7 to 10 mcg/kg/min, with the lowest dose requirement for neonates (birth to less than 28 days) and the highest dose requirement for children (greater than 2 years up to 11 years).

When halothane is used for general anesthesia, patients ranging from 3 months old through adolescence can be administered

Rocuronium Bromide Injection maintenance doses of 0.075 to 0.125 mg/kg upon return of T₁ to 0.25% to provide clinical relaxation for 7 to 10 minutes. Alternatively, a continuous infusion of Rocuronium Bromide Injection initiated at a rate of 12 mcg/kg/min upon return of T₁ to 10% (one twitch present in train-of-four) may also be used to maintain neuromuscular blockade

