

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use SUCCINYLCHOLINE CHLORIDE INJECTION safely and effectively. See full prescribing information for SUCCINYLCHOLINE CHLORIDE INJECTION.

**Succinylcholine Chloride Injection, USP for intravenous or intramuscular use**  
Initial U.S. Approval: 1952

<b>WARNING: VENTRICULAR DYSRHYTHMIAS, CARDIAC ARREST, AND DEATH FROM HYPERKALEMIC RHABDOMYOLYSIS IN PEDIATRIC PATIENTS</b> <i>See full prescribing information for complete boxed warning.</i>
<b>• Acute rhabdomyolysis with hyperkalemia followed by ventricular dysrhythmias, cardiac arrest, and death has occurred after use in apparently healthy pediatric patients who were subsequently found to have undiagnosed skeletal muscle myopathy. (5.1)</b>
<b>• When a healthy-appearing pediatric patient develops cardiac arrest soon after administration of succinylcholine chloride, not felt to be due to other causes, immediate treatment for hyperkalemia should be instituted. In the presence of signs of malignant hyperthermia, appropriate treatment should be instituted concurrently. (5.1)</b>
<b>• Reserve use of succinylcholine chloride in pediatric patients for emergency intubation or instances where immediate securing of the airway is necessary, or for intramuscular use when a suitable vein is inaccessible. (5.1)</b>

## RECENT MAJOR CHANGES

Contraindications (4)	11/2022
Warnings and Precautions, Malignant Hyperthermia (5.5)	11/2022

## INDICATIONS AND USAGE

Succinylcholine chloride is a depolarizing neuromuscular blocker indicated in adults and pediatric patients:

- as an adjunct to general anesthesia (1)
- to facilitate tracheal intubation (1)
- to provide skeletal muscle relaxation during surgery or mechanical ventilation. (1)

## DOSSAGE AND ADMINISTRATION

- For intravenous or intramuscular use only. (2.1)
- Individualize dosage after careful assessment of the patient. (2.1)
- Accidental administration of neuromuscular blocking agents may be fatal. Store succinylcholine chloride with the cap and ferrule intact and in a manner that minimizes the possibility of selecting the wrong product. (2.1)
- See full prescribing information for succinylcholine chloride dosage recommendations, preparation instructions, and administration information. (2.2, 2.3, 2.4, 2.5, 2.6)

<b>FULL PRESCRIBING INFORMATION: CONTENTS*</b>
<b>WARNING: VENTRICULAR DYSRHYTHMIAS, CARDIAC ARREST, AND DEATH FROM HYPERKALEMIC RHABDOMYOLYSIS IN PEDIATRIC PATIENTS</b>
<b>1 INDICATIONS AND USAGE</b>
<b>2 DOSAGE AND ADMINISTRATION</b>
2.1 Important Dosage and Administration Information
2.2 Dosage Recommendations for Intravenous Use in Adults
2.3 Dosage Recommendations for Intravenous Use in Pediatric and Pediatric Patients
2.4 Dosage Recommendations for Intramuscular Use in Adults and Pediatric Patients
2.5 Preparation of Succinylcholine Chloride Injection
2.6 Drug Incompatibility
<b>3 DOSAGE FORMS AND STRENGTHS</b>
<b>4 CONTRAINDICATIONS</b>
<b>5 WARNINGS AND PRECAUTIONS</b>
5.1 Ventricular Dysrhythmias, Cardiac Arrest, and Death From Hyperkalemic Rhabdomyolysis in Pediatric Patients
5.2 Anaphylaxis
5.3 Risk of Death due to Medication Errors
5.4 Hyperkalemia
5.5 Malignant Hyperthermia
5.6 Bradycardia
5.7 Increase in Intraocular Pressure
5.8 Prolonged Neuromuscular Block due to Phase II Block and Tachyphylaxis
5.9 Risk of Prolonged Neuromuscular Block in Patients with Reduced Plasma Cholinesterase Activity
<b>6 DRUG INTERACTIONS</b>
6.1 Drugs that May Enhance the Neuromuscular Blocking Action of Succinylcholine Chloride
<b>7 DRUG INTERACTIONS</b>
7.1 Drugs that May Affect the Neuromuscular Blocking Action of Succinylcholine Chloride
<b>8 USE IN SPECIFIC POPULATIONS</b>
8.1 Pregnancy
8.2 Lactation
8.4 Pediatric Use
8.5 Geriatric Use
<b>10 OVERDOSAGE</b>
<b>11 DESCRIPTION</b>
<b>12 CLINICAL PHARMACOLOGY</b>
12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
12.5 Pharmacogenomics
<b>13 NONCLINICAL TOXICOLOGY</b>
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
<b>16 HOW SUPPLIED/STORAGE AND HANDLING</b>
* Sections or subsections omitted from the full prescribing information are not listed.

<b>FULL PRESCRIBING INFORMATION</b>
<b>WARNING: VENTRICULAR DYSRHYTHMIAS, CARDIAC ARREST, AND DEATH FROM HYPERKALEMIC RHABDOMYOLYSIS IN PEDIATRIC PATIENTS</b>
<b>• Acute rhabdomyolysis with hyperkalemia followed by ventricular dysrhythmias, cardiac arrest, and death has occurred after the administration of succinylcholine to apparently healthy pediatric patients who were subsequently found to have undiagnosed skeletal muscle myopathy, most frequently Duchenne muscular dystrophy [see Warnings and Precautions (5.1)].</b>
<b>• When a healthy appearing pediatric patient develops cardiac arrest within minutes after administration of succinylcholine chloride, not felt to be due to inadequate ventilation, oxygenation or anesthetic overdose, immediate treatment for hyperkalemia should be instituted. In the presence of signs of malignant hyperthermia, appropriate treatment should be instituted concurrently [see Warnings and Precautions (5.1)].</b>
<b>• Reserve the use of succinylcholine chloride in pediatric patients for emergency intubation or instances where immediate securing of the airway is necessary, e.g., laryngospasm, difficult airway, full stomach, or for intramuscular use when a suitable vein is inaccessible [see Warnings and Precautions (5.1)].</b>

<b>1 INDICATIONS AND USAGE</b>
Succinylcholine chloride is indicated in adults and pediatric patients: <ul style="list-style-type: none"><li>as an adjunct to general anesthesia</li> <li>to facilitate tracheal intubation</li> <li>to provide skeletal muscle relaxation during surgery or mechanical ventilation.</li></ul>
<b>2 DOSAGE AND ADMINISTRATION</b>
<b>2.1 Important Dosage and Administration Information</b>
<ul style="list-style-type: none"><li>Succinylcholine chloride is for intravenous or intramuscular use only.</li> <li>Succinylcholine chloride must be titrated to effect by or under supervision of experienced clinicians who are familiar with its actions and with appropriate neuromuscular monitoring techniques.</li> <li>Succinylcholine chloride should be administered only by those skilled in the management of artificial respiration and only when facilities are instantly available for tracheal intubation and for providing adequate ventilation of the patient, including the administration of oxygen under positive pressure and the elimination of CO<sub>2</sub>. The clinician must be prepared to assist or control respiration.</li> <li>The dosage of succinylcholine chloride should be individualized and should always be determined by the clinician after careful assessment of the patient.</li> <li>To avoid distress to the patient, do not administer succinylcholine chloride before unconsciousness has been induced [see Warnings and Precautions (5.14)].</li> <li>The occurrence of bradycardias with administration of succinylcholine chloride may be reduced by pretreatment with anticholinergics (e.g., atropine) [see Warnings and Precautions (5.6)].</li> <li>Monitor neuromuscular function with a peripheral nerve stimulator when using succinylcholine chloride by infusion [see Dosage and Administration (2.2), Warnings and Precautions (5.8)].</li> <li>Visually inspect succinylcholine chloride for particulate matter and discoloration prior to administration, hence, solution and container permit. Do not administer solutions that are not clear and colorless.</li> <li>Succinylcholine chloride supplied in multiple-dose vials does not require dilution before use [see Dosage and Administration (2.5)].</li></ul>
<b>Risk of Medication Errors</b>
Accidental administration of neuromuscular blocking agents may be fatal. Store succinylcholine chloride with the cap and ferrule intact and in a manner that minimizes the possibility of selecting the wrong product [see Warnings and Precautions (5.3)].
<b>2.2 Dosage Recommendations for Intravenous Use in Adults</b>
<b>For Short Surgical Procedures</b>
The average dose required to produce neuromuscular blockade and to facilitate tracheal intubation is 0.6 mg/kg succinylcholine chloride given intravenously. The optimum intravenous dose of succinylcholine chloride will vary among patients and may be from 0.3 mg/kg to 1.1 mg/kg for adults. Following intravenous administration of doses in this range, neuromuscular blockade develops in about 1 minute; maximum blockade may persist for about 2 minutes, after which recovery takes place within 4 to 6 minutes. A 5 to 10 mg intravenous test dose of succinylcholine chloride may be used to determine the sensitivity of the patient and the individual recovery time [see Warnings and Precautions (5.9)].
<b>For Long Surgical Procedures</b>
<b>Continuous Intravenous Infusion</b>
The dosage of succinylcholine chloride administered by continuous intravenous infusion depends upon the duration of the surgical procedure and the need for muscle relaxation.
Diluted succinylcholine chloride solutions containing from 1 mg/mL to 2 mg/mL succinylcholine have commonly been used for continuous intravenous infusion [see Dosage and Administration (2.5)]. The more dilute solution (1 mg/mL) is probably preferable from the standpoint of ease of control of the rate of administration of succinylcholine chloride and, hence, of relaxation. This diluted succinylcholine chloride solution containing 1 mg/mL succinylcholine may be administered intravenously at a rate of 0.5 mg (0.5 mL) per minute to 10 mg (10 mL) per minute to obtain the required amount of relaxation. The amount required per minute will depend upon the individual response as well as the degree of relaxation required. The average rate of continuous intravenous infusion for an adult ranges between 2.5 mg per minute and 4.3 mg per minute.
Monitor neuromuscular function with a peripheral nerve stimulator when using succinylcholine chloride by infusion in order to avoid overdose, detect development of Phase II block, follow its rate of recovery, and assess the effects of reversing agents [see Warnings and Precautions (5.8)].
<b>Intermittent Intravenous Injection</b>
Intermittent intravenous injections of succinylcholine chloride may also be used to provide muscle relaxation for long procedures. An intravenous injection of 0.3 mg/kg to 1.1 mg/kg may be given initially, followed, at appropriate intervals, by further intravenous injections of 0.04 mg/kg to 0.07 mg/kg to maintain the degree of relaxation required.
<b>2.3 Dosage Recommendations for Intravenous Use in Pediatric Patients</b>
For emergency tracheal intubation or in instances where immediate securing of the airway is necessary, the intravenous dose of succinylcholine chloride is 2 mg/kg for infants and other small pediatric patients; for older pediatric patients and adolescents the intravenous dose is 1 mg/kg [see Warnings and Precautions (5.1)]. Use in Specific Populations (8.4)]. The effective dose of succinylcholine chloride in pediatric patients may be higher than that predicted by body weight dosing alone. For example, the usual adult intravenous dose of 0.6 mg/kg is comparable to a dose of 2 mg/kg to 3 mg/kg in neonates and infants up to 6 months of age and 1 mg/kg to 2 mg/kg in infants up to 2 years of age [see Clinical Pharmacology (12.3)].
<b>2.4 Dosage Recommendations for Intramuscular Use in Adults and Pediatric Patients</b>
If a suitable vein is inaccessible, Succinylcholine Chloride Injection may be administered intramuscularly at a dose of up to 3 mg/kg to 4 mg/kg to infants, older pediatric patients, or adults. The total dose administered by the intramuscular route should not exceed 150 mg. The onset of effect of succinylcholine given intramuscularly is usually observed in about 2 to 3 minutes.
<b>2.5 Preparation of Succinylcholine Chloride Injection</b>
Succinylcholine Chloride Injection supplied in multiple-dose vials does not require dilution before use. Succinylcholine Chloride Injection may be diluted to 1 mg/mL or 2 mg/mL in a solution such as: <ul style="list-style-type: none"><li>5% Dextrose Injection, USP or</li> <li>0.9% Sodium Chloride Injection, USP</li></ul>
Prepare the diluted succinylcholine chloride solution for single patient use only. Store the diluted succinylcholine chloride solution in a refrigerator [2°C to 8°C (36°F to 46°F)] and use within 24 hours after preparation. Visually inspect the diluted succinylcholine chloride solution for particulate matter and discoloration prior to administration. Do not administer solutions that are not clear and colorless. Discard any unused portion of the diluted succinylcholine chloride solution.

<b>DOSSAGE FORMS AND STRENGTHS</b>
Injection: <ul style="list-style-type: none"><li>200 mg/10 mL (20 mg/mL) in multiple-dose flitop vials (3)</li></ul>
<b>CONTRAINDICATIONS</b>
<ul style="list-style-type: none"><li>Skeletal muscle myopathies (4)</li> <li>Known hypersensitivity to succinylcholine (4)</li> <li>After the acute phase of injury following major burns, multiple trauma, extensive denervation of skeletal muscle, or upper motor neuron injury (4)</li> <li>Known or suspected genetic susceptibility to malignant hyperthermia (4)</li></ul>
<b>WARNINGS AND PRECAUTIONS</b>
<ul style="list-style-type: none"><li><b>Anaphylaxis:</b> Severe anaphylactic reactions to neuromuscular blocking agents, including succinylcholine, have been reported. Some cases have been life-threatening and fatal. Take necessary precautions, such as the immediate availability of appropriate emergency treatment. (5.2)</li> <li><b>Risk of Death due to Medication Errors:</b> Unintended administration of succinylcholine chloride may result in paralysis, respiratory arrest and death. Confirm proper selection of intended product and avoid confusion with other injectable solutions that are present in critical care and other clinical settings. (5.3)</li> <li><b>Hyperkalemia:</b> Succinylcholine chloride may induce serious cardiac arrhythmias or cardiac arrest due to hyperkalemia. (5.4)</li> <li><b>Malignant Hyperthermia:</b> Malignant hyperthermia may occur, especially in individuals with known or suspected susceptibility based on genetic factors or family history. Discontinue triggering agents, administer intravenous dantrolene sodium, and apply supportive therapies. (5.5)</li> <li><b>Bradycardia:</b> Intravenous bolus administration may result in profound bradycardia or, rarely, asystole. The incidence is higher following a second dose of succinylcholine. Pretreatment with anticholinergic agents (e.g., atropine) may reduce the occurrence of bradycardias. (5.6)</li></ul>

## ADVERSE REACTIONS

Adverse reactions reported with succinylcholine are cardiac arrest, malignant hyperthermia, arrhythmias, bradycardia, tachycardia, hypertension, hypotension, hyperkalemia, prolonged respiratory depression or apnea. (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

<b>DRUG INTERACTIONS</b>
Drugs that May Enhance the Neuromuscular Blocking Action of Succinylcholine: promazine, oxytocin, aprotinin, certain non-penicillin antibiotics, quinidine, β-adrenergic blockers, procainamide, lidocaine, trimethaphan, lithium carbonate, magnesium salts, quinine, chloroquine, isoflurane, desflurane, metoclopramide, terbutaline, and drugs that reduce plasma cholinesterase activity. (7.1)
<b>Revised: 11/2023</b>

5.10 Risk of Additional Trauma in Patients With Fractures or Muscle Spasms
5.11 Increase in Intraocular Pressure
5.12 Risk of Aspiration due to Increase in Intra gastric Pressure
5.13 Prolonged Neuromuscular Block in Patients with Hypokalemia or Hypocalcemia
5.14 Risks due to Inadequate Anesthesia
<b>6 ADVERSE REACTIONS</b>
<b>7 DRUG INTERACTIONS</b>
7.1 Drugs that May Affect the Neuromuscular Blocking Action of Succinylcholine Chloride
<b>8 USE IN SPECIFIC POPULATIONS</b>
8.1 Pregnancy
8.2 Lactation
8.4 Pediatric Use
8.5 Geriatric Use
<b>10 OVERDOSAGE</b>
<b>11 DESCRIPTION</b>
<b>12 CLINICAL PHARMACOLOGY</b>
12.1 Mechanism of Action
12.2 Pharmacodynamics
12.3 Pharmacokinetics
12.5 Pharmacogenomics
<b>13 NONCLINICAL TOXICOLOGY</b>
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
<b>16 HOW SUPPLIED/STORAGE AND HANDLING</b>
* Sections or subsections omitted from the full prescribing information are not listed.

<b>2.6 Drug Incompatibility</b>
Succinylcholine Chloride Injection is acidic (pH is between 3.0 and 4.5) and may not be compatible with alkaline solutions having a pH greater than 8.5 (e.g., barbiturate solutions). Therefore, do not mix Succinylcholine Chloride Injection with alkaline solutions.
<b>3 DOSAGE FORMS AND STRENGTHS</b>
Succinylcholine Chloride Injection, USP is supplied as a clear, colorless solution as follows: 200 mg/10 mL (20 mg/mL) in multiple-dose flitop vials contains: 20 mg of succinylcholine chloride anhydrous (equivalent to 22 mg of succinylcholine chloride dihydrate).
<b>4 CONTRAINDICATIONS</b>
Succinylcholine chloride is contraindicated: <ul style="list-style-type: none"><li>in patients with skeletal muscle myopathies [see Warnings and Precautions (5.1)]</li> <li>in patients with known hypersensitivity to succinylcholine. Severe anaphylactic reactions to succinylcholine have been reported [see Warnings and Precautions (5.2)]</li> <li>after the acute phase of injury following major burns, multiple trauma, extensive denervation of skeletal muscle, or upper motor neuron injury, which may result in severe hyperkalemia and cardiac arrest [see Warnings and Precautions (5.4)]</li> <li>in patients with known or suspected genetic susceptibility to malignant hyperthermia [see Warnings and Precautions (5.5), Clinical Pharmacology (12.5)]</li></ul>
<b>5 WARNINGS AND PRECAUTIONS</b>
<b>5.1 Ventricular Dysrhythmias, Cardiac Arrest, and Death From Hyperkalemic Rhabdomyolysis in Pediatric Patients</b>
There have been reports of ventricular dysrhythmias, cardiac arrest, and death secondary to acute rhabdomyolysis with hyperkalemia in apparently healthy pediatric patients who received succinylcholine. Many of these pediatric patients were subsequently found to have a skeletal muscle myopathy such as Duchenne muscular dystrophy whose clinical signs were not obvious. The syndrome often presented as sudden cardiac arrest within minutes after the administration of succinylcholine. These pediatric patients were usually, but not exclusively, males, and most frequently 8 years of age or younger. There have also been reports in adolescents. There may be no signs or symptoms to alert the practitioner to which patients are at risk. A careful history and physical may identify developmental delays suggestive of a myopathy. A preoperative creatine kinase could identify some but not all patients at risk. When a healthy-appearing pediatric patient develops cardiac arrest within minutes after administration of succinylcholine chloride, not felt to be due to inadequate ventilation, oxygenation or anesthetic overdose, immediate treatment for hyperkalemia should be instituted. Due to the abrupt onset of this syndrome, routine resuscitative measures are likely to be unsuccessful. Careful monitoring of the electrocardiogram may alert the practitioner to peaked T-waves (an early sign). Administration of intravenous calcium, bicarbonate, and glucose with insulin, with hyperventilation have resulted in successful resuscitation in some of the reported cases. Extraordinary and prolonged resuscitative efforts have been effective in some cases. In addition, in the presence of signs of malignant hyperthermia, appropriate treatment should be initiated concurrently [see Warnings and Precautions (5.5)].
Because it is difficult to identify which patients are at risk, reserve the use of succinylcholine chloride in pediatric patients for emergency intubation or instances where immediate securing of the airway is necessary, e.g., laryngospasm, difficult airway, full stomach, or for intramuscular use when a suitable vein is inaccessible.
<b>5.2 Anaphylaxis</b>
Severe anaphylactic reactions to neuromuscular blocking agents, including succinylcholine, have been reported. These reactions have, in some cases, been life-threatening and fatal. Due to the potential severity of these reactions, the necessary precautions, such as the immediate availability of appropriate emergency treatment, should be taken. Allergic cross-reactivity between neuromuscular blocking agents, both depolarizing and non-depolarizing, has been reported in this class of drugs. Therefore, assess patients for previous anaphylactic reactions to other neuromuscular blocking agents before administering succinylcholine chloride.
<b>5.3 Risk of Death due to Medication Errors</b>
Administration of succinylcholine chloride results in paralysis, which may lead to respiratory arrest and death; this progression may be more likely to occur in a patient for whom it is not intended. Confirm proper selection of intended product and avoid confusion with other injectable solutions that are present in critical care and other clinical settings. If another healthcare provider is administering the product, ensure that the intended dose is clearly labeled and communicated.
<b>5.4 Hyperkalemia</b>
Succinylcholine chloride may induce serious cardiac arrhythmias or cardiac arrest due to hyperkalemia in patients with electrolyte abnormalities and those who may have digitalis toxicity. Succinylcholine chloride is contraindicated after the acute phase of injury following major burns, multiple trauma, extensive denervation of skeletal muscle, or upper motor neuron injury [see Contraindications (4)]. The risk of hyperkalemia in these patients increases over time and usually peaks at 7 to 10 days after the injury. The risk is dependent on the extent and location of the injury. The precise time of onset and the duration of the risk period are undetermined.
Patients with chronic abdominal infection, subarachnoid hemorrhage, or conditions causing degeneration of central and peripheral nervous systems are at an increased risk of developing severe hyperkalemia after succinylcholine chloride administration. Consider avoiding use of succinylcholine chloride in these patients until the patient's baseline potassium levels are within the normal range prior to succinylcholine chloride administration.
<b>5.5 Malignant Hyperthermia</b>
In susceptible individuals, succinylcholine may trigger malignant hyperthermia, a skeletal muscle hypermetabolic state leading to high oxygen demand. Fatal outcomes of malignant hyperthermia have been reported. The risk of developing malignant hyperthermia increases with the concomitant administration of succinylcholine and volatile anesthetic agents. Succinylcholine chloride can induce malignant hyperthermia in patients with known or suspected susceptibility based on genetic factors or family history, including those with certain inherited ryanodine receptor (RYR1) or dihydropyridine receptor (CACNA1S) variants. [see Contraindications (4), Clinical Pharmacology (12.5)]. Signs consistent with malignant hyperthermia may include hyperthermia, hypoxia, hypercapnia, muscle rigidity (e.g., jaw muscle spasm), tachycardia (e.g., particularly that unresponsive to deepening anesthesia or analgesic medication administration), tachypnea, cyanosis, arrhythmias, hypovolemia, and hemodynamic instability. Skin mottling, coagulopathies, and renal failure may occur later in the course of the hypermetabolic process. Successful treatment of malignant hyperthermia depends on early recognition of the clinical signs. If malignant hyperthermia is suspected, discontinue all triggering agents (i.e., volatile anesthetic agents and succinylcholine), administer intravenous dantrolene sodium, and initiate supportive therapies. Consult prescribing information for intravenous dantrolene sodium for additional information on patient management. Supportive therapies include administration of supplemental oxygen and respiratory support based on clinical need, maintenance of hemodynamic stability and adequate urinary output, management of fluid and electrolyte balance, correction of acid base derangements, and institution of measures to control rising temperature.
<b>5.6 Bradycardia</b>
Intravenous bolus administration of succinylcholine chloride in pediatric patients (including infants) may result in profound bradycardia or, rarely, asystole. In both adult and pediatric patients the incidence of bradycardia, which may progress to asystole, is higher following a second dose of succinylcholine. The incidence and severity of bradycardia is higher in pediatric patients than adults. Whereas bradycardia is common in pediatric patients after an initial dose of 1.5 mg/kg, bradycardia is seen in adults only after repeated exposure. Pretreatment with anticholinergic agents (e.g., atropine) may reduce the occurrence of bradyarrhythmias.
<b>5.7 Increase in Intraocular Pressure</b>
Succinylcholine causes an increase in intraocular pressure. Avoid succinylcholine chloride in instances in which an increase in intraocular pressure is undesirable (e.g., narrow angle glaucoma, penetrating eye injury) unless the potential benefit of its use outweighs the potential risk.

## 5.8 Prolonged Neuromuscular Block due to Phase II Block and Tachyphylaxis

When succinylcholine chloride is given over a prolonged period of time, the characteristic depolarization block of the myoneural junction (Phase I block) may change to a block with characteristics superficially resembling a non-depolarizing block (Phase II block). Prolonged respiratory muscle paralysis or weakness may be observed in patients manifesting this transition to Phase II block. Tachyphylaxis occurs with repeated administration [see Clinical Pharmacology (12.2)].

When Phase II block is suspected in cases of prolonged neuromuscular blockade, positive diagnosis should be made by peripheral nerve stimulation, prior to administration of any anticholinesterase drug. Reversal of Phase II block is a medical decision which must be made upon the basis of the patient, clinical pharmacology, and the experience and judgment of the clinician. The presence of Phase II block is indicated by fade of responses to successive stimuli (preferably “train of four”). The use of an anticholinesterase drug such as neostigmine to reverse Phase II block should be accompanied by appropriate doses of an anticholinergic drug to prevent disturbances of cardiac rhythm. After adequate reversal of Phase II block with an anticholinesterase agent, the patient should be continually observed for at least 1 hour for signs of return of muscle relaxation. Reversal should not be attempted unless: (1) a peripheral nerve stimulator is used to determine the presence of Phase II block (since anticholinesterase agents will potentiate succinylcholine-induced Phase I block), and (2) spontaneous recovery of muscle twitch has been observed for at least 20 minutes and has reached a plateau with further repeated dosing slowly; this delay is to ensure complete hydrolysis of succinylcholine by plasma cholinesterase prior to administration of the anticholinesterase agent. Should the type of block be misdiagnosed, depolarization of the type initially induced by succinylcholine (i.e., Phase I block) will be prolonged by an anticholinesterase agent.

## 5.9 Risk of Prolonged Neuromuscular Block in Patients with Reduced Plasma Cholinesterase Activity

Succinylcholine chloride is not recommended in patients with known reduced plasma cholinesterase (pseudocholinesterase) activity due to the likelihood of prolonged neuromuscular block following administration of succinylcholine chloride in such patients.

Plasma cholinesterase activity may be diminished in the presence of genetic abnormalities of plasma cholinesterase (e.g., patients heterozygous or homozygous for atypical plasma cholinesterase gene), pregnancy, severe liver or kidney disease, malignant tumors, infections, burns, anemia, decompensated heart disease, peptic ulcer, or myxedema. Plasma cholinesterase activity may also be diminished by chronic administration of oral contraceptives, glucocorticoids, or certain monamine oxidase inhibitors and by irreversible inhibitors of plasma cholinesterase (e.g., organophosphate insecticides, echothiophate, and certain antiangiastic drugs) [see Drug Interactions (7.1)].

Patients homozygous for atypical plasma cholinesterase gene (1 in 2,500 patients) are extremely sensitive to the neuromuscular blocking effect of succinylcholine. If succinylcholine chloride is administered to a patient homozygous for atypical plasma cholinesterase, resulting apnea or prolonged muscle paralysis should be treated with controlled respiration.

## 5.10 Risk of Additional Trauma in Patients With Fractures or Muscle Spasms

Succinylcholine chloride should be employed with caution in patients with fractures or muscle spasm because the initial muscle fasciculations may cause additional trauma. Monitor neuromuscular transmission and the development of fasciculations throughout the use of neuromuscular blocking agents.

## 5.11 Increase in Intraocular Pressure

Succinylcholine chloride may cause a transient increase in intraocular pressure; however, adequate anesthetic induction prior to administration of succinylcholine chloride will minimize this effect.

## 5.12 Risk of Aspiration due to Increase in Intra gastric Pressure

Succinylcholine may increase intragastric pressure, which could result in regurgitation and possible aspiration of stomach contents. Evaluate patients at risk for aspiration and reurgitation. Monitor patients during induction of anesthesia and neuromuscular blockade for clinical signs of vomiting and/or aspiration.

## 5.13 Prolonged Neuromuscular Block in Patients with Hypokalemia or Hypocalcemia

Neuromuscular blockade may be prolonged in patients with hypokalemia (e.g., after severe vomiting, diarrhea, digitalisation and diuretic therapy) or hypocalcemia (e.g., after massive transfusions). Correct severe electrolyte disturbances when possible. In order to help preclude possible prolongation of neuromuscar block, monitor neuromuscular transmission throughout the use of succinylcholine chloride.

## 5.14 Risks due to Inadequate Anesthesia

Neuromuscular blockade in the conscious patient can lead to distress. Use succinylcholine chloride in the presence of appropriate sedation or general anesthesia. Monitor patients to ensure that the level of anesthesia is adequate. In emergency situations, however, it may be necessary to administer succinylcholine chloride before unconsciousness is induced.

<b>6 ADVERSE REACTIONS</b>
The following clinically significant adverse reactions are discussed in greater detail in other sections of the labeling: <ul style="list-style-type: none"><li>Ventricular Dysrhythmias, Cardiac Arrest, and Death from Hyperkalemic Rhabdomyolysis in Pediatric Patients [see Warnings and Precautions (5.1)]</li> <li>Anaphylaxis [see Warnings and Precautions (5.2)]</li> <li>Hyperkalemia [see Warnings and Precautions (5.4)]</li> <li>Malignant Hyperthermia [see Warnings and Precautions (5.5)]</li> <li>Bradycardia [see Warnings and Precautions (5.6)]</li> <li>Increase in Intraocular Pressure [see Warnings and Precautions (5.7)]</li> <li>Prolonged Neuromuscular Block due to Phase II Block and Tachyphylaxis [see Warnings and Precautions (5.8)]</li></ul>
The following adverse reactions associated with the use of succinylcholine were identified in clinical studies or postmarketing reports. Because some of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.
<b>Cardiovascular disorders:</b> Cardiac arrest, arrhythmias, bradycardia, tachycardia, hypertension, hypotension
<b>Electrolyte disorders:</b> Hypokalemia
<b>Eye disorders:</b> Increased intraocular pressure
<b>Gastrointestinal disorders:</b> Excessive salivation
<b>Immune system disorders:</b> Hypersensitivity reactions including anaphylaxis (in some cases life-threatening and fatal)
<b>Musculoskeletal disorders:</b> Malignant hyperthermia, rhabdomyolysis with possible myoglobinuric acute renal failure, muscle fasciculation, jaw rigidity, postoperative muscle pain
<b>Respiratory disorders:</b> Prolonged respiratory depression or apnea
<b>Skin disorders:</b> Rash

**7 DRUG INTERACTIONS**

**7.1 Drugs that May Affect the Neuromuscular Blocking Action of Succinylcholine Chloride**

Drugs that may enhance the neuromuscular blocking action of succinylcholine include: promazine, oxytocin, aprotinin, certain non-penicillin antibiotics, quinidine, β-adrenergic blockers, procainamide, lidocaine, trimethaphan, lithium carbonate, magnesium salts, quinine, chloroquine, isoflurane, desflurane, metoclopramide, and terbutaline.

The neuromuscular blocking effect of succinylcholine may be enhanced by drugs that reduce plasma cholinesterase activity (e.g., chronically administered oral contraceptives, glucocorticoids, or certain monamine oxidase inhibitors) or by drugs that irreversibly inhibit plasma cholinesterase [see Warnings and Precautions (5.9)].

If other neuromuscular blocking agents are to be used during the same procedure, consider the possibility of a synergistic or antagonistic effect.

<b>8 USE IN SPECIFIC POPULATIONS</b>
<b>8.1 Pregnancy</b>
<b>Risk Summary</b>
Available data from published literature from case reports and case series over decades of use with succinylcholine during pregnancy have not identified a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. Succinylcholine is used commonly during delivery by caesarean section to provide muscle relaxation. If succinylcholine is used during labor and delivery, there is a risk for prolonged apnea in some pregnant women (see <i>Clinical Considerations</i> ). Animal reproduction studies have not been conducted with succinylcholine chloride.
The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.
<b>Clinical Considerations</b>
<b>Maternal Adverse Reactions</b>
Plasma cholinesterase levels are decreased by approximately 24% during pregnancy and for several days postpartum which can prolong the effect of succinylcholine. Therefore, some pregnant patients may experience prolonged apnea.
<b>Fetal/Neonatal Adverse Reactions</b>
Apnea and flaccidity may occur in the newborn after repeated high doses to, or in the presence of atypical plasma cholinesterase in, the mother.
<b>Labor or Delivery</b>
Succinylcholine is commonly used to provide muscle relaxation during delivery by caesarean section. Succinylcholine is known to cross the placental barrier in an amount that is dependent on the concentration gradient between the maternal and fetal circulation.

<b>8.2 Lactation</b>
<b>Risk Summary</b>
There are no data on the presence of succinylcholine or its metabolite in either human or animal milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for succinylcholine chloride and any potential adverse effects on the breastfed infant from succinylcholine chloride or from the underlying maternal condition.
<b>8.4 Pediatric Use</b>
Safety and effectiveness of succinylcholine chloride have been established in pediatric patient age groups, neonate to adolescent. Because of a risk of ventricular dysrhythmias, cardiac arrest, and death from hyperkalemic rhabdomyolysis in pediatric patients, reserve the use of succinylcholine chloride in pediatric patients for emergency intubation or instances where immediate securing of the airway is necessary, e.g., laryngospasm, difficult airway, full stomach, or for intramuscular use when a suitable vein is inaccessible [see Warnings and Precautions (5.1)].
Intravenous bolus administration of succinylcholine chloride in pediatric patients (including infants) may result in profound bradycardia or, rarely, asystole. The incidence and severity of bradycardia is higher in pediatric patients than adults [see Warnings and Precautions (5.6)].
The effective dose of succinylcholine chloride in pediatric patients may be higher than that predicted by body weight dosing alone [see Dosage and Administration (2.3)].

<b>8.5 Geriatric Use</b>
Clinical studies of succinylcholine chloride did not include sufficient numbers of subjects aged 65 years and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.
In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.
<b>10 OVERDOSAGE</b>
Overdosage with succinylcholine chloride may result in neuromuscular block beyond the time needed for surgery and anesthesia. This may be manifested by skeletal muscle weakness, decreased respiratory reserve, low tidal volume, or apnea. The primary treatment is maintenance of a patent airway and respiratory support until recovery of normal respiration is assured. Depending on the dose and duration of succinylcholine chloride administration, the characteristic neuromuscular block (Phase I) may change to a block with characteristics superficially resembling a non-depolarizing block (Phase II) [see Warnings and Precautions (5.8)].
<b>11 DESCRIPTION</b>
Succinylcholine Chloride Injection, USP is a sterile, nonpyrogenic solution to be used as a short-acting, depolarizing neuromuscular blocker for intravenous or intramuscular use. Succinylcholine Chloride Injection contains succinylcholine chloride as the active pharmaceutical ingredient.

Succinylcholine Chloride, USP is chemically designated C<sub>14</sub>H<sub>24</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>4</sub> and its molecular weight is 361.31. The chemical name of succinylcholine chloride is ethanaminium, 2,2'-[-(1,4-dioxo-1,4-butanediyloxy)]bis[N,N-trimethyl-], dichloride. Succinylcholine chloride is a dicationic base consisting of the dichloride salt of the dicholine ester of succinic acid. It is a white, odorless, slightly bitter powder, very soluble in water. It has the following structural formula:

