


**Cetorelix Acetate for Injection, 0.25 mg
FOR SUBCUTANEOUS USE ONLY****DESCRIPTION**

Cetorelix acetate for injection is a synthetic decapeptide with gonadotropin-releasing hormone (GnRH) antagonistic activity. Cetorelix acetate is an analog of native GnRH with substitutions of amino acids at positions 1, 2, 3, 6, and 10. The molecular formula is Acetyl-D-3-(2'-naphtyl)-alanine-D-4-chlorophenylalanine-D-3-(3'-pyridyl)-alanine-L-serine-L-tyrosine-D-citruline-L-leucine-L-arginine-L-proline-D-alanine-amide, and the molecular weight is 1431.06, calculated as the anhydrous free base. The structural formula is as follows:

Cetorelix acetate

Cetorelix Acetate for Injection 0.25 mg is a sterile lyophilized powder intended for subcutaneous injection after reconstitution with Sterile Water for Injection, USP, that comes supplied in a 1.0 mL pre-filled syringe. Each vial of cetorelix acetate for injection 0.25 mg contains 0.26-0.27 mg cetorelix acetate, equivalent to 0.25 mg cetorelix, and 54.8 mg mannitol.

CLINICAL PHARMACOLOGY

GnRH induces the production and release of luteinizing hormone (LH) and follicle stimulating hormone (FSH) from the gonadotrophic cells of the anterior pituitary. Due to a positive estradiol (E2) feedback at midcycle, GnRH liberation is enhanced resulting in an LH-surge. This LH-surge induces the ovulation of the dominant follicle, resumption of oocyte meiosis and subsequently luteinization as indicated by rising progesterone levels.

Cetorelix acetate for injection competes with natural GnRH for binding to membrane receptors on pituitary cells and thus controls the release of LH and FSH in a dose-dependent manner. The onset of LH suppression is approximately one hour with the 3 mg dose and two hours with the 0.25 mg dose. This suppression is maintained by continuous treatment and there is a more pronounced effect on LH than on FSH. An initial release of endogenous gonadotropins has not been detected with cetorelix acetate for injection, which is consistent with an antagonist effect.

The effects of cetorelix acetate for injection on LH and FSH are reversible after discontinuation of treatment. In women, cetorelix acetate for injection delays the LH-surge, and consequently ovulation, in a dose-dependent fashion. FSH levels are not affected at the doses used during controlled ovarian stimulation. Following a single 3 mg dose of cetorelix acetate for injection, duration of action of at least 4 days has been established. A dose of cetorelix acetate for injection 0.25 mg every 24 hours has been shown to maintain the effect.

Pharmacokinetics

The pharmacokinetic parameters of single and multiple doses of cetorelix acetate for injection in adult healthy female subjects are summarized in Table 1.

Table 1: Pharmacokinetic parameters of Cetorelix Acetate for Injection following 3 mg single or 0.25 mg single and multiple (daily for 14 days) subcutaneous (sc) administration.

	Single dose 3 mg	Single dose 0.25 mg	Multiple dose 0.25 mg
No. of subjects	12	12	12
t _{max} * [h]	1.5 (0.5-2)	1.0 (0.5-1.5)	1.0 (0.5-2)
t _{1/2} * [h]	62.8 (38.2-108)	5.0 (2.4-48.8)	20.6 (4.1-179.3)
C _{max} [ng/ml]	28.5 (22.5-36.2)	4.97 (4.17-5.92)	6.42 (5.18-7.96)
AUC [ng·h/ml]	536 (451-636)	31.4 (23.4-42.0)	44.5 (36.7-54.2)
CL [†] [ml/min·kg]	1.28 [‡]		
Vz [‡] [l/kg]	1.16 [‡]		

t_{max} Time to reach observed maximum plasma concentration

t_{1/2} Elimination half-life

C_{max} Maximum plasma concentration; multiple dose C_{ss, max}

AUC Area under the curve; single dose AUC_{0-inf}; multiple dose AUC₀₋₁₄

CL Total plasma clearance

Vz Volume of distribution

Geometric mean (95% CI_n),

* median (min-max)

† arithmetic mean,

‡ Based on iv administration (n=6, separate study 0013)

Absorption

Cetorelix acetate for injection is rapidly absorbed following subcutaneous injection, maximal plasma concentrations being achieved approximately one to two hours after administration. The mean absolute bioavailability of cetorelix acetate for injection following subcutaneous administration to healthy female subjects is 85%.

Distribution

The volume of distribution of cetorelix acetate for injection following a single intravenous dose of 3 mg is about 1 l/kg. *In vitro* protein binding to human plasma is 86%.

Cetorelix acetate for injection concentrations in follicular fluid and plasma were similar on the day of oocyte pick-up in patients undergoing controlled ovarian stimulation. Following subcutaneous administration of cetorelix acetate for injection 0.25 mg and 3 mg, plasma concentrations of cetorelix were below or in the range of the lower limit of quantitation on the day of oocyte pick-up and embryo transfer.

Metabolism

After subcutaneous administration of 10 mg cetorelix acetate for injection to females and males, cetorelix acetate for injection and small amounts of (1-9), (1-7), (1-6), and (1-4) peptides were found in bile samples over 24 hours.

In *in vitro* studies, cetorelix acetate for injection was stable against phase I- and phase II-metabolism. Cetorelix acetate for injection was transformed by peptidases, and the (1-4) peptide was the predominant metabolite.

Excretion

Following subcutaneous administration of 10 mg cetorelix to males and females, only unchanged cetorelix was detected in urine. In 24 hours, cetorelix and small amounts of the (1-9), (1-7), (1-6), and (1-4) peptides were found in bile samples. 2-4% of the dose was eliminated in the urine as unchanged cetorelix, while 5-10% was eliminated as cetorelix and the four metabolites in bile. Therefore, only 7-14% of the total dose was recovered as unchanged cetorelix and metabolites in urine and bile up to 24 hours. The remaining portion of the dose may not have been recovered since bile and urine were not collected for a longer period of time.

Special Populations

Pharmacokinetic investigations have not been performed either in subjects with impaired renal or liver function, or in the elderly, or in children (see PRECAUTIONS).

Pharmacokinetic differences in different races have not been determined.

There is no evidence of differences in pharmacokinetic parameters for cetorelix acetate for injection between healthy subjects and patients undergoing controlled ovarian stimulation.

Drug-Drug Interactions

No formal drug-drug interaction studies have been performed with cetorelix acetate for injection (see PRECAUTIONS).

Clinical Studies

Seven hundred thirty two (732) patients were treated with cetorelix acetate for injection in five (two Phase 2 dose-finding and three Phase 3) clinical trials. The clinical trial population consisted of Caucasians (95.5%) and Black, Asian, Arabian and others (4.5%). Women were between 19 and 40 years of age (mean: 32). The studies excluded subjects with polycystic ovary syndrome (PCOS), subjects with low or no ovarian reserve, and subjects with stage III-IV endometriosis.

Two dose regimens were investigated in these clinical trials, either a single dose per treatment cycle or multiple dosing. In the Phase 2 studies, a single dose of 3 mg was established as the minimal effective dose for the inhibition of premature LH surges with a protection period of at least 4 days. When cetorelix acetate for injection is administered in a multidose regimen, 0.25 mg was established as the minimal effective dose. The extent and duration of LH-suppression is dose dependent.

In the Phase 3 program, efficacy of the single 3 mg dose regimen of cetorelix acetate for injection and the multiple 0.25 mg dose regimen of cetorelix acetate for injection was established separately in two adequate and well controlled clinical studies utilizing active comparators. A third non-comparative clinical study evaluated only the multiple 0.25 mg dose regimen of cetorelix acetate for injection. The ovarian stimulation treatment with recombinant FSH or human menopausal gonadotropin (hMG) was initiated on day 2 or 3 of a normal menstrual cycle. The dose of gonadotropins was administered according to the individual patient's disposition and response.

In the single dose regimen study, cetorelix acetate for injection 3 mg was administered on the day of controlled ovarian stimulation when adequate estradiol levels (400 pg/mL) were obtained, usually on day 7 (range day 5-12). If hCG was not given within 4 days of the 3 mg dose of cetorelix acetate for injection, then 0.25 mg of cetorelix acetate for injection was administered daily beginning 96 hours after the 3 mg injection until and including the day of hCG administration.

In the two multiple dose regimen studies, cetorelix acetate for injection 0.25 mg was started on day 5 or 6 of COS. Both gonadotropins and cetorelix acetate for injection were continued daily (multiple dose regimen) until the injection of human chorionic gonadotropin (hCG).

Oocyte pick-up (OPU) followed by *in vitro* fertilization (IVF) or intracytoplasmic sperm injection (ICSI) as well as embryo transfer (ET) were subsequently performed. The results for cetorelix acetate for injection are summarized below in Table 2.

Table 2: Results of Phase 3 Clinical Studies with Cetorelix Acetate for Injection 3 mg in a single dose (sd) regimen and 0.25 mg in a multiple dose (md) regimen

Parameter	Cetorelix Acetate for Injection 3 mg (sd, active comparator study)	Cetorelix Acetate for Injection 0.25 mg (md, active comparator study)	Cetorelix Acetate for Injection 0.25 mg (md, non-comparative study)
No. of subjects	115	159	303
hCG administered [%]	98.3	96.2	96.0
Oocyte pick-up [%]	98.3	94.3	93.1
LH-surge [%] (LH ≥ 10 U/L and P** ≥ 1 ng/mL) [†]	0.0	1.9	1.0
Serum E ₂ [pg/ml] at day hCG ^{‡,§}	1125 (470-2952)	1064 (341-2531)	1185 (311-3676)
Serum LH [U/L] at day hCG ^{‡,§}	1.0 (0.5-2.5)	1.5 (0.5-7.6)	1.1 (0.5-3.5)
No. of follicles ≥11 mm at day hCG [†]	11.2±5.5	10.8±5.2	10.4±4.5
No. of oocytes: IVF [†] ICSI [†]	9.2±5.2 10.0±4.2	7.6±4.3 10.1±5.6	8.5±5.1 9.3±5.9
Fertilization rate: IVF [†] ICSI [†]	0.48±0.33 0.66±0.29	0.62±0.26 0.63±0.29	0.60±0.26 0.61±0.25
No. of embryos transferred [†]	2.6±0.9	2.1±0.6	2.7±1.0
Clinical pregnancy rate [%] per attempt per subject with ET	22.6 26.3	20.8 24.1	19.8 23.3

* Progesterone

† Following initiation of cetorelix acetate for injection therapy

‡ Morning values

§ Median with 5th – 95th percentiles

¶ Mean ± standard deviation

In addition to IVF and ICSI, one pregnancy was obtained after intrauterine insemination. In the five Phase 2 and Phase 3 clinical trials, 184 pregnancies have been reported out of a total of 732 patients (including 21 pregnancies following the replacement of frozen-thawed embryos).

In the 3 mg regimen, 9 patients received an additional dose of 0.25 mg of cetorelix acetate for injection and two other patients received two additional doses of 0.25 mg cetorelix acetate for injection. The median number of days of cetorelix acetate for injection multiple dose treatment was 5 (range 1-15) in both studies.

No drug related allergic reactions were reported from these clinical studies.

INDICATIONS AND USAGE

Cetorelix acetate for injection is indicated for the inhibition of premature LH surges in women undergoing controlled ovarian stimulation.

CONTRAINDICATIONS

Cetorelix acetate for injection is contraindicated under the following conditions:

1. Hypersensitivity to cetorelix acetate, extrinsic peptide hormones or mannitol.
2. Known hypersensitivity to GnRH or any other GnRH analogs.
3. Known or suspected pregnancy, and lactation (see PRECAUTIONS).
4. Severe renal impairment

WARNINGS

Cetorelix acetate for injection should be prescribed by physicians who are experienced in fertility treatment. Before starting treatment with cetorelix acetate for injection, pregnancy must be excluded (see CONTRAINDICATIONS and PRECAUTIONS).

PRECAUTIONS**General**

Cases of hypersensitivity reactions, including anaphylactoid reactions with the first dose, have been reported during post-marketing surveillance (see ADVERSE REACTIONS). A severe anaphylactic reaction associated with cough, rash, and hypotension, was observed in one patient after seven months of treatment with cetorelix acetate for injection (10 mg/day) in a study for an indication unrelated to infertility.

Special care should be taken in women with signs and symptoms of active allergic conditions or known history of allergic predisposition. Treatment with cetorelix acetate for injection is not advised in women with severe allergic conditions.

Information for Patients

Prior to therapy with cetorelix acetate for injection, patients should be informed of the duration of treatment and monitoring procedures that will be required. The risk of possible adverse reactions should be discussed (see ADVERSE REACTIONS). Cetorelix acetate for injection should not be prescribed if a patient is pregnant.

If cetorelix acetate for injection is prescribed to patients for self-administration, information for proper use is given in the Patient Leaflet (see below).

Laboratory Tests

After the exclusion of preexisting conditions, enzyme elevations (ALT, AST, GGT, alkaline phosphatase) were found in 1-2% of patients receiving cetorelix acetate for injection during controlled ovarian stimulation. The elevations ranged up to three times the upper limit of normal. The clinical significance of these findings was not determined.

During stimulation with human menopausal gonadotropin, cetorelix acetate for injection had no notable effects on hormone levels aside from inhibition of LH surges.

Drug Interactions

No formal drug interaction studies have been performed with cetorelix acetate for injection.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term carcinogenicity studies in animals have not been performed with cetorelix acetate. Cetorelix acetate was not genotoxic *in vitro* (Ames test, HPRT test, chromosome aberration test) or *in vivo* (chromosome aberration test, mouse micronucleus test). Cetorelix acetate induced polyploidy in CHL-Chinese hamster lung fibroblasts, but not in V79-Chinese hamster lung fibroblasts, cultured peripheral human lymphocytes or in an *in vitro* micronucleus test in the CHL-cell line. Treatment with 0.46 mg/kg cetorelix acetate for 4 weeks resulted in complete infertility in female rats which was reversed 8 weeks after cessation of treatment.

Pregnancy**(see CONTRAINDICATIONS)**

Cetorelix acetate for injection is contraindicated in pregnant women.

When administered to rats for the first seven days of pregnancy, cetorelix acetate did not affect the development of the implanted conceptus at doses up to 38 mcg/kg (approximately 1 times the recommended human therapeutic dose based on body surface area). However, a dose of 139 mcg/kg (approximately 4 times the human dose) resulted in a resorption rate and a postimplantation loss of 100%. When administered from day 6 to near term to pregnant rats and rabbits, very early resorptions and total implantation losses were seen in rats at doses from 4.6 mcg/kg (0.2 times the human dose) and in rabbits at doses from 6.8 mcg/kg (0.4 times the human dose). In animals that maintained their pregnancy, there was no increase in the incidence of fetal abnormalities.

The fetal resorption observed in animals is a logical consequence of the alteration in hormonal levels effected by the antagonistic properties of cetorelix acetate for injection, which could result in fetal loss in humans as well. Therefore, this drug should not be used in pregnant women.

Nursing Mothers

It is not known whether cetorelix acetate for injection is excreted in human milk. Because many drugs are excreted in human milk, and because the effects of cetorelix acetate for injection on lactation and/or the breast-fed child have not been determined, cetorelix acetate for injection should not be used by nursing mothers.

Geriatric Use

Cetorelix acetate for injection is not intended to be used in subjects aged 65 and over.

ADVERSE REACTIONS

The safety of cetorelix acetate for injection in 949 patients undergoing controlled ovarian stimulation in clinical studies was evaluated. Women were between 19 and 40 years of age (mean: 32). 94.0% of them were Caucasian. Cetorelix acetate for injection was given in doses ranging from 0.1 mg to 5 mg as either a single or multiple dose.

Table 3 shows systemic adverse events, reported in clinical studies without regard to causality, from the beginning of cetorelix acetate for injection treatment until confirmation of pregnancy by ultrasound at an incidence ≥ 1% in cetorelix acetate for injection treated subjects undergoing COS.

Table 3: Adverse Events in ≥1%

(WHO preferred term)	Cetorelix Acetate for Injection N=949 % (n)
Ovarian Hyperstimulation Syndrome*	3.5 (33)
Nausea	1.3 (12)
Headache	1.1 (10)

* Intensity moderate or severe, or WHO Grade II or III, respectively

Local site reactions (e.g. redness, erythema, bruising, itching, swelling, and pruritus) were reported. Usually, they were of a transient nature, mild intensity and short duration. During post-marketing surveillance, cases of mild to moderate Ovarian Hyperstimulation syndrome and cases of hypersensitivity reactions including anaphylactoid reactions have been reported.

Two stillbirths were reported in Phase 3 studies of cetorelix acetate for injection.

Congenital Anomalies

Clinical follow-up studies of 316 newborns of women administered cetorelix acetate for injection were reviewed. One infant of a set of twin neonates was found to have anencephaly at birth and died after four days. The other twin was normal. Developmental findings from ongoing baby follow-up included a child with a ventricular septal defect and another child with bilateral congenital glaucoma.

Four pregnancies that resulted in therapeutic abortion in Phase 2 and Phase 3 controlled ovarian stimulation studies had major anomalies (diaphragmatic hernia, trisomy 21, Klinefelter syndrome, polydactyly, and trisomy 18). In three of these four cases, intracytoplasmic sperm injection (ICSI) was the fertilization method employed; in the fourth case, *in vitro* fertilization (IVF) was the method employed.

The minor congenital anomalies reported include: supernumerary nipple, bilateral strabismus, imperforate hymen, congenital nevi, hemangiomas, and QT syndrome.

The causal relationship between the reported anomalies and cetorelix acetate for injection is unknown. Multiple factors, genetic and others (including, but not limited to ICSI, IVF, gonadotropins, and progesterone) make causal attribution difficult to study.

To report SUSPECTED ADVERSE REACTIONS contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

There have been no reports of overdosage with cetorelix acetate for injection 0.25 mg or 3 mg in humans. Single doses up to 120 mg cetorelix acetate for injection have been well tolerated in patients treated for other indications without signs of overdosage.

DOSAGE AND ADMINISTRATION

Ovarian stimulation therapy with gonadotropins (FSH, hMG) is started on cycle Day 2 or 3. The dose of gonadotropins should be adjusted according to individual response. Cetorelix acetate for injection 0.25 mg may be administered subcutaneously once daily during the early- to mid-follicular phase.

Cetorelix acetate for Injection 0.25 mg is administered on either stimulation day 5 (morning or evening) or day 6 (morning) and continued daily until the day of hCG administration.

When assessment by ultrasound shows a sufficient number of follicles of adequate size, hCG is administered to induce ovulation and final maturation of the oocytes. No hCG should be administered if the ovaries show an excessive response to the treatment with gonadotropins to reduce the chance of developing ovarian hyperstimulation syndrome (OHSS).

Administration

Cetorelix acetate for Injection 0.25 mg can be administered by the patient herself after appropriate instructions by her doctor.

Directions for using Cetorelix Acetate for Injection 0.25 mg with the enclosed needles and pre-filled syringe:

1. Wash hands thoroughly with soap and water.
2. Flip off the plastic cover of the vial and wipe the aluminum ring and the rubber stopper with an alcohol swab.
3. Twist the injection needle with the yellow mark (20 gauge) on the pre-filled syringe.
4. Push the needle through the center of the rubber stopper of the vial and slowly inject the solvent into the vial.
5. Leaving the syringe in the vial, gently swirl the vial until the solution is clear and without residues. Avoid forming bubbles.
6. Draw the total contents of the vial into the syringe. If necessary, invert the vial and pull back the needle as far as needed to withdraw the entire contents of the vial. Discard unused portion.
7. Replace the needle with the yellow mark by the injection needle with the grey mark (27 gauge).
8. Invert the syringe and push the plunger until all air bubbles have been expelled.
9. Choose an injection site in the lower abdominal area, preferably around, but staying at least one inch away from the navel. Choose a different injection site each day to minimize local irritation. Use a second alcohol swab to clean the skin at the injection site and allow alcohol to dry. Gently pinch up the skin surrounding the site of injection.
10. Inject the prescribed dose as directed by your doctor, nurse or pharmacist.
11. Use the syringe and needles only once. Dispose of the syringe and needles properly after use. If available, use a medical waste container for disposal.

HOW SUPPLIED

Cetorelix Acetate for Injection 0.25 mg is available in a carton of one packaged tray.

Presentation	Unit of Sale	Description
Cetorelix Acetate for Injection Kit	NDC 65219-292-77	Each packaged tray contains: One glass vial containing 0.26 - 0.27 mg cetorelix acetate (corresponding to 0.25 mg cetorelix) (NDC 65219-290-02). One pre-filled glass syringe with 1 mL of Sterile Water for Injection, USP (NDC 65219-286-01). One 20 gauge needle (yellow) and One 27 gauge needle (grey)

Storage

Store Cetorelix Acetate for Injection 0.25 mg refrigerated, 2-8°C (36-46°F). Store the packaged tray in the outer carton in order to protect from light.

Rx only

Manufactured for:

FRESENIUS KABI

Lake Zurich, IL 60047

www.fresenius-kabi.com/us

Made in India

4 5 1 8 0 1

Revised: 04/2024

Patient Leaflet Cetorelix Acetate for Injection 0.25 mg

Active ingredient: cetorelix acetate

Summary

Cetorelix Acetate for Injection blocks the effects of a natural hormone, called gonadotropin-releasing hormone (GnRH). GnRH controls the secretion of another hormone, called luteinizing hormone (LH), which induces ovulation during the menstrual cycle. During hormone treatment for ovarian stimulation, premature ovulation may lead to eggs that are not suitable for fertilization. Cetorelix Acetate for Injection blocks such undesirable premature ovulation.

Uses

Cetorelix Acetate for Injection is used to prevent premature ovulation during controlled ovarian stimulation.

General Cautions

Do not use Cetorelix Acetate for Injection if you

- have kidney disease
- are allergic to cetorelix acetate, mannitol or exogenous peptide hormones (medicines similar to cetorelix acetate for injection) or
- are pregnant, or think that you might be pregnant, or if you are breast-feeding.

Consult your doctor before taking Cetorelix Acetate for Injection if you have had severe allergic reactions.

Proper Use

Ovarian stimulation therapy is started on cycle Day 2 or 3. Cetorelix Acetate for Injection 0.25 mg is injected under the skin once daily, as directed by your physician. When an ultrasound examination shows that you are ready, another drug (hCG) is injected to induce ovulation.

How should you use Cetorelix Acetate for Injection?

You may self-inject cetorelix acetate for injection after special instruction from your doctor.

To fully benefit from cetorelix acetate for injection, please read carefully and follow the instructions given below, unless your doctor advises you otherwise.

Cetorelix acetate for injection is for injection under the skin of the lower abdominal area, preferably around, but staying at least one inch away from the belly button. Choose a different injection site each day to minimize local irritation.

Dissolve cetorelix acetate for injection powder only with the water contained in the pre-filled syringe. Do not use a cetorelix acetate for injection solution if it contains particles or if it is not clear.

Before you inject cetorelix acetate for injection yourself, please read the following instructions carefully:

Directions for using Cetorelix Acetate for Injection 0.25 mg with the enclosed needles and pre-filled syringe:


1. Wash your hands thoroughly with soap and water.




2. On a clean flat surface, lay out everything you need (one vial of powder, one pre-filled syringe, one injection needle with a yellow mark, and one injection needle with a grey mark).



3. Flip off the plastic cover of the vial. Wipe the aluminum ring and the rubber stopper with an alcohol swab.




4. Take the injection needle with the yellow mark and remove the wrapping. Take the pre-filled syringe and remove the cover. Twist the needle on the syringe and remove the cover of the needle.




5. Push the needle through the center of the rubber stopper of the vial. Inject the water into the vial by slowly pushing down on the plunger of the syringe.




6. Leave the syringe in the vial. While carefully holding the syringe and vial, swirl gently to mix the powder and water together. When it is mixed, it will look clear and have no particles in it. Do not shake or you will create bubbles in your medicine.



7. Draw the total contents of the vial into the syringe. If liquid is left in the vial, invert the vial, pull back the needle until the opening of the needle is just inside the stopper. If you look from the side through the gap in the stopper, you can control the movement of the needle and the liquid. It is important to withdraw the entire contents of the vial. Discard unused portion.



8. Detach the syringe from the needle and lay down the syringe. Take the injection needle with the grey mark and remove its wrapping. Twist the needle on the syringe and remove the cover of the needle.



9. Invert the syringe and push the plunger until all air bubbles have been pushed out. Do not touch the needle or allow the needle to touch any surface.



10. Choose an injection site in the lower abdominal area, preferably around, but at least one inch away from the belly button. Choose a different injection site each day to minimize local irritation. Take a second alcohol swab and clean the skin at the injection site and allow alcohol to dry. Inject the prescribed dose as directed by your doctor, nurse or pharmacist.



11. Use the syringe and needles only once. Dispose of the syringe and needles immediately after use (put the covers on the needles to avoid injury). A medical waste container should be used for disposal.



SPECIAL ADVICE

What do you do if you have used too much Cetorelix Acetate for Injection?

Contact your doctor in case of overdosage immediately to check whether an adjustment of the further ovarian stimulation procedure is required.

Possible Side Effects

Mild and short lasting reactions may occur at the injection site like reddening, itching, and swelling. Nausea and headache have also been reported.

Call your doctor if you have any side effect not mentioned in this leaflet or if you are unsure about the effect of this medicine.

Storage

How is Cetorelix Acetate for Injection to be stored?

Store Cetorelix Acetate for Injection in a cool dry place protected from excess moisture and heat.

Store Cetorelix Acetate for Injection 0.25 mg in the refrigerator at 2-8°C (36-46°F). Keep the packaged tray in the outer carton in order to protect it from light.

How long may Cetorelix Acetate for Injection be stored?

Do not use the Cetorelix Acetate for Injection powder or the pre-filled syringe after the expiration date, which is printed on the labels and on the carton, and dispose of the vial and the syringe properly.

How long can you keep Cetorelix Acetate for Injection after preparation of the solution?

The solution should be used immediately after preparation.

Store the medicine out of the reach of children.

If you suspect that you may have taken more than the prescribed dose of this medicine, contact your doctor immediately. This medicine was prescribed for your particular condition. Do not use it for another condition or give the drug to others.

This leaflet provides a summary of the information about Cetorelix Acetate for Injection. Medicines are sometimes prescribed for uses other than those listed in the Leaflet. If you have any questions or concerns, or want more information about Cetorelix Acetate for Injection, contact your doctor or pharmacist.

This Leaflet has been approved by the U.S. Food and Drug Administration.

Manufactured for:

FRESENIUS KABI

Lake Zurich, IL 60047

www.fresenius-kabi.com/us

Made in India

4 5 1 8 0 1

Revised: 04/2024