

451851 /Issued: June 2025

Rocuronium Bromide Injection

HIGHLIGHTS OF PRESCRIBING INFORMATION: These highlights do not include all the information needed to use **ROCURONIUM BROMIDE INJECTION** safely and effectively. See full prescribing information for **ROCURONIUM BROMIDE INJECTION**.

ROCURONIUM BROMIDE Injection, for intravenous use
Initial U.S. Approval: 1994

INDICATIONS AND USAGE

Rocuronium Bromide Injection is a nondepolarizing neuromuscular blocking agent indicated as an adjunct to general anesthesia to facilitate both rapid sequence and routine tracheal intubation, and to provide skeletal muscle relaxation during surgery or mechanical ventilation. (1)

DOSAGE AND ADMINISTRATION

Rocuronium Bromide Injection should only be administered by experienced clinicians or trained individuals supervised by an experienced clinician familiar with the use, actions, characteristics, and complications of neuromuscular blocking agents. (2.1)

- Individualize the dose for each patient. (2.1)
- Peripheral nerve stimulator recommended for determination of drug response and need for additional doses, and to evaluate recovery. (2.1)
- Store Rocuronium Bromide Injection with cap and ferrule intact and in a manner that minimizes the possibility of selecting the wrong product. (2.1)
- Tracheal intubation: Rocuronium Bromide Injection dose is 0.6 mg/kg. (2.2)
- Rapid sequence intubation: 0.6 to 1.2 mg/kg. (2.3)
- Maintenance doses: Guided by response to prior dose, not administered until recovery is evident. (2.4)
- Continuous infusion: Initial rate of 10 to 12 mcg/kg/min. Start only after early evidence of spontaneous recovery from an intubating dose. (2.5)

DOSAGE FORMS AND STRENGTHS

- 50 mg/5 mL (10 mg/mL). (3)
- 100 mg/10 mL (10 mg/mL). (3)

CONTRAINDICATIONS

Hypersensitivity (e.g., anaphylaxis) to rocuronium bromide or other neuromuscular blocking agents. (4)

FULL PRESCRIBING INFORMATION: CONTENTS*

| | |
|------|---|
| 1 | INDICATIONS AND USAGE |
| 2 | DOSAGE AND ADMINISTRATION |
| 2.1 | Important Dosing and Administration Information |
| 2.2 | Dose for Tracheal Intubation |
| 2.3 | Rapid Sequence Intubation |
| 2.4 | Maintenance Dosing |
| 2.5 | Use by Continuous Infusion |
| 2.6 | Dosing in Specific Populations |
| 2.7 | Preparation for Administration of Rocuronium Bromide Injection |
| 3 | DOSAGE FORMS AND STRENGTHS |
| 4 | CONTRAINDICATIONS |
| 5 | WARNINGS AND PRECAUTIONS |
| 5.1 | Appropriate Administration and Monitoring |
| 5.2 | Individualized Dosing |
| 5.3 | Risk of Death due to Medication Errors |
| 5.4 | Need for Adequate Anesthesia |
| 5.5 | Residual Paralysis |
| 5.6 | Long-Term Use in an Intensive Care Unit |
| 5.7 | Malignant Hyperthermia (MH) |
| 5.8 | Prolonged Circulation Time |
| 5.9 | QT Interval Prolongation |
| 5.10 | Conditions/Drugs Causing Potentiation of, or Resistance to, Neuromuscular Block |
| 5.11 | Incompatibility with Alkaline Solutions |
| 5.12 | Increase in Pulmonary Vascular Resistance |
| 5.13 | Use in Patients with Myasthenia |
| 5.14 | Extravasation |
| 6 | ADVERSE REACTIONS |
| 6.1 | Clinical Trials Experience |
| 6.2 | Postmarketing Experience |
| 7 | DRUG INTERACTIONS |
| 7.1 | Antibiotics |
| 7.2 | Anticonvulsants |
| 7.3 | Inhalation Anesthetics |

FULL PRESCRIBING INFORMATION

- 1 INDICATIONS AND USAGE
- Rocuronium Bromide Injection is indicated as an adjunct to general anesthesia to facilitate both rapid sequence and routine tracheal intubation, and to provide skeletal muscle relaxation during surgery or mechanical ventilation.
- 2 DOSAGE AND ADMINISTRATION
- 2.1 Important Dosing and Administration Information
- Rocuronium Bromide Injection is for intravenous use only.
- Rocuronium Bromide Injection should only be administered by experienced clinicians or trained individuals supervised by an experienced clinician familiar with the use, actions, characteristics, and complications of neuromuscular blocking agents. Doses of Rocuronium Bromide Injection should be individualized and a peripheral nerve stimulator should be used to monitor drug effect, need for additional doses, adequacy of spontaneous recovery or antagonism, and to provide early evidence of the complications of overdose if additional doses are administered.
- The dosage information which follows is derived from studies based upon units of drug per unit of body weight. It is intended to serve as an initial guide to clinicians familiar with other neuromuscular blocking agents to acquire experience with Rocuronium Bromide Injection.
- In patients in whom potentiation of, or resistance to, neuromuscular block is anticipated, a dose adjustment should be considered [see Dosage and Administration (2.6), Warnings and Precautions (5.10, 5.13), Drug Interactions (7.2, 7.3, 7.4, 7.5, 7.6, 7.8, 7.10), Use in Specific Populations (6.6)].
- Risk of Medication Errors
- Accidental administration of neuromuscular blocking agents may be fatal. Store Rocuronium Bromide Injection with the cap and ferrule intact and in a manner that minimizes the possibility of selecting the wrong product [see Warnings and Precautions (5.3)].
- 2.2 Dose for Tracheal Intubation
- The recommended initial dose of Rocuronium Bromide Injection, regardless of anesthetic technique, is 0.6 mg/kg. Neuromuscular block sufficient for intubation (80% block or greater) is attained in a median (range) time of 1 (0.4-6) minute(s) and most patients have intubation completed within 2 minutes. Maximum blockade is achieved in most patients in less than 3 minutes. This dose may be expected to provide 31 (15-85) minutes of clinical relaxation under opioid/nitrous oxide/oxygen anesthesia. Under halothane, isoflurane, and enflurane anesthesia, some extension of the period of clinical relaxation should be expected [see Drug Interactions (7.3)].
- A lower dose of Rocuronium Bromide Injection (0.45 mg/kg) may be used. Neuromuscular block sufficient for intubation (80% block or greater) is attained in a median (range) time of 1.3 (0.8-6.2) minute(s), and most patients have intubation completed within 2 minutes. Maximum blockade is achieved in most patients in less than 4 minutes. This dose may be expected to provide 22 (12-31) minutes of clinical relaxation under opioid/nitrous oxide/oxygen anesthesia. Patients receiving this low dose of 0.45 mg/kg who achieve less than 90% block (about 16% of these patients) may have a more rapid time to 25% recovery, 12 to 15 minutes.
- A large bolus dose of 0.9 or 1.2 mg/kg can be administered under opioid/nitrous oxide/oxygen anesthesia without adverse effects to the cardiovascular system [see Clinical Pharmacology (12.2)].
- 2.3 Rapid Sequence Intubation
- In appropriately premedicated and adequately anesthetized patients, Rocuronium Bromide Injection 0.6 to 1.2 mg/kg will provide excellent or good intubating conditions in most patients in less than 2 minutes [see Clinical Studies (14.1)].
- 2.4 Maintenance Dosing
- Maintenance doses of 0.1, 0.15, and 0.2 mg/kg Rocuronium Bromide Injection, administered at 25% recovery of control T₁ (defined as 3 twitches of train-of-four), provide a median (range) of 12 (2-31),

- 2.6 Dosage in Specific Populations
- Pediatric Patients
- The recommended initial intubation dose of Rocuronium Bromide Injection is 0.6 mg/kg; however, a lower dose of 0.45 mg/kg may be used depending on anesthetic technique and the age of the patient.
- For sevoflurane (induction) Rocuronium Bromide Injection doses of 0.45 mg/kg and 0.6 mg/kg in general produce excellent to good intubating conditions within 75 seconds. When halothane is used, a 0.6 mg/kg dose of Rocuronium Bromide Injection resulted in excellent to good intubating conditions within 60 seconds.
- The time to maximum block for an intubating dose was shortest in infants (28 days up to 3 months) and longest in neonates (birth to less than 28 days). The duration of clinical relaxation following an intubating dose is shortest in children (greater than 2 years up to 11 years) and longest in infants.
- When halothane is used for general anesthesia, patients ranging from 3 months old through adolescence can be administered Rocuronium Bromide Injection maintenance doses of 0.075 to 0.125 mg/kg upon return of T₁ to 0.25% to provide clinical relaxation for 7 to 10 minutes. Alternatively, a continuous infusion of Rocuronium Bromide Injection initiated at a rate of 12 mcg/kg/min upon return of T₁ to 10% (one twitch present in train-of-four) may also be used to maintain neuromuscular blockade in pediatric patients.
- When halothane is used for general anesthesia, patients ranging from 3 months old through adolescence can be administered Rocuronium Bromide Injection maintenance doses of 0.075 to 0.125 mg/kg upon return of T₁ to 0.25% to provide clinical relaxation for 7 to 10 minutes. Alternatively, a continuous infusion of Rocuronium Bromide Injection initiated at a rate of 12 mcg/kg/min upon return of T₁ to 10% (one twitch present in train-of-four) may also be used to maintain neuromuscular blockade in pediatric patients.
- Additional information for administration to pediatric patients of all age groups is presented elsewhere in the label [see Clinical Pharmacology (12.2)].
- The infusion of Rocuronium Bromide Injection must be individualized for each patient. The rate of administration should be adjusted according to the patient's twitch response as monitored with the use of a peripheral nerve stimulator. Spontaneous recovery and reversal of neuromuscular blockade following discontinuation of Rocuronium Bromide Injection must be expected to proceed at rates comparable to that following comparable total doses administered by repetitive bolus injections [see Clinical Pharmacology (12.2)].
- Inhalation anesthetics, particularly enflurane and isoflurane, may enhance the neuromuscular blocking action of nondepolarizing muscle relaxants. In the presence of steady-state concentrations of enflurane or isoflurane, it may be necessary to reduce the rate of infusion by 30% to 50%, at 45 to 60 minutes after the intubating dose.
- Spontaneous recovery and reversal of neuromuscular blockade following discontinuation of Rocuronium Bromide Injection infusion may be expected to proceed at rates comparable to that following comparable total doses administered by repetitive bolus injections [see Clinical Pharmacology (12.2)].
- Infusion solutions of Rocuronium Bromide Injection can be prepared by mixing Rocuronium Bromide Injection with an appropriate infusion solution such as 5% glucose in water or lactated Ringers [see Dosage and Administration (2.7)]. These infusion solutions should be used within 24 hours of mixing. Unused portions of infusion solutions should be discarded.
- Infusion rates of Rocuronium Bromide Injection can be individualized for each patient using the following tables for 3 different concentrations of Rocuronium Bromide Injection solution as guidelines:
- Table 1: Infusion Rates Using Rocuronium Bromide Injection (0.5 mg/ mL)*
- | Patient Weight (kg) (lbs) | Drug Delivery Rate (mcg/kg/min) | | | | | | | | | | | |
|---------------------------|---------------------------------|------|----|------|------|------|------|-----|-------|-------|-------|----|
| | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 12 | 14 | 16 | 18 | 20 |
| | Infusion Delivery Rate (mL/hr) | | | | | | | | | | | |
| 10 | 22 | 4.8 | 6 | 7.2 | 8.4 | 9.6 | 10.8 | 12 | 14.4 | 16.8 | 19.2 | |
| 15 | 33 | 7.2 | 9 | 10.8 | 12.6 | 14.4 | 16.2 | 18 | 21.6 | 25.2 | 28.8 | |
| 20 | 44 | 9.6 | 12 | 14.4 | 16.8 | 19.2 | 21.6 | 24 | 28.8 | 33.6 | 38.4 | |
| 25 | 55 | 12 | 15 | 18 | 21 | 24 | 27 | 30 | 36 | 42 | 48 | |
| 35 | 77 | 16.8 | 21 | 25.2 | 29.4 | 33.6 | 37.8 | 42 | 50.4 | 58.8 | 67.2 | |
| 50 | 110 | 24 | 30 | 36 | 42 | 48 | 54 | 60 | 72 | 84 | 96 | |
| 60 | 132 | 28.8 | 36 | 43.2 | 50.4 | 57.6 | 64.8 | 72 | 86.4 | 100.8 | 115.2 | |
| 70 | 154 | 33.6 | 42 | 50.4 | 58.8 | 67.2 | 75.6 | 84 | 100.8 | 117.6 | 134.4 | |
| 80 | 176 | 38.4 | 48 | 57.6 | 67.2 | 76.8 | 86.4 | 96 | 115.2 | 134.4 | 153.6 | |
| 90 | 198 | 43.2 | 54 | 64.8 | 75.6 | 86.4 | 97.2 | 108 | 129.6 | 151.2 | 172.8 | |
| 100 | 220 | 48 | 60 | 72 | 84 | 96 | 108 | 120 | 144 | 168 | 192 | |
- * 50 mg Rocuronium Bromide Injection in 100 mL solution.
- Table 2: Infusion Rates Using Rocuronium Bromide Injection (1 mg/ mL)*
- | Patient Weight (kg) (lbs) | Drug Delivery Rate (mcg/kg/min) | | | | | | | | | | | |
|---------------------------|---------------------------------|------|------|------|------|------|------|----|------|------|------|----|
| | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 12 | 14 | 16 | 18 | 20 |
| | Infusion Delivery Rate (mL/hr) | | | | | | | | | | | |
| 10 | 22 | 2.4 | 3 | 3.6 | 4.2 | 4.8 | 5.4 | 6 | 7.2 | 8.4 | 9.6 | |
| 15 | 33 | 3.6 | 4.5 | 5.4 | 6.3 | 7.2 | 8.1 | 9 | 10.8 | 12.6 | 14.4 | |
| 20 | 44 | 4.8 | 6 | 7.2 | 8.4 | 9.6 | 10.8 | 12 | 14.4 | 16.8 | 19.2 | |
| 25 | 55 | 6 | 7.5 | 9 | 10.5 | 12 | 13.5 | 15 | 18 | 21 | 24 | |
| 35 | 77 | 8.4 | 10.5 | 12.6 | 14.7 | 16.8 | 18.9 | 21 | 25.2 | 29.4 | 33.6 | |
| 50 | 110 | 12 | 15 | 18 | 21 | 24 | 27 | 30 | 36 | 42 | 48 | |
| 60 | 132 | 14.4 | 18 | 21.6 | 25.2 | 28.8 | 32.4 | 36 | 43.2 | 50.4 | 57.6 | |
| 70 | 154 | 16.8 | 21 | 25.2 | 29.4 | 33.6 | 37.8 | 42 | 50.4 | 58.8 | 67.2 | |
| 80 | 176 | 19.2 | 24 | 28.8 | 33.6 | 38.4 | 43.2 | 48 | 57.6 | 67.2 | 76.8 | |
| 90 | 198 | 21.6 | 27 | 32.4 | 37.8 | 43.2 | 48.6 | 54 | 64.8 | 75.6 | 86.4 | |
| 100 | 220 | 24 | 30 | 36 | 42 | 48 | 54 | 60 | 72 | 84 | 96 | |
- * 100 mg Rocuronium Bromide Injection in 100 mL solution.
- Table 3: Infusion Rates Using Rocuronium Bromide Injection (5 mg/ mL)*
- | Patient Weight (kg) (lbs) | Drug Delivery Rate (mcg/kg/min) | | | | | | | | | | | |
|---------------------------|---------------------------------|-----|-----|-----|-----|-----|------|------|------|------|------|----|
| | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 12 | 14 | 16 | 18 | 20 |
| | Infusion Delivery Rate (mL/hr) | | | | | | | | | | | |
| 10 | 22 | 0.5 | 0.6 | 0.7 | 0.8 | 1 | 1.1 | 1.2 | 1.4 | 1.7 | 1.9 | |
| 15 | 33 | 0.7 | 0.9 | 1.1 | 1.3 | 1.4 | 1.6 | 1.8 | 2.2 | 2.5 | 2.9 | |
| 20 | 44 | 1 | 1.2 | 1.4 | 1.7 | 1.9 | 2.2 | 2.4 | 2.9 | 3.4 | 3.8 | |
| 25 | 55 | 1.2 | 1.5 | 1.8 | 2.1 | 2.4 | 2.7 | 3 | 3.6 | 4.2 | 4.8 | |
| 35 | 77 | 1.7 | 2.1 | 2.5 | 2.9 | 3.4 | 3.8 | 4.2 | 5 | 5.9 | 6.7 | |
| 50 | 110 | 2.4 | 3 | 3.6 | 4.2 | 4.8 | 5.4 | 6 | 7.2 | 8.4 | 9.6 | |
| 60 | 132 | 2.9 | 3.6 | 4.3 | 5 | 5.8 | 6.5 | 7.2 | 8.6 | 10.1 | 11.5 | |
| 70 | 154 | 3.4 | 4.2 | 5 | 5.9 | 6.7 | 7.6 | 8.4 | 10.1 | 11.8 | 13.4 | |
| 80 | 176 | 3.8 | 4.8 | 5.8 | 6.7 | 7.7 | 8.6 | 9.6 | 11.5 | 13.4 | 15.4 | |
| 90 | 198 | 4.3 | 5.4 | 6.5 | 7.6 | 8.6 | 9.7 | 10.8 | 13 | 15.1 | 17.3 | |
| 100 | 220 | 4.8 | 6 | 7.2 | 8.4 | 9.6 | 10.8 | 12 | 14.4 | 16.8 | 19.2 | |
- * 500 mg Rocuronium Bromide Injection in 100 mL solution.

Drug Admixture Incompatibility

Rocuronium Bromide Injection is physically incompatible when mixed with the following drugs:

| | |
|---------------|---------------------------------|
| amphotericin | hydrocortisone sodium succinate |
| aztrexam | insulin |
| azathioprine | Intralipid |
| cefazolin | ketorolac |
| cloxacinil | lorazepam |
| dexamethasone | methohexital |
| diazepam | methyprednisolone |
| erythromycin | thiopental |
| famotidine | trimethoprim |
| furosemide | vancomycin |

If Rocuronium Bromide Injection is administered via the same infusion line that is also used for other drugs, it is important that this infusion line is adequately flushed between administration of Rocuronium Bromide Injection and drugs for which incompatibility with Rocuronium Bromide Injection has been demonstrated or for which compatibility with Rocuronium Bromide Injection has not been established.

Infusion solutions should be used within 24 hours of mixing. Unused portions of infusion solutions should be discarded.

Rocuronium Bromide Injection should not be mixed with alkaline solutions [see Warnings and Precautions (5.11)].

Visual Inspection

Parenteral drug products should be inspected visually for particulate matter and clarity prior to administration whenever solution and container permit. Do not use solution if particulate matter is present.

- 3
- DOSAGE FORMS AND STRENGTHS
- Rocuronium Bromide Injection is a clear, colorless to yellow/orange solution, free from visible particulate matter available as:
- 50 mg/5 mL (10 mg/mL), single-dose vials
 - 100 mg/10 mL (10 mg per mL), single-dose vials

- 4
- CONTRAINDICATIONS
- Rocuronium Bromide Injection is contraindicated in patients known to have hypersensitivity (e.g., anaphylaxis) to rocuronium bromide or other neuromuscular blocking agents [see Warnings and Precautions (5.2)].

WARNINGS AND PRECAUTIONS

- 5.1
- Appropriate Administration and Monitoring
- Rocuronium Bromide Injection should be administered in carefully adjusted dosages by or under the supervision of experienced clinicians who are familiar with the drug's actions and the possible complications of its use. Rocuronium Bromide Injection should not be administered unless facilities for intubation, mechanical ventilation, oxygen therapy, and an antagonist are immediately available. It is recommended that clinicians administering neuromuscular blocking agents such as Rocuronium Bromide Injection employ a peripheral nerve stimulator to monitor drug effect, need for additional doses, adequacy of spontaneous recovery or antagonism, and to decrease the complications of overdose if additional doses are administered.
- 5.2
- Anaphylaxis
- Severe anaphylactic reactions to neuromuscular blocking agents, including Rocuronium Bromide Injection, have been reported. These reactions have, in some cases (including cases with Rocuronium Bromide Injection), been life threatening and fatal. Due to the potential severity of these reactions, the necessary precautions, such as the immediate availability of appropriate emergency treatment, should be taken. Precautions should also be taken in those patients who have had previous anaphylactic reactions to other neuromuscular blocking agents, since cross-reactivity between neuromuscular blocking agents, both depolarizing and nondepolarizing, has been reported.

- 5.3
- Risk of Death due to Medication Errors
- Administration of Rocuronium Bromide Injection results in paralysis, which may lead to respiratory arrest and death, a progression that may be more likely to occur in a patient for whom it is not intended. Confirm proper selection of intended product and avoid confusion with other injectable solutions that are present in critical care and other clinical settings. If another healthcare provider is administering Rocuronium Bromide Injection, ensure that the intended dose is clearly labeled and communicated.
- 5.4
- Need for Adequate Anesthesia
- Rocuronium Bromide Injection has no known effect on consciousness, pain threshold, or cerebation. Therefore, its administration must be accompanied by adequate anesthesia or sedation.
- 5.5
- Residual Paralysis
- To prevent complications resulting from residual paralysis from Rocuronium Bromide Injection, it is recommended to extubate only after the patient has recovered sufficiently from neuromuscular block. Geriatric patients (65 years or older) may be at increased risk for residual neuromuscular block. Other factors which could cause residual paralysis (such as inadequate anesthesia, inadequate ventilation, or drug interactions or patient condition) should also be considered. If not used as part of standard clinical practice the use of a reversal agent should be considered, especially in those cases where residual paralysis is more likely to occur.
- 5.6
- Long-Term Use in an Intensive Care Unit
- Rocuronium Bromide Injection has not been studied for long-term use in the intensive care unit (ICU). As with other nondepolarizing neuromuscular blocking drugs, apparent tolerance to Rocuronium Bromide Injection may develop during chronic administration in the ICU. While the mechanism for development of this resistance is not known, receptor up-regulation may be a contributing factor. It is strongly recommended that neuromuscular transmission be monitored continuously during Rocuronium Bromide Injection administration and recovery with the help of a nerve stimulator. Additional doses of Rocuronium Bromide Injection or any other neuromuscular blocking agent should not be given until there is a definite response (one twitch of the train-of-four) to nerve stimulation. Prolonged paralysis and/or skeletal muscle weakness may be noted during initial attempts to wean intubated patients from ventilators who have chronically received neuromuscular blocking drugs in the ICU.

Myopathy after long-term administration of other nondepolarizing neuromuscular blocking agents in the ICU alone or in combination with corticosteroid therapy has been reported. Therefore, for patients receiving both neuromuscular blocking agents (including Rocuronium Bromide Injection) and corticosteroids, the period of use of the neuromuscular blocking agent should be limited as much as possible and the drug should be used only under the close supervision of the prescribing physician, the specific advantages of the drug outweigh the risk.

- 5.7
- Malignant Hyperthermia
- Rocuronium Bromide Injection has not been studied in malignant hyperthermia (MH)-susceptible patients. In an animal study in MH-susceptible swine, the administration of Rocuronium Bromide Injection did not appear to trigger malignant hyperthermia. Because Rocuronium Bromide Injection is always used with other agents, and the occurrence of MH during anesthesia is possible even in the absence of known triggering agents, clinicians should be familiar with early signs, confirmatory diagnosis, and treatment of MH prior to the start of any anesthetic [see Adverse Reactions (6.2)].
- 5.8
- Prolonged Circulation Time
- Conditions associated with an increased circulatory delayed time, e.g., cardiovascular disease or advanced age, in patients treated with Rocuronium Bromide Injection may be associated with a delay in onset time [see Dosage and Administration (2.6)].

- 5.9
- QT Interval Prolongation
- The overall analysis of ECG data in pediatric patients indicates that the concomitant use of Rocuronium Bromide Injection with general anesthetic agents can prolong the QTc interval [see Clinical Studies (14.3)].

- 5.10
- Conditions/Drugs Causing Potentiation of, or Resistance to, Neuromuscular Block
- Potentiation
- Nondepolarizing neuromuscular blocking agents have been found to exhibit profound neuromuscular blocking effects in cachectic or debilitated patients, patients with neuromuscular diseases, and patients with carcinomatosis.
- Certain inhalation anesthetics, particularly enflurane and isoflurane, antibiotics, magnesium salts, lithium, local anesthetics, procainamide, and quinine have been shown to increase the duration of neuromuscular block and decrease infusion requirements of neuromuscular blocking agents [see Drug Interactions (7.3)].
- In these or other patients in whom potentiation of neuromuscular block or difficulty with reversal may be anticipated, a decrease from the recommended initial dose of Rocuronium Bromide Injection should be considered [see Dosage and Administration (2.6)].
- Resistance
- Resistance to nondepolarizing agents, consistent with up-regulation of skeletal muscle acetylcholine receptors, is associated with burns, disuse atrophy, denervation, and direct muscle trauma. Receptor up-regulation may also contribute to the resistance to nondepolarizing muscle relaxants which sometimes develops in patients with cerebral palsy, patients chronically receiving anticonvulsant agents such as carbamazepine or phenytoin, or with chronic exposure to nondepolarizing agents. When Rocuronium Bromide Injection is administered to these patients, shorter durations of neuromuscular block may occur, and infusion rates may be higher due to the development of resistance to nondepolarizing muscle relaxants.

Potentiation or Resistance

Severe acid-base and/or electrolyte abnormalities may potentiate or cause resistance to the neuromuscular blocking action of Rocuronium Bromide Injection. No data are available in such patients and no dosing recommendations can be made.

Resistance to Rocuronium Bromide Injection

Rocuronium Bromide Injection-induced neuromuscular blockade was modified by alkalosis and acidosis in experimental pigs. Both respiratory and metabolic acidosis prolonged the recovery time. The potency of Rocuronium Bromide Injection was significantly enhanced in metabolic acidosis and alkalosis but was reduced in respiratory alkalosis. In addition, experience with other drugs has suggested that acute (e.g., diarrhea) or chronic (e.g., adrenocortical insufficiency) electrolyte imbalance may alter neuromuscular blockade. Since electrolyte imbalance and acid-base imbalance are usually mixed, either enhancement or inhibition may occur.

Resistance to Rocuronium Bromide Injection

Rocuronium Bromide Injection-induced neuromuscular blockade was modified by alkalosis and acidosis in experimental pigs. Both respiratory and metabolic acidosis prolonged the recovery time. The potency of Rocuronium Bromide Injection was significantly enhanced in metabolic acidosis and alkalosis but was reduced in respiratory alkalosis. In addition, experience with other drugs has suggested that acute (e.g., diarrhea) or chronic (e.g., adrenocortical insufficiency) electrolyte imbalance may alter neuromuscular blockade. Since electrolyte imbalance and acid-base imbalance are usually mixed, either enhancement or inhibition may occur.

- 5.11
- Incompatibility with Alkaline Solutions
- Rocuronium Bromide Injection, which has an acid pH, should not be mixed with alkaline solutions (e.g., barbiturate solutions) in the same syringe or administered simultaneously during intravenous infusion through the same needle.
- 5.12
- Increase in Pulmonary Vascular Resistance
- Rocuronium Bromide Injection may be associated with increased pulmonary vascular resistance, so caution is appropriate in patients with pulmonary hypertension or valvular heart disease [see Clinical Studies (14.1)].

- 5.13
- Use in Patients with Myasthenia
- In patients with myasthenia gravis or myasthenic (Eaton-Lambert) syndrome, small doses of nondepolarizing neuromuscular blocking agents, including Rocuronium Bromide Injection, may have profound effects. In such patients, a peripheral nerve stimulator and use of a small test dose may be of value in monitoring the response to administration of muscle relaxants.
- 5.14
- Extravasation
- If extravasation occurs after Rocuronium Bromide Injection administration, it may be associated with signs or symptoms of local irritation. The Rocuronium Bromide Injection or infusion should be terminated immediately and restarted in another vein.

- 6
- ADVERSE REACTIONS
- In clinical trials, the most common adverse reactions (2%) are transient hypotension and hypertension. The following adverse reactions are described, or described in greater detail, in other sections:
- Anaphylaxis [see Warnings and Precautions (5.2)]
 - Residual paralysis [see Warnings and Precautions (5.5)]
 - Myopathy [see Warnings and Precautions (5.6)]
 - Increased pulmonary vascular resistance [see Warnings and Precautions (5.12)]
- 6.1
- Clinical Trials Experience
- Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.
- Clinical studies of another rocuronium bromide injection product in the U.S. (n=1137) and Europe (n=1394) totaled 2531 patients. The following adverse reactions were reported in patients administered another rocuronium bromide injection (all events judged by investigators during the clinical trials to have a possible causal relationship):
- Adverse reactions in greater than 1% of patients: None
- Adverse reactions in less than 1% of patients (probably related or relationship unknown):
- Cardiovascular: arrhythmia, abnormal electrocardiogram, tachycardia
- Gastrointestinal: nausea, vomiting
- Respiratory: asthma (bronchospasm, wheezing, or rhinorrhea), hiccup
- Skin and Appendages: rash, injection site edema, pruritus

7.9 Propofol
The use of propofol for induction and maintenance of anesthesia does not alter the clinical duration or recovery characteristics following recommended doses of Rocuronium Bromide Injection.

7.10 Quinidine
Injection of quinidine during recovery from use of muscle relaxants is associated with recurrent paralysis. This possibility must also be considered for Rocuronium Bromide Injection [see **Warnings and Precautions (5.10)**].

7.11 Succinylcholine
The use of Rocuronium Bromide Injection before succinylcholine, for the purpose of attenuating some of the side effects of succinylcholine, has not been studied.

If Rocuronium Bromide Injection is administered following administration of succinylcholine, it should not be given until recovery from succinylcholine has been observed. The median duration of action of Rocuronium Bromide Injection 0.6 mg/kg administered after a 1 mg/kg dose of succinylcholine when T₁ returned to 75% of control was 36 minutes (range: 14-57, n=12) vs. 28 minutes (range: 17-51, n=12) without succinylcholine.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Available data from controlled trials and case series and over decades of use of rocuronium bromide in pregnant women have not identified a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. There are potential risks when rocuronium bromide is used during labor or delivery [see *Clinical Considerations*]. Based on data from umbilical cord blood sampling, rocuronium bromide is transferred across the placenta [see *Clinical Studies (14.11)*].

In animal reproduction studies, there was no evidence of teratogenicity when rocuronium bromide was administered intravenously to pregnant, conscious, nonventilated rats and rabbits at 15%-30% and 25%, respectively, the human intubation dose of 0.6-1.2 mg/kg during the period of organogenesis [see *Data*].

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Clinical Considerations

Labor or Delivery: Magnesium sulfate used in the management of pre-eclampsia or eclampsia in pregnancy may enhance neuromuscular blockade [see *Warning and Precautions 5.10*].

The use of another rocuronium bromide injection product in Cesarean section has been studied in a limited number of patients. Adverse events in this study included low APGAR scores in neonates at 5 minutes and poor intubation conditions in some pregnant women who received rocuronium bromide [see *Clinical Studies (14.11)*]. Rocuronium Bromide Injection is not recommended for rapid sequence induction in Cesarean section patients.

Data

Animal Data: Developmental toxicology studies have been performed with rocuronium bromide in pregnant, conscious, nonventilated rats and rabbits. Inhibition of neuromuscular function was the endpoint for high-dose selection. The maximum tolerated dose served as the high dose and was administered intravenously 3 times a day to rats (0.3 mg/kg, 15%-30% of human intubation dose of 0.6-1.2 mg/kg based on the body surface unit of mg/m²) from Day 6 to 17 and to rabbits (0.02 mg/kg, 25% human dose) from Day 6 to 18 of pregnancy. High-dose treatment caused acute symptoms of respiratory dysfunction due to the pharmacological activity of the drug. Teratogenicity was not observed in these animal species. The incidence of late embryonic death was increased at the high dose in rats, most likely due to oxygen deficiency. Therefore, this finding probably has no relevance for humans because immediate mechanical ventilation of the intubated patient will effectively prevent embryo-fetal hypoxia.

8.2 Lactation

Risk Summary

There are no available data on the presence of rocuronium bromide in human milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for rocuronium bromide and any potential adverse effects on the breastfed child from rocuronium bromide or from the underlying maternal condition.

8.4 Pediatric Use

The use of another rocuronium bromide injection product has been studied in 3 months to 14 years of age patients who received halothane anesthesia. Of the pediatric patients anesthetized with halothane who did not receive atropine for induction, about 80% experienced a transient increase (30% or greater) in heart rate after intubation. One of the 19 infants anesthetized with halothane and fentanyl who received atropine for induction experienced this magnitude of change [see *Dosage and Administration (2.6)*, *Clinical Studies (14.3)*].

Another rocuronium bromide injection product was also studied in pediatric patients up to 17 years of age, including neonates, under sevoflurane (induction) and isoflurane/nitrous oxide (maintenance) anesthesia. Onset time and clinical duration varied with dose, the age of the patient, and anesthetic technique. The overall analysis of ECG data in pediatric patients indicated that the concomitant use of Rocuronium Bromide Injection with general anesthetic agents can prolong the QTc interval. The data also suggest that Rocuronium Bromide Injection may increase heart rate. However, it was not possible to conclusively identify an effect of rocuronium bromide independent of that of anesthesia and other factors. Additionally, when examining plasma levels of rocuronium bromide in correlation to QTc interval prolongation, no relationship was observed [see *Dosage and Administration (2.6)*, *Warnings and Precautions (5.9)*, *Clinical Studies (14.3)*].

Rocuronium Bromide Injection is not recommended for rapid sequence intubation in pediatric patients. Recommendations for use in pediatric patients are discussed in other sections of labeling [see *Dosage and Administration (2.6)* *Clinical Pharmacology (12.2)*].

8.5 Geriatric Use

Another rocuronium bromide injection product was administered to 140 geriatric patients (65 years of age or older) in U.S. clinical trials and 128 geriatric patients in European clinical trials. The observed pharmacokinetic profile of rocuronium bromide for geriatric patients (n=20) was similar to that for other adult surgical patients [see *Clinical Pharmacology (12.3)*]. However, onset time and duration of action of rocuronium bromide were slightly longer for geriatric patients (n=43) in clinical trials. Clinical experiences and recommendations for use of Rocuronium Bromide Injection in geriatric patients are discussed in other sections of the labeling [see *Dosage and Administration (2.6)*, *Warnings and Precautions (5.5)*, *Clinical Pharmacology (12.2)*, *Clinical Studies (14.2)*].

8.6 Patients with Hepatic Impairment

Because rocuronium bromide is primarily excreted by the liver, it should be used with caution in patients with clinically significant hepatic impairment. Another rocuronium bromide injection product (0.6 mg/kg) was studied in a limited number of patients (n=9) with clinically significant hepatic impairment under steady-state isoflurane anesthesia. After rocuronium bromide injection administration (0.6 mg/kg), the median (range) clinical duration of 60 (35-166) minutes was moderately prolonged compared to 42 minutes in patients with normal hepatic function. The median recovery time of 53 minutes was also prolonged in patients with cirrhosis compared to 20 minutes in patients with normal hepatic function. Four of 8 patients with cirrhosis, who received the other rocuronium bromide injection product (0.6 mg/kg) under opioid/nitrous oxide/oxygen anesthesia, did not achieve complete block. These findings are consistent with the increase in volume of distribution at steady state observed in patients with significant hepatic impairment [see *Clinical Pharmacology (12.3)*]. If used for rapid sequence induction in patients with ascites, an increased initial dosage of Rocuronium Bromide Injection may be necessary to assure complete block. Duration will be prolonged in these cases. The use of Rocuronium Bromide Injection doses higher than 0.6 mg/kg has not been studied [see *Dosage and Administration (2.6)*].

8.7 Patients with Renal Impairment

Due to the limited role of the kidney in the excretion of rocuronium bromide, usual dosing recommendations Rocuronium Bromide Injection for should be followed. In patients with renal dysfunction, the duration of neuromuscular blockade was not prolonged; however, there was substantial individual variability (range: 22-90 minutes) [see *Clinical Pharmacology (12.3)*].

10 OVERDOSAGE

Overdosage with neuromuscular blocking agents may result in neuromuscular block beyond the time needed for surgery and anesthesia. The primary treatment is maintenance of a patent airway, controlled ventilation, and adequate sedation until recovery of normal neuromuscular function is assured. Once evidence of recovery from neuromuscular block is observed, further recovery may be facilitated by administration of an anticholinesterase agent in conjunction with an appropriate anticholinergic agent.

Reversal of Neuromuscular Blockade: Anticholinesterase agents should not be administered prior to the demonstration of some spontaneous recovery from neuromuscular blockade. The use of a nerve stimulator to document recovery is recommended.


Patients should be evaluated for adequate clinical evidence of neuromuscular recovery, e.g., 5-second head lift, adequate phonation, ventilation, and upper airway patency. Ventilation must be supported while patients exhibit any signs of muscle weakness.

Recovery may be delayed in the presence of debilitation, carcinomatosis, and concomitant use of certain drugs which enhance neuromuscular blockade or separately cause respiratory depression. Under such circumstances the management is the same as that of prolonged neuromuscular blockade.

11 DESCRIPTION

Rocuronium bromide is a nondepolarizing neuromuscular blocking agent with a rapid to intermediate onset depending on dose and intermediate duration. Rocuronium bromide is chemically designated as 1-[1-(7-(acetyloxy)-3 α -hydroxy-2 β -(4-morpholinyl)-5 α -androstan-16 β -yl]-1-(2-propenyl) pyridinium bromide.

The structural formula is:



Rocuronium bromide USP is an almost white to pale yellow, hygroscopic powder. Rocuronium bromide is soluble in water and slightly soluble in methanol. The chemical structure of rocuronium bromide is C₃₂H₄₂BrN₃O₄ with a molecular weight of 609.70. The partition coefficient of rocuronium bromide in n-octanol/water is 0.5 at 20°C.

Rocuronium Bromide Injection is supplied as a sterile, nonpyrogenic, isotonic solution that is clear, colorless to yellow/orange, for intravenous use only. Each mL contains 10 mg rocuronium bromide (equivalent to 8.69 mg of rocuronium) and inactive ingredients 17.1 mL hydrochloric acid, 8 mg sodium chloride. The solution pH is adjusted to 2.8 to 3.2 with hydrochloric acid and/or sodium hydroxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Rocuronium Bromide Injection is a nondepolarizing neuromuscular blocking agent with a rapid to intermediate onset depending on dose and intermediate duration. It acts by competing for cholinergic receptors at the motor end-plate. This action is antagonized by acetylcholinesterase inhibitors, such as neostigmine and edrophonium.

12.2 Pharmacodynamics

The Rocuronium Bromide Injection dose required to produce 95% suppression (ED₉₅) of the first [T₁] mechanomyographic [MMG] response of the adductor pollicis muscle [thumb] to indirect supra-maximal train-of-four stimulation of the ulnar nerve) during opioid/nitrous oxide/oxygen anesthesia is approximately 0.3 mg/kg. Patient variability around the ED₉₅ dose suggests that 50% of patients will exhibit T₁ depression of 91% to 97%.

Table 4 presents intubating conditions in patients with intubation initiated at 60 to 70 seconds.

Table 4: Percent of Excellent or Good Intubating Conditions and Median (Range) Time to Completion of Intubation in Patients with Intubation Initiated at 60 to 70 Seconds

| rocuronium bromide injection Dose (mg/kg) Administered Over 5 sec | Percent of Patients with Excellent or Good Intubating Conditions | Time to Completion of Intubation (min) |
|---|--|--|
| Adults* 18 to 64 yrs 0.45 (n=43) 0.6 (n=51) | 86% 96% | 1.6 (1.0-7.0) 1.6 (1.0-3.2) |
| Infants† 3 mo to 1 yr 0.6 (n=18) | 100% | 1.0 (1.0-1.5) |
| Pediatric‡ 1 to 12 yrs 0.6 (n=12) | 100% | 1.0 (0.5-2.3) |

* Excludes patients undergoing Cesarean section.

† Pediatric patients were under halothane anesthesia.

‡ Excellent intubating conditions=jaw relaxed, vocal cords apart and immobile, no diaphragmatic movement.

Good intubating conditions=same as excellent but with some diaphragmatic movement.

Table 5 presents the time to onset and clinical duration for the initial dose of Rocuronium Bromide Injection under opioid/nitrous oxide/oxygen anesthesia in adults and geriatric patients, and under halothane anesthesia in pediatric patients.

Table 5: Median (Range) Time to Onset and Clinical Duration Following Initial (Intubating) Dose During Opioid/Nitrous Oxide/Oxygen Anesthesia (Adults) and Halothane Anesthesia (Pediatric Patients)

| rocuronium bromide injection Dose (mg/kg) Administered Over 5 sec | Time to ≥80% Block (min) | Time to Maximum Block (min) | Clinical Duration (min) |
|---|--|---|--|
| Adults 18 to 64 yrs 0.45 (n=50) 0.6 (n=142) 0.9 (n=20) 1.2 (n=18) | 1.3 (0.8-6.2) 1.0 (0.4-6.0) 1.1 (0.3-3.8) 0.7 (0.4-1.7) | 3.0 (1.3-8.2) 1.8 (0.6-13.0) 1.4 (0.8-6.2) 1.0 (0.6-4.7) | 22 (12-31) 31 (15-85) 58 (27-111) 67 (38-160) |
| Geriatric ≥65 yrs 0.6 (n=31) 0.9 (n=5) 1.2 (n=7) | 2.3 (1.0-8.3) 2.0 (1.0-3.0) 2.0 (0.8-3.5) | 3.7 (1.3-11.3) 2.5 (1.2-5.0) 1.3 (1.2-4.7) | 46 (22-73) 62 (49-75) 94 (64-136) |
| Infants 3 mo to 1 yr 0.6 (n=9) | — | 0.8 (0.3-3.0) 0.7 (0.5-0.8) | 41 (24-68) 40 (27-70) |
| Pediatric 1 to 12 yrs 0.6 (n=27) 0.6 (n=18) | 0.8 (0.4-2.0) — | 1.0 (0.5-3.3) 0.5 (0.3-1.0) | 26 (17-39) 30 (17-56) |

n=the number of patients who had time to maximum block recorded.

Clinical duration=time until return to 25% of control T₁. Patients receiving doses of 0.45 mg/kg who achieved less than 90% block (16% of these patients) had about 12 to 15 minutes to 25% recovery.

Table 6 presents the time to onset and clinical duration for the initial dose of Rocuronium Bromide Injection (rocuronium bromide) Injection under sevoflurane (induction) and isoflurane/nitrous oxide (maintenance) anesthesia in pediatric patients.

Table 6: Median (Range) Time to Onset and Clinical Duration Following Initial (Intubating) Dose During Sevoflurane (Induction) and Isoflurane/Nitrous Oxide (maintenance) Anesthesia (Pediatric Patients)

| rocuronium bromide injection Dose (mg/kg) Administered Over 5 sec | Time to Maximum Block (min) | Time to Reappearance T ₂₅ (min) |
|---|---|--|
| Neonates birth to <28 days 0.45 (n=5) 0.6 (n=10) 1 (n=6) | 1.1 (0.6-2.2) 1.0 (0.2-2.1) 0.6 (0.3-1.8) | 40.3 (32.5-62.6) 49.7 (16.6-119.0) 41.4 (32.6-136.3) |
| Infants 28 days to ≤3 mo 0.45 (n=9) 0.6 (n=11) 1 (n=5) | 0.5 (0.4-1.3) 0.4 (0.2-0.8) 0.3 (0.2-0.7) | 49.1 (13.5-79.9) 58.8 (32.3-87.8) 103.3 (90.8-155.4) |
| Toddlers >3 mo to ≤2 yrs 0.45 (n=17) 0.6 (n=29) 1 (n=15) | 0.8 (0.3-1.9) 0.6 (0.2-1.6) 0.5 (0.2-1.5) | 39.2 (16.9-59.4) 44.2 (18.9-68.8) 72.0 (36.2-128.2) |
| Children >2 yrs to ≤11 yrs 0.45 (n=14) 0.6 (n=37) 1 (n=16) | 0.9 (0.4-1.9) 0.6 (0.2-1.4) 0.7 (0.4-1.2) | 21.5 (17.5-38.0) 36.7 (20.1-65.9) 53.1 (31.2-89.9) |
| Adolescents >11 to ≤17 yrs 0.45 (n=18) 0.6 (n=31) 1 (n=14) | 1.0 (0.5-1.7) 0.9 (0.2-2.1) 0.7 (0.5-1.2) | 37.5 (18.3-65.7) 41.4 (16.3-91.2) 67.1 (25.6-93.8) |

n=the number of patients with the highest number of observations for time to maximum block or reappearance T₂₅.

The time to 80% or greater block and clinical duration as a function of dose are presented in **Figures 1 and 2**.

Figure 1: Time to 80% or Greater Block vs. Initial Dose of rocuronium bromide injection by Age Group (Median, 25th and 75th Percentile, and Individual Values)






Figure 2: Duration of Clinical Effect vs. Initial Dose of rocuronium bromide injection by Age Group (Median, 25th and 75th Percentile, and Individual Values)



The clinical durations for the first 5 maintenance doses, in patients receiving 5 or more maintenance doses are represented in **Figure 3** [see *Dosage and Administration (2.4)*].

Figure 3: Duration of Clinical Effect vs. Number of rocuronium bromide injection Maintenance Doses, by Dose



Once spontaneous recovery has reached 25% of control T₁, the neuromuscular block produced by rocuronium bromide injection is readily reversed with anticholinesterase agents, e.g., edrophonium or neostigmine.

The median spontaneous recovery from 25% to 75% T₁ was 13 minutes in adult patients. When neuromuscular block was reversed in 36 adults at a T₁ of 22% to 27%, recovery to a T₁ of 89 (50-132)% and T₄/T₁ of 69 (38-92)% was achieved within 5 minutes. Only 5 of 320 adults reversed received an additional dose of reversal agent. The median (range) dose of neostigmine was 0.04 (0.01-0.09) mg/kg and the median (range) dose of edrophonium was 0.5 (0.3-1.0) mg/kg.

In geriatric patients (n=51) reversed with neostigmine, the median T₄/T₁ increased from 40% to 88% in 5 minutes.

In clinical trials with halothane, pediatric patients (n=27) who received 0.5 mg/kg edrophonium had increases in the median T₄/T₁ from 37% at reversal to 93% after 2 minutes. Pediatric patients (n=58) who received 1 mg/kg edrophonium had increases in the median T₄/T₁ from 72% at reversal to 100% after 2 minutes. Infants (n=10) who were reversed with 0.03 mg/kg neostigmine recovered from 25% to 75% T₁ within 4 minutes.

There were no reports of less than satisfactory clinical recovery of neuromuscular function.

The neuromuscular blocking action of Rocuronium Bromide Injection may be enhanced in the presence of potent inhalation anesthetics [see *Drug Interactions (7.3)*].

Hemodynamics

There were no dose-related effects on the incidence of changes from baseline (30% or greater) in mean arterial blood pressure (MAP) or heart rate associated with rocuronium bromide injection administration over the dose range of 0.12 to 1.2 mg/kg (4 x ED₉₅) within 5 minutes after rocuronium bromide injection administration and prior to intubation. Increases or decreases in MAP were observed in 2% to 5% of geriatric and other adult patients, and in about 1% of pediatric patients. Heart rate changes (30% or greater) occurred in 0% to 2% of geriatric and other adult patients. Tachycardia (30% or greater) occurred in 12 of 127 pediatric patients. Most of the pediatric patients developing tachycardia were from a single study where the patients were anesthetized with halothane and who did not receive atropine for induction [see *Clinical Studies (14.3)*]. In US studies, laryngoscopy and tracheal intubation following rocuronium bromide injection administration were accompanied by transient tachycardia (30% or greater increases) in about one-third of adult patients under opioid/nitrous oxide/oxygen anesthesia. Animal studies with methoxyflurane, isoflurane, and sevoflurane anesthesia following Rocuronium Bromide Injection administration is less than rocuronium but greater than pancuronium. The tachycardia observed in some patients may result from this vagal blocking activity.

Histamine Release

In studies of histamine release, clinically significant concentrations of plasma histamine occurred in 1 of 88 patients. Clinical signs of histamine release (flushing, rash, or bronchospasm) associated with the administration of Rocuronium Bromide Injection were assessed in clinical trials and reported in 9 of 1137 (0.8%) patients.

12.3 Pharmacokinetics

Adult and Geriatric Patients

In an effort to maximize the information gathered in the *in vivo* pharmacokinetic studies, the data from the studies was used to develop population-based estimates of the pharmacokinetic parameters represented (e.g., geriatric, pediatric, renal, and hepatic impairment). These population-based estimates and a measure of the estimate variability are contained in the following section.

Following intravenous administration of rocuronium bromide injection, plasma levels of rocuronium follow a three-compartment open model. The rapid distribution half-life is 1 to 2 minutes and the slower distribution half-life is 14 to 18 minutes. Rocuronium is approximately 30% bound to human plasma proteins. In geriatric and other adult patients undergoing either opioid/nitrous oxide/oxygen or inhalational anesthesia, the observed pharmacokinetic profile was essentially unchanged [see *Dosage and Administration (2.6)*].

Table 7: Mean (SD) Pharmacokinetic Parameters in Adults (n=22; ages 27 to 58 yrs) and Geriatric (n=20; 65 yrs or greater) During Opioid/Nitrous Oxide/Oxygen Anesthesia

| PK Parameters | Adults (Ages 27-58 yrs) | Geriatrics (≥65 yrs) |
|---|-------------------------|----------------------|
| Clearance (L/kg/hr) | 0.25 (0.08) | 0.21 (0.06) |
| Volume of Distribution at Steady State (L/kg) | 0.25 (0.04) | 0.22 (0.03) |
| t _{1/2} β Elimination (hr) | 1.4 (0.4) | 1.5 (0.4) |

In general, studies with normal adult subjects did not reveal any differences in the pharmacokinetics of rocuronium due to gender.

Studies of distribution, metabolism, and excretion in cats and dogs indicate that rocuronium is eliminated primarily by the liver. The rocuronium analog 17-desacetyl-rocuronium, a metabolite, has been rarely observed in the plasma or urine of humans administered single doses of 0.5 to 1 mg/kg with or without a subsequent infusion (for use of 17-desacetyl-rocuronium). In the 17-desacetyl-rocuronium study, approximately one-twentieth the neuromuscular blocking potency of rocuronium. The effects of renal failure and hepatic disease on the pharmacokinetics and pharmacodynamics of rocuronium in humans are consistent with these findings.

In general, patients undergoing cadaver kidney transplant have a small reduction in clearance which is offset pharmacokinetically by a corresponding increase in volume, such that the net effect is an unchanged plasma half-life. Patients with demonstrated liver cirrhosis have a marked increase in their volume of distribution resulting in a plasma half-life approximately twice that of patients with normal hepatic function. **Table 8** shows the pharmacokinetic parameters in subjects with either impaired renal or hepatic function.

Table 8: Mean (SD) Pharmacokinetic Parameters in Adults with Normal Renal and Hepatic Function (n=10, ages 23 to 65), Renal Transplant Patients (n=10, ages 21 to 45), and Hepatic Dysfunction Patients (n=9, ages 31 to 67) During Isoflurane Anesthesia

| PK Parameters | Normal Renal and Hepatic Function | Renal Transplant Patients | Hepatic Dysfunction Patients |
|---|-----------------------------------|---------------------------|------------------------------|
| Clearance (L/kg/hr) | 0.16 (0.05)* | 0.13 (0.04) | 0.13 (0.06) |
| Volume of Distribution at Steady State (L/kg) | 0.26 (0.03) | 0.34 (0.11) | 0.53 (0.14) |
| t _{1/2} β Elimination (hr) | 2.4 (0.8)* | 2.4 (1.1) | 4.3 (2.6) |

* Differences in the calculated t_{1/2} β and CI between this study and the study in young adults vs. geriatrics (≥65 years) is related to the different sample populations and anesthetic techniques.

The net result of these findings is that subjects with renal failure have clinical durations that are similar to but somewhat more variable than the duration that one would expect in subjects with normal renal function. Hepatically impaired patients, due to the large increase in volume, may demonstrate clinical durations approaching 1.5 times that of subjects with normal hepatic function. In both populations the clinician should individualize the dose to the needs of the patient [see *Dosage and Administration (2.6)*].

Tissue redistribution accounts for most (about 80%) of the initial amount of rocuronium administered. As tissue compartments fill with continued dosing (4-8 hours), less drug is redistributed away from the site of action and, for this reason, the rate to maintain neuromuscular blockade falls to about 20% of the initial infusion rate. The use of a loading dose and a smaller infusion rate reduces the need for adjustment of dose.

Pediatric Patients

Under halothane anesthesia, the clinical duration of effects of rocuronium bromide injection did not vary with age in patients 4 months to 8 years of age. The terminal half-life and other pharmacokinetic parameters of rocuronium in these pediatric patients are presented in **Table 9**.

Table 9: Mean (SD) Pharmacokinetic Parameters of Rocuronium in Pediatric Patients (ages 3 to less than 12 mos, n=6; 1 to less than 3 yrs, n=5; 3 to less than 8 yrs, n=7) During Halothane Anesthesia

| PK Parameters | Patient Age Range | 3 to <12 mos | 1 to <3 yrs | 3 to <8 yrs |
|---|-------------------|--------------|-------------|-------------|
| Clearance (L/kg/hr) | | 0.35 (0.08) | 0.32 (0.07) | 0.44 (0.16) |
| Volume of Distribution at Steady State (L/kg) | | 0.30 (0.04) | 0.26 (0.06) | 0.21 (0.03) |
| t _{1/2} β Elimination (hr) | | 1.3 (0.5) | 1.1 (0.7) | 0.8 (0.3) |

Pharmacokinetics of rocuronium bromide injection were evaluated using a population analysis of the pooled pharmacokinetic datasets from 2 trials under sevoflurane (induction) and isoflurane/nitrous oxide (maintenance) anesthesia. All pharmacokinetic parameters were found to be linearly proportional to body weight. In patients under the age of 18 years, clearance (CL) and volume of distribution (V_{ss}) increase with bodyweight (kg) and age (years). As a result, the terminal half-life of rocuronium bromide injection decreases with increasing age from 1.1 hour to 0.7-0.8 hour. **Table 10** presents the pharmacokinetic parameters in the different age groups in the studies with sevoflurane (induction) and isoflurane/nitrous oxide (maintenance) anesthesia.

Table 10: Mean (SD) Pharmacokinetic Parameters of Rocuronium in Pediatric Patients During Sevoflurane (Induction) and Isoflurane/Nitrous Oxide (maintenance) Anesthesia

| PK Parameters | Patient Age Range | | | | |
|-------------------------------|-------------------|-------------------|-----------------|--------------|---------------|
| | Birth to <28 days | 28 days to <3 mos | 3 mos to <2 yrs | 2 to <11 yrs | 11 to <17 yrs |
| CL (L/kg/hr) | 0.31 (0.07) | 0.30 (0.08) | 0.33 (0.10) | 0.35 (0.09) | 0.29 (0.14) |
| Volume of Distribution (L/kg) | 0.42 (0.06) | 0.31 (0.03) | 0.23 (0.03) | 0.18 (0.02) | 0.18 (0.01) |
| t _{1/2} β (hr) | 1.1 (0.2) | 0.9 (0.3) | 0.8 (0.2) | 0.7 (0.2) | 0.8 (0.3) |