

www.fresenius-kabi.com/us

GTIN: 00363323821104

For subcutaneous or intravenous use.

3.5 mg per vial

Bortezomib
for Injection

Bortezomib for Injection

3.5 mg per vial

For subcutaneous or
intravenous use.

Usual Dosage: See package insert.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from light (keep in outer carton). Discard unused portion.

The container closure is not made with natural rubber latex.



Lake Zurich, IL 60047

www.fresenius-kabi.com/us

Reconstitution Information

SUBCUTANEOUS INJECTION ONLY

0.9% NaCl Add **1.4 mL**
0.9% Sodium Chloride

To make **2.5 mg / mL**
final concentration

INTRAVENOUS INJECTION ONLY

0.9% NaCl Add **3.5 mL**
0.9% Sodium Chloride

To make **1 mg / mL**
final concentration

Sample Label. Please see
package insert
for complete
prescribing information.

NDC 63323-821-10

Bortezomib for Injection

3.5 mg per vial

For subcutaneous or
intravenous use.

Caution: Cytotoxic Agent

Rx only
Single Dose Vial
Discard unused portion.



See Side Panel for
Reconstitution Information

861210

Sterile, lyophilized.

Each vial contains: Bortezomib 3.5 mg and mannitol 35 mg.

Reconstitution Information:

Subcutaneous injection only: Add 1.4 mL of 0.9% Sodium Chloride to each 3.5 mg single dose vial for the final concentration of 2.5 mg/mL. See package insert.

Intravenous injection only: Add 3.5 mL of 0.9% Sodium Chloride to each 3.5 mg single dose vial for the final concentration of 1 mg/mL. See package insert.



3 63323-821-10 4

621572



3 63323-821-10 4