



Do not use Diluents
containing Dextrose
(α -D-glucose)

10 Single-Dose Vials

Discard unused portion.

Preservative free.

For intravenous infusion after dilution.

Caspofungin Acetate for Injection

NDC 63323-356-10

Caspofungin Acetate for Injection

50 mg per vial*

*Each vial contains: 50 mg of caspofungin equivalent to 55.5 mg of caspofungin acetate.

Inactive ingredients: Arginine 100 mg and hydrochloric acid/sodium hydroxide to adjust pH.

Reconstitution: Reconstitute with 10.8 mL of diluent to obtain a concentration of 5 mg per mL. Requires further dilution prior to infusion. There are no preservatives or bacteriostatic agents in this product.

Usual dosage: See package insert.

STORE AT: 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

The container closure is not made with natural rubber latex.

LOT/EXP

621514B



Caspofungin Acetate for Injection

50 mg per vial*



(01)20363323356105



Lake Zurich, IL 60047

www.fresenius-kabi.com/us

Sample label. Please see package insert
for complete prescribing information.

NDC 63323-356-10

Caspofungin Acetate for Injection

50 mg per vial*

For intravenous infusion after dilution.

Preservative free.

Discard unused portion.

Rx only

10 Single-Dose Vials

Do not use Diluents
containing Dextrose
(α -D-glucose)



Caspofungin Acetate for Injection

50 mg per vial*

10 Single-Dose Vials

