



Do not use Diluents
containing Dextrose
(α-D-glucose)

For intravenous infusion after dilution.
Preservative free.
Discard unused portion.
Rx only
10 Single-Dose Vials

NDC 63323-358-10
**Caspofungin
Acetate**
for Injection
70 mg per vial*

**Caspofungin
Acetate**
for Injection

70 mg per vial*

*Each vial contains: 70 mg of caspofungin equivalent to 77.7 mg of caspofungin acetate.
Inactive ingredients: Arginine 140 mg and hydrochloric acid/sodium hydroxide to adjust pH.
Reconstitution: Reconstitute with 10.8 mL of diluent to obtain a concentration of 7 mg per mL. Requires further dilution prior to infusion. There are no preservatives or bacteriostatic agents in this product.
Usual dosage: See package insert.
STORE AT: 20° to 25°C (68° to 77°F) (see USP Controlled Room Temperature).
The container closure is not made with natural rubber latex.

LOT/EXP

621512B



**Caspofungin
Acetate**
for Injection

70 mg per vial*



(01)20363323358109



Lake Zurich, IL 60047

www.fresenius-kabi.com/us

NDC 63323-358-10

**Caspofungin
Acetate**
for Injection

70 mg per vial*

For intravenous infusion after dilution.
Preservative free.
Discard unused portion.

Rx only
10 Single-Dose Vials

Do not use Diluents
containing Dextrose
(α-D-glucose)



**Caspofungin
Acetate**
for Injection

70 mg per vial*

10 Single-Dose Vials



**Sample label. Please see package insert
for complete prescribing information.**