



Cisatracurium
Besylate Injection, USP
200 mg per 20 mL*
(10 mg per mL)
For ICU use only.
For intravenous injection.
Preservative free.
Warning: Paralyzing Agent. Causes Respiratory Arrest. Facilities must be immediately available for artificial respiration.
10 x 20 mL
Single Dose Vials
Rx only

Sample label. Please see
package insert for complete
prescribing information.

Cisatracurium
Besylate Injection, USP

200 mg per 20 mL*
(10 mg per mL)

For ICU use only.



3 NDC 63323-418-20 5

621469C

Cisatracurium
Besylate Injection, USP

200 mg per 20 mL*
(10 mg per mL)

For ICU use only.



Lake Zurich, IL 60047

www.fresenius-kabi.us

Cisatracurium
Besylate Injection, USP

200 mg per 20 mL*
(10 mg per mL)

For ICU use only.

For intravenous injection.

Preservative free.

Warning: Paralyzing Agent. Causes Respiratory Arrest. Facilities must be immediately available for artificial respiration.

10 x 20 mL
Single Dose Vials
Rx only

Sterile, Nonpyrogenic.

*Each mL contains: cisatracurium besylate equivalent to 10 mg cisatracurium, water for injection. Benzenesulfonic acid is added to adjust the pH.

Usual dosage: See package insert.

REFRIGERATE AT: 2° to 8°C (36° to 46°F). Protect from light. Retain in carton until time of use. Do not freeze. Upon removal from refrigeration to room temperature, use within 21 days even if rerefrigerated. The container closure is not made with natural rubber latex.



Cisatracurium
Besylate Injection, USP

200 mg per 20 mL*
(10 mg per mL)

For ICU use only.



3 NDC 63323-418-20 5

10 x 20 mL
Single Dose Vials
Rx only

