


Sample Label. Please see package insert for complete prescribing information.

509930

CTIN: 0036323599300

2 grams per vial

tor Injection, USP Mesylate **Deferoxamine**



NDC 63323-599-30

Deferoxamine Mesylate

for Injection, USP

2 grams per vial

For subcutaneous, intramuscular or intravenous use.

Deferoxamine Mesylate

for Injection, USP

2 grams per vial

STORE AT: 20°C to 25°C (68°F to 77°F) [see USP Controlled Room Temperature1. Keep this and all drugs out of the reach of children. The container closure is not made with natural

NDC 63323-599-30

Deferoxamine Mesylate

for Injection, USP

2 grams per vial



FRESENIUS Lake Zurich, IL 60047 for Injection, USP 2 grams per vial

Deferoxamine

Mesylate

Sterile.

509930

Each vial contains: Deferoxamine mesylate USP.

2 grams in lyophilized form. Deferoxamine mesylate reconstituted with sterile water for injection should be used immediately after reconstitution (commencement of treatment within 3 hours) for microbiological safety. When reconstituted under validated aseptic conditions (in a sterile

product may be stored at room temperature a maximum period of 24 hours before use. Do not refrigerate reconstituted solution. Turbid solutions should not be used.

laminar flow hood using aseptic technique), the

Usual dosage: See package insert.

2 gram vial is for single use only.

Single-Dose Vial







rubber latex.

