

**Sample label.
Please see
package insert
for complete
prescribing
information.**

APP Rx only

• Use strict aseptic technique
• Contains EDTA, which inhibits
microbial growth up to 12 hours,
discard within 12 hours of
opening

Ten 20 mL vials
For Single Patient Use Only

SHAKE WELL BEFORE USING

FOR INTRAVENOUS ADMINISTRATION

200 mg per 20 mL
(10 mg per mL)

DIPRIVAN[®]
(Propofol) INJECTABLE
EMULSION, USP

NDC 63323-269-29 260929

M027702/01 US
621280A



NDC 63323-269-29 260929
DIPRIVAN[®]
(Propofol) INJECTABLE
EMULSION, USP

FOR INTRAVENOUS ADMINISTRATION
SHAKE WELL BEFORE USING

200 mg per 20 mL
(10 mg per mL)

Ten 20 mL vials
For Single Patient Use Only



Rx only APP

Sterile, nonpyrogenic
Dosage: See package insert.
In addition to the active component, propofol, the formulation contains: soybean oil (100 mg/mL), glycerol (22.5 mg/mL), egg lecithin (12 mg/mL) and disodium edetate (0.005%); with sodium hydroxide to adjust pH.
DIPRIVAN Injection should be administered only by persons trained in the administration of general anesthesia and not involved in the conduct of the surgical/diagnostic procedure.
Patients should be continuously monitored and facilities for maintenance of a patent airway, artificial ventilation, and oxygen enrichment and circulatory resuscitation must be immediately available.
Store between 4° to 25°C (40° to 77°F). Do not freeze.

DIPRIVAN is a trademark of
Fresenius Kabi USA, LLC.
Manufactured for:
APP
Fresenius Kabi USA, LLC
Lake Zurich, IL 60047
Made in Austria

NDC 63323-269-29 260929
DIPRIVAN[®]
(Propofol) INJECTABLE
EMULSION, USP

FOR INTRAVENOUS ADMINISTRATION
SHAKE WELL BEFORE USING

200 mg per 20 mL
(10 mg per mL)

Ten 20 mL vials
For Single Patient Use Only

• Use strict aseptic technique
• Contains EDTA, which inhibits
microbial growth up to 12 hours,
discard within 12 hours of
opening

Rx only

NDC 63323-269-29 260929
DIPRIVAN[®]
(Propofol) INJECTABLE
EMULSION, USP

200 mg per 20 mL
(10 mg per mL)

FOR INTRAVENOUS ADMINISTRATION
SHAKE WELL BEFORE USING
Ten 20 mL vials
For Single Patient Use Only

• Use strict aseptic technique
• Contains EDTA, which inhibits
microbial growth up to 12 hours,
discard within 12 hours of
opening

Rx only

APP
3 63323 26929 6