EXP S/N

Manufactured for:

| FRESENIUS | KABI
| Lake Zurich, |L 60047

NDC 63323-269-50 500 mg per 50 mL

Sample label.

package insert

for complete

prescribing

information.

Please see

FOR INTRAVENOUS ADMINISTRATION Twenty 50 mL Vials Single-Dose Vials

Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

SHAKE WELL BEFORE USING TWenty 50 mL Vials Single-Dose Vials 500 mg per 50 mL

FOR INTRAVENOUS ADMINISTRATION



DIPRIVAN should be administered only by persons trained in the administration of general anesthesia and not involved in the conduct of the surgical/diagnostic procedure. **Facilities for maintenance of a patent airway, artificial ventilation, and oxygen enrichment and circulatory resuscitation must be immediately available.** Store between 4°C to 25°C (40°F to 77°F). Do not freeze.



discard within 12 hours of opening microbial growth up to 12 hours, Contains EDTA, which inhibits Use strict aseptic technique

Single-Dose Vials Twenty 50 mL Vials SHAKE WELL BEFORE USING FOR INTRAVENDUS ADMINISTRATION

> (10 mg per mL) 200 mg per 50 mL

Propofol) emulsion, usp



NDC 63323-269-50 260950 DIPRIVAN® (Proposol) injectable emulsion, usp

500 mg per 50 mL (10 mg per mL)

FOR INTRAVENOUS ADMINISTRATION

SHAKE WELL BEFORE USING

Twenty 50 mL Vials **Single-Dose Vials** 

- Use strict aseptic technique
- Contains EDTA, which inhibits microbial growth up to 12 hours, discard within 12 hours of opening

Rx only

