## LLG/LL67883353119 (50 mg per mL) 2.5 grams per 50 mL

Injection, USP **Fluorouracil** 

Sterile, Nonpyrogenic

Each mL contains: 50 mg fluorouracil; pH is adjusted to approximately 9.2 with sodium hydroxide.

The container closure is not made with natural rubber latex.

**FRESENIUS** Lake Zurich, IL 60047 If a prompt fluid transfer is not possible, a maximum of 4 hours from the time of initial entry is allowed to complete the transferring operations. Discard the contents no

later than 4 hours after initial closure puncture. (See Package Insert for complete dosage information and proper use of container.)

Store at 20°C to 25°C (68°F to 77°F)

Usual Dosage: Swab stopper with an antiseptic solution. A single entry

through the vial closure should be

dispensing set or transfer device.

made utilizing a suitable sterile

Isee USP Controlled Room

Temperature]. DO NOT FREEZE. PROTECT FROM LIGHT.

**FRESENIUS** 

for complete prescribing information.

101751

Sample label. Please see package insert

## **Fluorouracil** Injection, USP

NDC 63323-117-51

PHARMACY BULK PACKAGE-Not for Direct Infusion

## 2.5 grams per 50 mL

(50 mg per mL) FOR INTRAVENOUS USE ONLY

## 50 mL Rx only





