

NDC 65219-383-50

**Rx Only**

# Iodixanol Injection, USP

**320 mg Iodine/mL**

**For Intra-arterial and  
Intravenous Use Only**

in polypropylene bottle

**10 × 150 mL Single-Dose Bottles**

**Discard Unused Portion**

**Sterile Aqueous Injection**

NOT FOR INTRATHECAL USE

For CT of the head and body, iodixanol injection, USP may be used with an automated contrast injection system or contrast media management system cleared for use with iodixanol injection, USP.

Each 1 mL contains 652 mg of iodixanol (320 mg organically bound iodine), 0.044 mg calcium chloride dihydrate, 1.11 mg sodium chloride, 1.2 mg tromethamine, and 0.1 mg edetate calcium disodium. The pH is adjusted to 7.4 with hydrochloric acid and/or sodium hydroxide to achieve a range between pH 6.8 and 7.7. Osmolality is 290 mOsm/kg water. No preservative added. Each bottle for one procedure only. If used with an automated contrast management system, maximum use time is 4 hours after initial puncture. Discard unused portion.

Usual Dosage: For indications and dosage, see package insert.

Protect from light if removed from carton. Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Do not freeze.

Manufactured for:



Lake Zurich, IL 60047

[www.fresenius-kabi.com/us](http://www.fresenius-kabi.com/us)

For inquiries call 1-800-551-7176

Made in China

42635A

SP23EUS02

**Sample Label.  
Please see  
package insert  
for complete  
prescribing  
information.**



(01)20365219383509