

Manufactured by:
**FRESENIUS
KABI**
Uppsala, Sweden
www.freseniuskabinutrition.com

KABIVEN®

**Amino Acids, Electrolytes, Dextrose, and Lipid Injectable Emulsion,
for intravenous use**

(3.3%, 0.7%*, 10.8% and 3.9%), No sulfites added CENTRAL INFUSION

2053 mL



**TO ACTIVATE BAG: WITH PORTS POINTING AWAY FROM YOU, ROLL *TIGHTLY* FROM TOP OF BAG
DOWN TOWARD PORTS UNTIL BOTH VERTICAL SEALS BREAK AND ENTIRE CONTENTS ARE WHITE.**



R_x only

SEE INSTRUCTIONS FOR USE



READ THIS

• INSPECT BAG PRIOR TO ACTIVATION.



DISCARD IF:

- More than one chamber is **WHITE**.
- Solution is **YELLOW**.
- Seals are already **BROKEN**.

• REMOVE BAG FROM OVERPOUCH.

• ACTIVATE BAG.

- Roll *tightly* from top of bag down toward ports.
- *Apply pressure* until both vertical seals break and entire contents are white.



• INSPECT BAG TO CONFIRM ACTIVATION.

An activated bag has both vertical seals broken from bends to ports and entire contents are white.

• IDENTIFY CORRECT PORT FOR ADDITIVES AND INFUSION.

After mixing, the final admixture in this bag provides:

TOTAL CALORIES	1820 kcal	*	
AMINO ACIDS	68 g	SODIUM	64 mEq
DEXTROSE	220 g	POTASSIUM	47 mEq
LIPIDS	80 g	MAGNESIUM	16 mEq
		ACETATE	78 mEq
		CHLORIDE	92 mEq
		SULFATE	16 mEq
		CALCIUM	8 mEq (4 mmol)
		PHOSPHOROUS	20 mmol

Contents of each 100 mL of the mixed injection:
Soybean Oil, USP 3.9 g, Dextrose Monohydrate, USP 10.8 g.
Essential Amino Acids: Lysine, USP (added as the hydrochloride salt) 263 mg, Phenylalanine, USP 231mg, Leucine, USP 231mg, Valine, USP 213 mg, Histidine, USP 199 mg, Threonine, USP 164 mg, Methionine, USP 164 mg, Isoleucine, USP 164 mg, Tryptophan, USP 55 mg.
Nonessential Amino Acids: Alanine, USP 467 mg, Arginine, USP 330 mg, Glycine, USP 231 mg, Proline, USP 199 mg, Glutamic Acid 164 mg, Serine, USP 131 mg, Aspartic Acid, USP 99 mg, Tyrosine, USP 6.7 mg. **Electrolytes:** Sodium Acetate Trihydrate, USP 239 mg, Potassium Chloride, USP 174 mg, Sodium Glycero-phosphate Anhydrous 147 mg, Magnesium Sulfate Heptahydrate, USP 96 mg, Calcium Chloride Dihydrate, USP 29 mg. Also Contains: Purified egg phospholipids, glycerol, water for injection. Contains not more than 25 mcg/L of aluminum. pH adjusted with glacial acetic acid and sodium hydroxide.

NDC 63323-712-20

Sterile. Single dose container. For Dosage Information, see prescribing information.

For use only with calibrated infusion device. Discontinue infusion if adverse reaction occurs.

Recommended Storage: Store at 5°C to 25°C (41°F to 77°F). Avoid excessive heat. Protect from freezing. If accidentally frozen, discard bag.

Refrigerate activated bag at 2° to 8°C (36° to 46°F) for up to 7 days.

After removal from refrigerated storage, the activated bag should be infused within 48 hours. Discard unused portion.

339 893

2000

1500

1000

750

500

**BREAK VERTICAL SEALS FROM
HERE DOWN TO PORTS**

Additive compatibility, consult pharmacist.
Use Aseptic Technique.



ADDITIVE PORT

402550D



INFUSION PORT

NEVER USE THIS PORT

