

Batch no:
Manuf. date:
Use before:

Sample Label.
Please see package insert
for complete prescribing information.

Manufactured by:

**FRESENIUS
KABI**
Uppsala, Sweden
www.freseniuskabinutrition.com

KABIVEN®

**Amino Acids, Electrolytes, Dextrose, and Lipid Injectable Emulsion,
for intravenous use**

(3.3%, 0.7%*, 10.8% and 3.9%), No sulfites added CENTRAL INFUSION

2566 mL



**TO ACTIVATE BAG: WITH PORTS POINTING AWAY FROM YOU, ROLL *TIGHTLY* FROM TOP OF BAG
DOWN TOWARD PORTS UNTIL BOTH VERTICAL SEALS BREAK AND ENTIRE CONTENTS ARE WHITE.**



R_x only

SEE INSTRUCTIONS FOR USE



READ THIS



• **INSPECT BAG PRIOR TO ACTIVATION.
DISCARD IF:**

- More than one chamber is **WHITE**.
- Solution is **YELLOW**.
- Seals are already **BROKEN**.

• **REMOVE BAG FROM OVERPOUCH.**

• **ACTIVATE BAG.**

- Roll ***tightly*** from top of bag down toward ports.
- ***Apply pressure*** until both vertical seals break and entire contents are white.



• **INSPECT BAG TO CONFIRM
ACTIVATION.**

An activated bag has both vertical seals broken from bends to ports and entire contents are white.

• **IDENTIFY CORRECT PORT FOR
ADDITIVES AND INFUSION.**



NEVER USE THIS PORT



**BREAK VERTICAL SEALS FROM HERE
DOWN TO PORTS**

Additive compatibility, consult pharmacist.
Use Aseptic Technique.



ADDITIVE PORT

NDC 63323-712-25

Sterile. Single dose container. For Dosage Information, see prescribing information.

For use only with calibrated infusion device. Discontinue infusion if adverse reaction occurs.

Recommended Storage: Store at 5°C to 25°C (41°F to 77°F). Avoid excessive heat. Protect from freezing. If accidentally frozen, discard bag.

Refrigerate activated bag at 2° to 8°C (36° to 46°F) for up to 7 days.

After removal from refrigerated storage, the activated bag should be infused within 48 hours. Discard unused portion.

2500

2000

1500

1000

750

500

339 894

402551D



INFUSION PORT