

FRESENIUS
KABI



Contents: 10 x 50 mg Single Dose Vials

Rx only

For intravenous infusion only.
Discard unused portion.

50 mg per vial

Micafungin for Injection

NDC 63323-728-10

NDC 63323-728-10

Micafungin for Injection

50 mg per vial

For intravenous infusion only.
Discard unused portion.

Contents: 10 x 50 mg Single Dose Vials

Sterile.

Each vial contains: 50 mg micafungin equivalent to 50.86 mg micafungin sodium; 200 mg lactose; with citric acid and/or sodium hydroxide (used for pH adjustment).

Reconstitution: Reconstitute by aseptically adding 5 mL of 0.9% sodium chloride injection, USP (**without a bacteriostatic agent**) or 5 mL of 5% dextrose injection, USP and gently swirling the vial so that each mL contains 10 mg of micafungin. Visually inspect the vial for particulate matter. The reconstituted product may be stored in the original vial for up to 24 hours at room temperature, 25°C (77°F).

Requires further dilution prior to infusion.

For dosage, reconstitution and dilution, **see package insert.**

STORE AT: 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light.

Preservative free.

Discard unused portion.



(01)20363323728100

FRESENIUS
KABI

PSCAR-009953-01

NDC 63323-728-10

Micafungin for Injection

50 mg per vial



(01)20363323728100

www.fresenius-kabi.com/us

NDC 63323-728-10

Micafungin for Injection

50 mg per vial

For intravenous infusion only.
Discard unused portion.

Rx only

Contents: 10 x 50 mg Single Dose Vials

FRESENIUS
KABI

**Sample Label. Please see
package insert for complete
prescribing information.**

Mfd. for:

FRESENIUS
KABI
Lake Zurich, IL 60047
Made in India
Code No.: AP/DRUGS/103/97

621407A

For product inquiry: 1-800-551-7176 or www.fresenius-kabi.com/us



(01)20363323728100