



Rx only
Contents: **10 x 100 mg** Single Dose Vials

For intravenous infusion only.
Discard unused portion.

100 mg per vial

Micafungin
for Injection

NDC 63323-729-10

NDC 63323-729-10

Micafungin
for Injection

100 mg per vial

For intravenous infusion only.
Discard unused portion.

Contents: **10 x 100 mg** Single Dose Vials



Sterile.

Each vial contains: 100 mg micafungin equivalent to 101.73 mg micafungin sodium; 200 mg lactose; with citric acid and/or sodium hydroxide (used for pH adjustment).

Reconstitution: Reconstitute by aseptically adding 5 mL of 0.9% sodium chloride injection, USP (**without a bacteriostatic agent**) or 5 mL of 5% dextrose injection, USP and gently swirling the vial so that each mL contains 20 mg of micafungin. Visually inspect the vial for particulate matter. The reconstituted product may be stored in the original vial for up to 24 hours at room temperature, 25°C (77°F).

Requires further dilution prior to infusion.

For dosage, reconstitution and dilution, **see package insert.**

STORE AT: 25°C (77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Protect from light.

Preservative free.

Discard unused portion.

1312000579-02

NDC 63323-729-10

Micafungin
for Injection

100 mg per vial

www.fresenius-kabi.com/us

NDC 63323-729-10

Micafungin
for Injection

100 mg per vial

For intravenous infusion only.
Discard unused portion.

Rx only

Contents: **10 x 100 mg** Single Dose Vials



**Sample label. Please see
package insert for complete
prescribing information.**

Mfd. for:

 **FRESENIUS
KABI**
Lake Zurich, IL 60047
Made in India
Code No.: AP/DRUGS/103/97

6 2 1 4 0 8 B

For product inquiry: 1-800-551-7176 or www.fresenius-kabi.com/us



(01)20363323729107