

10089561Z59E00 N119

PSCAR-010511-00

Sample Label.
Please see
package insert
for complete
prescribing
information.

NDC 65219-**568**-00 Rx only

Mitomycin for Injection, USP

40 mg/vial

**WARNING: MUST BE
ADMINISTERED INTRAVENOUSLY
TO AVOID TISSUE DAMAGE.**

Caution: Cytotoxic Agent

Single-Dose Vial



This vial contains mitomycin 40 mg and mannitol 80 mg. Add 80 mL Sterile Water for Injection, USP. Shake to dissolve. If product does not dissolve immediately, allow to stand at room temperature until solution is obtained. Each mL of reconstituted solution will contain 0.5 mg mitomycin and have a pH between 6.0 and 8.0.

USUAL DOSAGE:

See package insert for full prescribing information.

STORAGE:

Retain the container in the carton. Store dry powder at 25°C, excursion permitted between 15°C and 30°C, protected from light. Avoid excessive heat, over 40°C (104°F). Protect reconstituted solution from light. Store solution under refrigeration 2° to 8°C (36° to 46°F), discard after 14 days. If unrefrigerated, discard after 7 days.

NDC 65219-**568**-00 Rx only

Mitomycin for Injection, USP

40 mg/vial

**WARNING: MUST BE
ADMINISTERED INTRAVENOUSLY
TO AVOID TISSUE DAMAGE.**

Caution: Cytotoxic Agent

Single-Dose Vial



Manufactured for:



Lake Zurich, IL 60047

Made in India

621864

Code No.: AP/DRUGS/03/2011



(01)00365219568001

www.fresenius-kabi.com/us