

Sample label. Please see package insert
for complete prescribing information.

855040/04

Nafcillin
for Injection, USP

2 grams per vial

Buffered - for intravenous or
intramuscular use.

621001C



10 x 2 grams
Vials

Made in Italy

Mfd. for:
**FRESENIUS
KABI**
Lake Zurich, IL 60047
by 181

Sterile.
When reconstituted with 6.6 mL diluent (SEE INSERT - INTRAMUSCULAR ROUTE), each vial contains 8 mL solution. Each mL of solution contains 250 mg nafcillin. The sodium content is 65.8 mg [2.9 mEq] per gram of nafcillin. The product is buffered with 40 mg sodium citrate per gram. Read accompanying insert for complete stability data.
Usual adult dosage: 500 mg every 4 to 6 hours. Read accompanying insert for directions for IM or IV use.
Before reconstitution store sterile powder at **20° to 25°C (68° to 77°F)** [see USP Controlled Room Temperature]. The container closure is not made with natural rubber latex.

www.fresenius-kabi.us

NDC 63323-328-20

Nafcillin
for Injection, USP

2 grams per vial

Buffered - for intravenous or intramuscular use.

10 x 2 grams
Vials



**FRESENIUS
KABI**

NDC 63323-328-20

Nafcillin
for Injection, USP

2 grams per vial

Buffered - for intravenous or intramuscular use.

Each vial contains nafcillin sodium, as the monohydrate equivalent to
2 grams of nafcillin.

**FRESENIUS
KABI**

Rx only