

GTIN: 00363323378059

[www.fresenius-kabi.com/us](http://www.fresenius-kabi.com/us)

For Subcutaneous or Intravenous Use

1,000 mcg per 5 mL (200 mcg per mL)

Octreotide Acetate Injection

NDC 63323-378-05

## Octreotide Acetate Injection

1,000 mcg per 5 mL  
(200 mcg per mL)



3 63323-378-05 9

 **FRESENIUS  
KABI**  
Lake Zurich, IL 60047  
[www.fresenius-kabi.com/us](http://www.fresenius-kabi.com/us)

NDC 63323-378-05

## Octreotide Acetate Injection

1,000 mcg per 5 mL  
(200 mcg per mL)

Sterile.

**Each mL contains:** Octreotide (as acetate), 200 mcg; sodium chloride, 7 mg; glacial acetic acid USP, 2 mg; sodium acetate trihydrate USP, 2 mg; phenol USP, 5 mg; water for injection, q.s.

**Usual dosage:** See package insert.

5 mL  
Multiple Dose Vial

 **FRESENIUS  
KABI**

NDC 63323-378-05

## Octreotide Acetate Injection

1,000 mcg per 5 mL  
(200 mcg per mL)

For Subcutaneous or Intravenous Use

**STORE REFRIGERATED AT 2°C TO 8°C  
(36°F TO 46°F); PROTECT FROM  
LIGHT. AFTER INITIAL USE, DISCARD  
WITHIN 14 DAYS.**

5 mL  
Multiple Dose Vial  
Rx only

 **FRESENIUS  
KABI**

NDC 63323-378-05

## Octreotide Acetate Injection

1,000 mcg per 5 mL  
(200 mcg per mL)

**Storage:** Refrigerate at 2°C to 8°C (36°F to 46°F); protect from light. After initial use, discard within 14 days. At room temperature, (20°C to 30°C or 70°F to 86°F), Octreotide Acetate is stable for 14 days if protected from light. The container closure is not made with natural rubber latex.

5 mL  
Multiple Dose Vial

 **FRESENIUS  
KABI**

62850G