



621172B

Sample Label. Please see package insert
for complete prescribing information.

NDC 63323-134-10

Pemetrexed for Injection, USP

100 mg per vial

For intravenous use only
Caution: Cytotoxic Agent

134010

NDC 63323-134-10

134010

Pemetrexed for Injection, USP

100 mg per vial

For intravenous use only
Caution: Cytotoxic Agent
Usual Dosage: See insert.

The container closure is not made with natural rubber latex.

Storage: Store powder at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Store reconstituted and infusion solutions at refrigerated 2° to 8°C (36° to 46°F), or 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Single-Dose Vial
Rx onlySingle-Dose Vial
Rx only

Mfd. by:
**FRESENIUS
KABI**
Lake Zurich, IL 60047

www.fresenius-kabi.com/us

GTIN: 00363323134105

Pemetrexed
for Injection, USP
100 mg per vial



3 63323-134-10 5

Pemetrexed for Injection, USP

100 mg per vial

For intravenous use only
Caution: Cytotoxic Agent

Each vial contains: pemetrexed disodium equivalent to 100 mg pemetrexed, 106 mg of mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.

To Reconstitute: Add 4.2 mL of 0.9% Sodium Chloride Injection (preservative free) to make a solution containing 25 mg/mL pemetrexed. Reconstituted solution must be further diluted before use (see accompanying literature). Administer infusion solution within 24 hours after initial reconstitution. Discard unused portion.

Single-Dose Vial
Rx only

621172B