

Sample Label. Please see package insert for complete prescribing information.

NDC 63323-450-50

134150

Pemetrexed for Injection, USP

500 mg per vial

For intravenous use only
Caution: Cytotoxic Agent

Single-Dose Vial
Rx only



NDC 63323-450-50 134150

Pemetrexed for Injection, USP

500 mg per vial

For intravenous use only
Caution: Cytotoxic Agent

Usual Dosage: See insert.

The container closure is not made with natural rubber latex.

Storage: Store powder at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Store reconstituted and infusion solutions at refrigerated 2° to 8°C (36° to 46°F), or 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Single-Dose Vial
Rx only



Pemetrexed for Injection, USP

500 mg per vial

For intravenous use only
Caution: Cytotoxic Agent



3 63323-450-50 2

Mfd. by:
 FRESENIUS
KABI
Lake Zurich, IL 60047

www.fresenius-kabi.com/us

621173B

Pemetrexed for Injection, USP

500 mg per vial

For intravenous use only
Caution: Cytotoxic Agent

Each vial contains: pemetrexed disodium equivalent to 500 mg pemetrexed, 500 mg of mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.

To Reconstitute: Add 20 mL of 0.9% Sodium Chloride Injection (preservative free) to make a solution containing 25 mg/mL pemetrexed. Reconstituted solution must be further diluted before use (see accompanying literature).

Administer infusion solution within 24 hours after initial reconstitution. Discard unused portion.

Single-Dose Vial
Rx only

