

NDC 63323-113-10

11310

PENTAM® 300

*(pentamidine isethionate
for injection)*

300 mg

Lyophilized

**For Intramuscular or
Intravenous Use**

Rx only

10 Single Dose Vials

Preservative Free

Sterile, Nonpyrogenic

Each vial contains: Pentamidine Isethionate
300 mg.

Dosage: Recommended regimen for adults
and pediatric patients is 4 mg/kg once a day
for 14 days. The benefits and risks of therapy
with pentamidine isethionate for more than
14 days are not well defined.

ADMINISTRATION: Intramuscular Injection:

The contents of 1 vial (300 mg) should
be dissolved in 3 mL of Sterile Water for
Injection, USP. The calculated daily dose
should then be withdrawn and administered
by deep intramuscular injection.

Intravenous Injection: The contents of 1 vial
should first be dissolved in 3-5 mL of Sterile
Water for Injection, USP or 5% Dextrose
Injection, USP only. The calculated dose
of pentamidine isethionate should then be

Sample label.

Please see package insert for complete prescribing information.

withdrawn and diluted further in 50-250 mL
of 5% dextrose solution. **The diluted
intravenous solutions containing
pentamidine isethionate should be
infused over a period of 60 minutes.**

For additional information: See Package
Insert.

Store dry product at 20° to 25°C (68°
to 77°F) [see USP Controlled Room
Temperature]. Protect dry product and
reconstituted solution from light.



**FRESENIUS
KABI**

Lake Zurich, IL 60047

www.fresenius-kabi.com/us

42956A



(01)20363323113104