

Batch no:  
Manuf. date:  
Use before:

Sample Label.  
Please see package insert  
for complete prescribing information.

Manufactured by:

**FRESENIUS  
KABI**  
Uppsala, Sweden  
www.freseniuskabinutrition.com

**PERIKABIVEN®**

Amino Acids, Electrolytes, Dextrose, and Lipid Injectable Emulsion,  
for intravenous use

(2.4%, 0.5%\*, 7.5% and 3.5%), No sulfites added

**PERIPHERAL INFUSION**

**1440 mL**



TO ACTIVATE BAG: WITH PORTS POINTING AWAY FROM YOU, ROLL *TIGHTLY* FROM TOP OF BAG  
DOWN TOWARD PORTS UNTIL BOTH VERTICAL SEALS BREAK AND ENTIRE CONTENTS ARE WHITE.



R<sub>x</sub> only

SEE INSTRUCTIONS FOR USE

**STOP READ THIS**



• **INSPECT BAG PRIOR TO ACTIVATION.**  
**DISCARD IF:**

- More than one chamber is **WHITE**.
- Solution is **YELLOW**.
- Seals are already **BROKEN**.

• **REMOVE BAG FROM OVERPOUCH.**

• **ACTIVATE BAG.**

- Roll *tightly* from top of bag down toward ports.
- *Apply pressure* until both vertical seals break and entire contents are white.



• **INSPECT BAG TO CONFIRM ACTIVATION.**

An activated bag has both vertical seals broken from bends to ports and entire contents are white.

• **IDENTIFY CORRECT PORT FOR ADDITIVES AND INFUSION.**

**NEVER USE THIS PORT**



After mixing, the final admixture in this bag provides:

TOTAL CALORIES	1010 kcal	*
AMINO ACIDS	34 g	SODIUM 32 mEq
DEXTROSE	107 g	POTASSIUM 24 mEq
LIPIDS	51 g	MAGNESIUM 8 mEq
		ACETATE 39 mEq
		CHLORIDE 46 mEq
		SULFATE 8 mEq
		CALCIUM 4 mEq (2 mmol)
		PHOSPHOROUS 11 mmol

Osmolarity: 750 mOsmol/L;

pH 5.6

Contents of each 100 mL of the mixed injection:  
Soybean Oil, USP 3.5 g, Dextrose Monohydrate, USP 7.5 g.  
**Essential Amino Acids:** Lysine, USP (added as the hydrochloride salt) 187 mg, Phenylalanine, USP 164 mg, Leucine, USP 164 mg, Valine, USP 152 mg, Histidine, USP 141 mg, Threonine, USP 116 mg, Methionine, USP 116 mg, Isoleucine, USP 116 mg, Tryptophan, USP 40 mg. **Nonessential Amino Acids:** Alanine, USP 333 mg, Arginine, USP 235 mg, Glycine, USP 164 mg, Proline, USP 141 mg, Glutamic Acid 116 mg, Serine, USP 94 mg, Aspartic Acid, USP 71 mg, Tyrosine, USP 4.8 mg. **Electrolytes:** Sodium Acetate Trihydrate, USP 170 mg, Potassium Chloride, USP 124 mg, Sodium Glycero-phosphate Anhydrous 105 mg, Magnesium Sulfate Heptahydrate, USP 68 mg, Calcium Chloride Dihydrate, USP 20 mg.

Also Contains: Purified egg phospholipids, glycerol, water for injection. Contains not more than 25 mcg/L of aluminum. pH adjusted with glacial acetic acid and sodium hydroxide.

**BREAK VERTICAL SEALS FROM  
HERE DOWN TO PORTS**

Additive compatibility, consult pharmacist.  
Use Aseptic Technique.



**ADDITIVE PORT**

NDC 63323-714-14

Sterile. Single dose container. For Dosage Information, see prescribing information.

For use only with calibrated infusion device. Discontinue infusion if adverse reaction occurs.

Recommended Storage: Store at 5°C to 25°C (41°F to 77°F). Avoid excessive heat. Protect from freezing. If accidentally frozen, discard bag.

Refrigerated storage after mixing limited to 24 hours. Discard unused portion.

402552C



1000

750

500

250



**INFUSION PORT**

339 839