

Batch no:  
Manuf. date:  
Use before:

PERIKABIVEN®

Amino Acids, Electrolytes, Dextrose, and Lipid Injectable Emulsion,  
for intravenous use

(2.4%, 0.5%\*, 7.5% and 3.5%), No sulfites added

1920 mL

PERIPHERAL INFUSION

3 63323 71419 2

ROLL

TO ACTIVATE BAG: WITH PORTS POINTING AWAY FROM YOU, ROLL *TIGHTLY* FROM TOP OF BAG  
DOWN TOWARD PORTS UNTIL BOTH VERTICAL SEALS BREAK AND ENTIRE CONTENTS ARE WHITE.

ROLL

R<sub>x</sub> only

SEE INSTRUCTIONS FOR USE

STOP

READ THIS

• INSPECT BAG PRIOR TO ACTIVATION.  
DISCARD IF:  
- More than one chamber is **WHITE**.  
- Solution is **YELLOW**.  
- Seals are already **BROKEN**.

• REMOVE BAG FROM OVERPOUCH.

• ACTIVATE BAG.  
- Roll *tightly*  
from top of  
bag down  
toward ports.  
- *Apply pressure*  
until both  
vertical seals  
break and  
entire contents  
are white.

• INSPECT BAG TO CONFIRM  
ACTIVATION.  
An activated bag has both vertical  
seals broken from bends to ports  
and entire contents are white.

• IDENTIFY CORRECT PORT FOR  
ADDITIVES AND INFUSION.

↑

↑

↑

↑

↑

↑

↑

↑

ROLL

NEVER USE THIS PORT

After mixing, the final admixture in this bag provides:

TOTAL CALORIES	1346 kcal	*
AMINO ACIDS	45 g	SODIUM 42 mEq
DEXTROSE	143 g	POTASSIUM 33 mEq
LIPIDS	68 g	MAGNESIUM 11 mEq
		ACETATE 52 mEq
		CHLORIDE 61 mEq
		SULFATE 11 mEq
Osmolarity: 750 mOsmol/L;		CALCIUM 5 mEq (3 mmol)
pH 5.6		PHOSPHOROUS 14 mmol

Contents of each 100 mL of the mixed injection:  
Soybean Oil, USP 3.5 g, Dextrose Monohydrate, USP 7.5 g.  
**Essential Amino Acids:** Lysine, USP (added as the hydrochloride salt) 187 mg, Phenylalanine, USP 164 mg, Leucine, USP 164 mg, Valine, USP 152 mg, Histidine, USP 141 mg, Threonine, USP 116 mg, Methionine, USP 116 mg, Isoleucine, USP 116 mg, Tryptophan, USP 40 mg. **Nonessential Amino Acids:** Alanine, USP 333 mg, Arginine, USP 235 mg, Glycine, USP 164 mg, Proline, USP 141 mg, Glutamic Acid 116 mg, Serine, USP 94 mg, Aspartic Acid, USP 71 mg, Tyrosine, USP 4.8 mg. **Electrolytes:** Sodium Acetate Trihydrate, USP 170 mg, Potassium Chloride, USP 124 mg, Sodium Glycero-phosphate Anhydrous 105 mg, Magnesium Sulfate Heptahydrate, USP 68 mg, Calcium Chloride Dihydrate, USP 20 mg.  
Also Contains: Purified egg phospholipids, glycerol, water for injection. Contains not more than 25 mcg/L of aluminum. pH adjusted with glacial acetic acid and sodium hydroxide.

↓ BREAK VERTICAL SEALS FROM  
HERE DOWN TO PORTS ↓

Additive compatibility, consult pharmacist.  
Use Aseptic Technique.

ADDITIVE PORT

NDC 63323-714-19

Sterile. Single dose container. For Dosage  
Information, see prescribing information.

For use only with calibrated infusion device.  
Discontinue infusion if adverse reaction  
occurs.

Recommended Storage: Store at 5°C to 25°C  
(41°F to 77°F). Avoid excessive heat. Protect  
from freezing. If accidentally frozen, discard  
bag.

Refrigerated storage after mixing limited to  
24 hours. Discard unused portion.

402553C

LATEX

DEHP

05

PP

ROLL

INFUSION PORT

339 840

1500

1000

750

500