

NDC 65219-512-50

Contains 25 of NDC 65219-512-02

Potassium Acetate Injection, USP

100 mEq/50 mL (2 mEq/mL)

Pharmacy Bulk Package—Not for Direct Infusion

**CAUTION: MUST BE DILUTED.
FOR INTRAVENOUS USE.***

50 mL
Rx only

Each mL contains potassium acetate, 196 mg. May contain acetic acid for pH adjustment. 4 mOsmol/mL (calc). pH 6.2 (5.5 to 8.0). **Usual dosage:** See insert. Sterile, nonpyrogenic.

Aseptically add to a suitable intravenous solution. Contains no bacteriostat; use promptly; discard unused portion. Use only if clear and seal is intact and undamaged.

*For preparing intravenous admixtures only. See insert for complete dosage information and proper use of this container. Discard vial 4 hours after initial entry. A single entry through the vial closure should be made with a sterile dispensing set or transfer device. Transfer individual doses to appropriate intravenous infusion solutions. Use of a syringe with needle is not recommended. The above process should be carried out under a laminar flow hood using aseptic technique.

Use Aseptic Technique

Remove cover from fliptop vial and cleanse stopper with antiseptic. Add to a suitable solution in an intravenous container.

Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]



**FRESENIUS
KABI**

Lake Zurich IL, 60047

Sample label. Please see package insert
for complete prescribing information.

www.fresenius-kabi.com/us

421086



(01)20365219512503